

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

ZOLIAR[®] 800 DF

HERBICIDE

ACTIVE CONSTITUENT: 800 g/kg NORFLURAZON



A pre-emergence selective residual herbicide for a programmed approach to Nutgrass (Cyperus spp) and other grass and broadleaf weed control in Asparagus, Citrus, Grapes, Nuts, Pome Fruit and Stone Fruit as per the Directions for Use.

***The bag must not be sold separately.
DO NOT destroy box while product still remains.***

***In a transport emergency dial 000, Police or Fire Brigade
For specialist advice in an emergency only, call 1800 033 111 (24 hours)***

AgNova Technologies Pty Ltd
ABN 70 097 705 158
Suite 3, 935 Station Street
Box Hill North, Vic. 3129 Australia
Phone (03) 9899 8100
agnova.com.au



GENERAL INSTRUCTIONS

Asparagus: To avoid crop damage it is advisable to only treat a small number of plants to ascertain their reaction before treating the whole crop.

Citrus, grapes, apples, pears, stone fruit and almond orchards: Zoliar 800 DF is a surface applied residual pre-emergence herbicide. Residual weed control can be expected for up to 6 months, dependent on rate applied and soil conditions. The 2.5 kg/ha rate is most effective on coarse or light loamy soils when harder to control weeds are not important. Extended period of weed control can be achieved by using rates higher than 2.5 kg. The period of control will then be determined by soil type, irrigation/rainfall patterns, time of treatment and other soil and management factors.

Rainfall or irrigation within 2 or 3 weeks after application is necessary for activation.

Areas to be treated should be free of weeds and excessive mulch. Where weeds are already growing in the area to be treated, an appropriate "knockdown herbicide" must be applied either before or, if compatible, combined as a tank mix with this product. This product should be applied strategically at mid spring for full summer weed control, and early autumn when winter control is required. Where strip treatment is applied, the recommended rate per hectare must relate only to the area of the treated strip. If residual or "knockdown herbicides" are added, consider all label recommendations and adhere to the crop recommendations and directions that are most restrictive.

Mixing

Fill the spray or mixing tank to 2/3 full and start agitation. Add required amount of ZOLIAR 800 DF to tank whilst maintaining continuous agitation. Add additional compatible products (if required), then balance of water until tank is full. Maintain agitation throughout mixing, standing and spraying operations.

If tank mixing, products should be added in the following order, after ZOLIAR 800 DF:

1. Wettable powders and water dispersible granules
2. Flowable liquids
3. Emulsifiable concentrates
4. Surfactants

Application

Citrus, grapes, apples, pears, stone fruit and almond orchards: Apply ZOLIAR 800 DF as a directed spray, avoiding contact with foliage or fruit using a fixed boom sprayer. Apply 300 to 500 L of spray/ha, have good tank agitation, use 50 mesh nozzle strainers. Shut off spray booms when starting, stopping or slowing down to avoid overdosing. Wash out equipment thoroughly with water after use.

Compatibility

ZOLIAR 800 DF is compatible with most "knockdown" and residual herbicides in situations where appropriate recommendations exist for such herbicides in the crop concerned. Tank mixtures with residual herbicides are primarily made to improve control of particular weed species. Observe mixing instructions.

DIRECTIONS FOR USE

Restraints:

- DO NOT rotate to alternate crops in fields treated with ZOLIAR 800 DF
- DO NOT incorporate in a manner which would place untreated soil on top of treated soil.
- DO NOT apply to nursery stock.
- DO NOT use more than 4.2 kg/ha of ZOLIAR 800DF in any one year.
- DO NOT apply to trees or vines unless they have been established in the orchard or vineyard for 18 months.

Crop	Weed	State	Rate /ha	WHP	Critical Comments
Asparagus	Johnson Grass, Nutgrass	Qld	2.5 kg	14 days	Apply using a boom spray to bare ground prior to weed emergence. Apply as a directed spray in 300 to 500 L water/hectare. Avoid contact with foliage or fruit. An application in early autumn will give winter weed control or in early to mid-spring will give full summer weed control.
Citrus Grapefruit, Lemons, Mandarins, Oranges (Navel and Valencia) Grapes Nuts Almonds only Pome Fruit Apples, Pears Stone Fruit Apricots, Cherries, Nectarines, Peaches, Plums	Annual Ryegrass, Barley Grass, Blackberry Nightshade, Brachiaria (Green Summer Grass), Caltrop, Capeweed, Chickweed, Common Sowthistle (Milk Thistle), Dandelion seedlings, Curled Dock seedlings, False Caper seedlings, Fat-Hen ⁺ , Indian Hedge Mustard, Innocent Weed (Spiny Burgrass), Medic, Hedge Mustards, Paspalum, Plantain seedlings, Pigweed (<i>Portulaca</i>), Prairie Grass, Prickly Lettuce, Great Brome (Ripgut Brome), Salvation Jane, Scarlet Pimpernel, Shepherd's Purse, Silver Grass, Skeleton Weed seedlings, Sorrel seedlings, Soursob ⁺ , Stinkgrass, Stinking Roger, Subterranean Clover, Summer Grass (Crabgrass), Three-cornered Jack (Doublegee, Spiny Emex) Variegated Thistle, Wild Oats, Wild Radish, Wild Turnip, Winter Grass, Wireweed, Witch Grass, Yellow Weed, Yorkshire Fog Grass Couch Grass ⁺ , Dandelion, Curled Dock, False Caper, Johnson Grass ⁺ , Skeleton Weed, Sorrel, Soursob	All states	5.0 kg		Note: <i>For asparagus only:</i> Allow newly planted fields (direct seeded, seedlings, or crowns) to become established for one year before application of ZOLIAR 800 DF. DO NOT apply within 14 days of harvest and before fern emerges. DO NOT spray on young ferns. DO NOT apply more than 2 applications per year. ZOLIAR 800 DF should not be applied if crop rotation is expected within 24 months. <i>For grapes only:</i> Not recommended for grapes grown in sand or loamy sand soils with less than 1% organic matter and pH greater than 7.5 as veinal chlorosis may occur. <i>For almonds and cherries only:</i> DO NOT use on coarse textured soil such as sand, loamy sand, gravelly sand and fine sand as veinal chlorosis may occur.

⁺ Suppression only

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

Asparagus: DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION

Other Crops: NOT REQUIRED WHEN USED AS DIRECTED.

RESISTANT WEEDS WARNING **GROUP F HERBICIDE**

ZOLIAR 800 DF Herbicide is a member of the pyridazinone group of herbicides. ZOLIAR 800 DF has the inhibitor of carotenoid biosynthesis mode of action. For weed resistance management ZOLIAR 800 DF is a Group F herbicide. Some naturally-occurring weed biotypes resistant to ZOLIAR 800 DF and other inhibitors of carotenoid biosynthesis herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by ZOLIAR 800 DF or other Group F herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, AgNova Technologies Pty Ltd accepts no liability for any losses that may result from the failure of ZOLIAR 800 DF to control resistant weeds.

PROTECTION OF LIVESTOCK

DO NOT graze livestock on treated areas.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

DO NOT drain or flush equipment or spray near desirable trees or other plants or where their roots may extend; or in locations where the product may be washed or moved into such situations.

DO NOT apply while fruit or nuts are on the ground at harvest.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with ZOLIAR 800 DF or used container. Avoid rapid flood irrigation which may erode treated soil from the field.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, well ventilated area, as cool as possible, out of direct sunlight. Keep out of reach of children. Single rinse containers before disposal. Add rinsings to the spray tank. DO NOT dispose of undiluted chemicals on site. Puncture and bury the containers in a local authority landfill. If not available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Avoid contact with skin and eyes. DO NOT inhale dust or spray mist. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 131 126.

SAFETY DATA SHEET

If additional hazard information is required refer to the Material Safety Data Sheet. For a copy visit our website at agnova.com.au

CONDITIONS OF SALE

AgNova Technologies Pty Ltd shall not be liable for any consequential or other loss or damage relating to the supply or subsequent handling or use of this product, unless such liability by law cannot be lawfully excluded or limited. All warranties, conditions or rights implied by statute or other law which may be lawfully excluded are so excluded. Where the liability of AgNova Technologies Pty Ltd for breach of any such statutory warranties and conditions cannot be lawfully excluded but may be limited to it re-supplying the product or an equivalent product or the cost of a product or an equivalent product, then the liability of AgNova Technologies Pty Ltd for any breach of such statutory warranty or condition is so limited.

® Zoliar is a registered trademark of Tessenderlo Kerley Inc.



Batch Number	
Date of Manufacture	