



**DANGEROUS POISON**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING  
CAN KILL IF SWALLOWED  
DO NOT PUT IN DRINK BOTTLES  
KEEP LOCKED UP



**PARAQUAT 250**

HERBICIDE

ACTIVE CONSTITUENT: 250g/L PARAQUAT PRESENT AS PARAQUAT DICHLORIDE

GROUP L HERBICIDE

FOR THE CONTROL OF A WIDE RANGE OF GRASSES  
AND BROADLEAF WEEDS AS PER DIRECTIONS FOR USE.

READ THE ATTACHED BOOKLET BEFORE USING THIS PRODUCT.

CONTENTS: 5, 20, 100, 110 and 200 LITRES

**4FARMERS PTY. LTD.**

A.C.N. 067 443 485

D.O.M

BATCH NO:

APVMA APPROVAL No:



**FOR USE ONLY AS AN AGRICULTURAL AND HORTICULTURAL HERBICIDE. THIS PRODUCT IS TO HAZARDOUS TO BE USED IN THE HOME GARDEN.**

**STORAGE AND DISPOSAL**

Store in the closed, original container in a dry, cool, well-ventilated locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. Do not store for prolonged periods in direct sunlight.

Do not repack or use container for any other purpose. Triple or preferably pressure rinse empty containers adding the rinsate to the spray tank. Do not dispose undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury containers in a local authority landfill. If not available bury the containers below 500mm in a disposal pit specifically set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

**For refillable containers:**

Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

**SAFETY DIRECTIONS**

Very dangerous, particularly the concentrate

Product is poisonous if swallowed. Will irritate the nose, throat and skin.

Attacks eyes, protect eyes while using. Avoid contact with eyes, skin and clothing.

When opening the container and preparing the product for use, wear elbow length PVC gloves and face shield or goggles. If product on skin, immediately wash area with soap and water. If clothing becomes contaminated with product, remove clothing immediately. If product in eyes, wash it out immediately with water. Avoid contact with spray mist. Do not inhale spray mist. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash contaminated clothing, gloves and face shield or goggles.

**Spray Application:**

DO NOT work in spray mist.

DO NOT continue to use if skin irritation or nosebleed occurs. This may be caused by exposure to spray mist as the result of incorrect use of equipment or adverse climatic conditions. Stop and review handling and spraying techniques before further spraying. If symptoms persist seek medical advice.

Where there is a risk of exposure to spray mist wear waterproof footwear and waterproof protective clothing, impervious gauntlet length gloves (rubber or PVC), goggles and a face mask and respirator covering nose and mouth and capable of filtering spray droplets. A high efficiency type particulate respirator is recommended, but in any event use a respirator which complies with the requirements of AS1716 (Standards Association of Australia). Further advice on safety equipment should be obtained from a safety equipment manufacturer.

Avoid contacting vegetation wet with spray, but if necessary to do so wear waterproof footwear and waterproof protective clothing and gloves.

**FIRST AID:**

If poisoning occurs, get to a doctor or hospital quickly. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

**MATERIAL SAFETY DATA SHEET:**

Additional information is listed in the Material Safety Data Sheet.

**CONDITIONS OF SALE:**

The use of 4 Farmers Paraquat 250 Herbicide, being beyond the control of the manufacturer, no warranty expressed or implied is given by 4 Farmers Pty. Ltd. regarding its suitability, fitness or efficacy for any purpose for which it is used by the buyer, whether in accordance with the directions or not and 4 Farmers Pty. Ltd. accepts no responsibility for any consequences whatsoever resulting from the use of this product.

**4FARMERS PTY LTD**  
**A.C.N. 067 443 485**  
**643 Murray Street**  
**WEST PERTH WESTERN AUSTRALIA 6005**

**Phone: (08) 9356 3445**  
**Fax: (08) 9356 3447**

Toxic 6 diamond

UN No. 3016

Pesticides, Bipyridium Liquid toxic (contains paraquat).

Hazchem Code 2X

In an Emergency Dial 000 Police or Fire Brigade.



**DANGEROUS POISON**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING  
CAN KILL IF SWALLOWED  
DO NOT PUT IN DRINK BOTTLES  
KEEP LOCKED UP



**PARAQUAT 250**

**HERBICIDE**

ACTIVE CONSTITUENT: 250g/L PARAQUAT PRESENT AS PARAQUAT DICHLORIDE

GROUP **L** HERBICIDE

FOR THE CONTROL OF A WIDE RANGE OF GRASSES  
AND BROADLEAF WEEDS AS PER DIRECTIONS FOR USE.

**READ THE ENTIRE LABEL LEAFLET BEFORE USING THIS PRODUCT**

**4FARMERS PTY. LTD.**  
A.C.N. 067 443 485

APVMA APPROVAL No:



## DIRECTIONS FOR USE

### RESTRAINTS: FOR GROUND APPLICATION ONLY

DO NOT use through aircraft, misting machines (except banana plantations) or hand-held ultra low volume applicators (CDA units).

DO NOT add wetter unless spraying at high volume, use an appropriate wetter as per directions for use.

DO NOT spray plants covered with heavy dew, waterlogged, under stress of any kind or covered with soil or dust.

Note: Rain following spraying will not effect results.

DO NOT sow or cultivate for 1 hour after spraying, but operations should commence within 7 days.

CROP USE OR SITUATION	WEEDS CONTROLLED	STATE	RATE/HA	CRITICAL COMMENTS
PERENNIAL PASTURES AND CROPS  Mature lucerne stands - Autumn/early winter  - Late Winter/Spring  Mature Perennial Grass Seed Crops -Cocksfoot, Perennial Ryegrass Phalaris and Demeter Fescue only Kikuyu/Paspalum - Pasture to suppress growth to oversow Winter Seed	Annual grass and some broad leaf weeds	Qld, Vic, SA, WA, Tas only	1.2 to 1.6L	Do not spray lucerne stands under 12 months old. NOTE: In some instances <i>H. glaucum</i> has shown signs of resistance to paraquat. If resistance is suspected, other products such as 'Fusilade' should be used for grass weed control. Check with your local Department of Agriculture. Use the higher rates for dense weed stands. For residual weed control or if Paterson's Curse, Shepherd's Purse and some other broadleaf weeds are present, add a compatible Diuron formulation at registered rates.
		NSW only	1.2L	
		Qld, SA, WA, Vic, NT, Tas Only	1.6 to 2.4L	
		All States	600mL to 1.2L	
		Qld, NSW, ACT only	1.6 or 2.4L	Use the high rate for February spraying and use the low rate for March.
Bananas	Annual Weed Control	Qld, NSW only	Knapsack: 100mL/100L Mister: 8mL/1L plus A suitable wetter	Spray weeds from after emergence and prior to reach 15cm in height. Repeat sprays as required. To minimise drift, select a spraying pressure less than 240kPa. Chemical contact with roots and peepers near the pseudo stem will damage crops. Refer to directions for use with misting machines
Hops	Annual grasses	Vic, Tas only	1.2 to 1.6L	Spray prior to crop emerging from winter dormancy, using directed inter-row application. Ensure even coverage of weeds with herbicide. A minimum of 250L per ha spray volume is recommended.
Orchards and Vineyards	Annual grasses and some broad leaf weeds	Qld, Vic, SA, WA, NT, Tas only	1.6 to 3.2L	Spray when annual weeds are present. Avoid contact with crop foliage. 4Farmers Paraquat 250 herbicide will not damage mature brown bark, but avoid spraying green tissue of trees and vines. Use the higher rate for dense growth.
		NSW only	1.7L	



CROP USE OR SITUATION	WEEDS CONTROLLED	STATE	RATE/HA	CRITICAL COMMENTS
Pastures: Controlling grass seed set in medic and sub-clover pastures.  Grass pasture desiccation (Hay Freezing) – To preserve the value of standing feed - To minimise the risk of ryegrass toxicity	Barley Grass *( <i>Hordeum leporinum</i> ) *( <i>H glaucum</i> ) Annual ryegrass *( <i>Lolium rigidum</i> )	All States	400mL	Best results are obtained if seed heads of the target grass species develop evenly. Graze pastures heavily in spring. Remove stock 2-3 weeks before grass maturity. Barley grass: Spray when all seed heads have emerged but prior to the heads showing signs of haying off. Other grass species: Spray pasture when all seed heads of the target species have emerged and heads are showing initial signs of haying off. N.B. Ensure boom is raised to give a double overlap at the level of the seed heads. To aid coverage, spray at pressures of 250-350 kPa and use 110015 nozzles or equivalent. Check for hardened seeds prior to returning stock to paddocks at least 1 day after spraying. This treatment may scorch medic and subclover species. A reduction of legume seedset may occur if plants are flowering.
			800mL	Graze heavily to encourage even head development of grass species. Remove stock approximately 1 month before anticipated maturity. Spray prior to haying off of grasses. Ensure the boom is raised to give a double overlap at the level of the seed heads. This treatment will minimise the formation of toxin causing annual ryegrass toxicity. If ryegrass pasture is sprayed after first heads emerge but prior to commencement of flowering. Heavy grazing is then required to prevent regrowth and the development of new seed heads.
Control of saffron thistle	Saffron thistle *( <i>Carthamus fanatus</i> )	SA, NSW, ACT only	400mL	Spray thistles as they are beginning to “run up” or in the early branching stage, but before flowering, when green and actively growing. Ensure good coverage with herbicide.
		WA only	800mL	Spray when primary flower first emerges. Ensure double spray overlap.
Cultivation aid: To prepare a clean bed for sowing and minimise cultivation	Annual grasses and broadleaf weeds	Qld, Vic, SA, Tas, NSW, NT, ACT only	1.2 to 1.6L	When cultivation is to follow spray start 1 hour after spraying, but complete within 7 days. To achieve a better seedbed in areas of heavy weed growth at spraying, delay cultivation for 3-5 days. With dense more mature weed stands use the higher rate. Wild Oats must have at least two leaves. When used with Diquat, the lower rate of 4Farmers Paraquat 250 Herbicide should control dense mature weeds. Use the higher Diquat rate for plants more than 10cm in diameter.
	Early Autumn sowing			
	Winter, spring and early summer sowing	1.6 to 2.4L		
	Capeweed or Erodium	Qld, Vic, SA, Tas, NSW only	1.2L or 1.6L plus 750mL or 1.5L/ha of Diquat	To remove old pasture residues graze heavily then, 3-5 days before spraying remove stock to allow weeds to freshen up.
Rice	Annual grasses and broadleaf weeds	Qld, NSW only	1.6L	Pre-sowing.
			800mL	Post-sowing but before crop emergence.
Sugar Cane Up to 5cm high	Annual grasses and some broadleaf weeds	Qld, NSW, NT only	1.2 to 1.6L per sprayed ha	Apply as a broadcast spray over the top of plant cane up to the 3-4leaf stage or ratoon cane up to 10cm high. Cane foliage will be scorched but new leaves will appear in 7-10 days. In plant cane between the 3-4 leaf stage and the formation of the true stem use a directed interspace spray. The Irvin spray boom (or other similar equipment) is the most suitable equipment to avoid excessive drift onto cane foliage while spraying the cane bases of plant and ratoon cane. After the formation of the true stem, which is resistant to 4Farmers Paraquat 250 Herbicide, the spray height can be raised to overlap the spray pattern to give weed control in the stool. Use the higher rate for dense, maturer weeds. 4Farmers Paraquat 250 Herbicide can be mixed with atrazine to give residual weed control when used as a blanket or directed spray. Refer to the atrazine label for specific rates. It may also be mixed with diuron for residual weed control.
			1.2 to 1.6L per sprayed ha plus 275g to 500g diuron	



CROP USE OR SITUATION	WEEDS CONTROLLED	STATE	RATE/HA	CRITICAL COMMENTS
Sugar Cane Up to 10cm high	Annual grasses and some broadleaf weeds	Qld, NSW, NT only	1.2 to 1.6L per sprayed ha plus 1kg diuron	To enhance the activity of 4Farmers Paraquat 250 Herbicide under favourable growing conditions and in open sunny conditions, add diuron at rates shown for weed size. Diuron WG at rates up to 500g/ha can be blanket sprayed. Use a directed spray for higher rates of diuron. Complete spray coverage is essential. For grasses up to 5cm high use a minimum of 250L-spray solution/ha; increase to 350L/ha for weeds up to 10cm high. Use a minimum spray volume of 400L/ha for weeds > 10cm high and for dense mature weeds.  <b>Always add Agral at 200mL.</b>
	Annual grasses and some broadleaf weeds with short residual control.		1.6L plus 2.8 to 3.9kg diuron	
Spray topping to reduce seed set  Chickpeas Faba Beans Field Peas Lupins Vetch	Annual Ryegrass	WA, SA, NSW, ACT, VIC only	400mL or 800mL	As an aid in managing annual ryegrass resistance. For use on escapes from a previous herbicide application in the current crop.  Spray the crop when ryegrass seed heads at the bottom of the plant have emerged and the majority are at or just past flowering but before haying off is evident (Oct – Nov). Use of the higher rate in these crops is usually more reliable and gives a greater reduction in seed set.  Reduction in crop yield (in excess of 25%) may occur, especially if the ryegrass is more advanced than the crop. The higher rate may further decrease yield.  Apply by ground boom only in 50 –100L Spray with calibrated boom spray raised to give double overlap at the level of ryegrass seed heads. Pressures of 250 –350kPa and use of 110015 or 02 nozzles will aid coverage.
Spray topping to reduce seed set  Pastures	Grasses (particularly annual ryegrass)	All States	400mL	Heavily graze paddocks during spring flush to encourage even head development. Remove stock 2-3 weeks before the anticipated maturity date of the target species. However, if this is not feasible through lack of stock it is preferable to allow the pasture to mature ungrazed. Delay spraying until the last seed heads at the bottom of the plant has emerged and initial signs of haying off appear. Spray with a calibrated boom spray raised to give double overlap at the level of the seed heads.
	Barley Grass			Manage paddocks as above. Spray after head emergence but when all seed heads are green and there is no sign of haying off. Inspect paddocks before returning stock. Provided spraying was carried out before hardening of grass seeds stock (excluding horses) may be returned 24 hours after spraying. When hardening seeds are present harrow to knock seed from the heads. Do not introduce lambs into paddock until safe from risk of seed injury. If seasonal conditions favour regeneration, stock should be returned to selectively graze new shoots. Spray with a calibrated boom spray raised to give double overlap at the level of the seed heads.
	Saffron Thistle	NSW, SA, ACT only		Spray after the plant begins to run to head until flowering.
HORTICULTURE Potatoes -In crop -Pre harvest desiccation Row crops - pre-plant or before crop emerges -post emergent interrow weed control	Annual grasses and some seeding broadleaf weeds	All States	1.2 to 1.6L	Spray before 25% crop emergence. Use the higher rate for dense weeds.
	Pre-harvest weed control		2.8L	Spray about one week before digging and after tops have died down.
	Annual grasses and some broad leaf weeds		1.2 to 1.6L	Do not spray less than three days before sowing or three days before crop emergence. Use the lower rate for early autumn applications. Once the crop seedlings have emerged or transplanting has occurred, shielded nozzles must be used to ensure that spray does not contact crop.
	Seedling Weeds			Use the lower rate for early autumn applications
	Older Weeds		2.4L	Use on more mature stages of weed growth.



CROP USE OR SITUATION	WEEDS CONTROLLED	STATE	RATE/HA	CRITICAL COMMENTS
Peanuts Post –Emergence (In-crop)	<i>Datura</i> spp. (2-4 Leaf)	Qld, NT only	400mL	Spray peanuts up to 7-8 leaf stage but before majority of plants are flowering. Foliage will be scorched following application but plants recover rapidly. Apply in 200 – 250L/ha for thorough coverage of weed foliage. A dense canopy of weeds may reduce weed control due to shielding. Add <b>100mL wetting agent?? 1000/100L</b> of spray mix.  Do not spray on peanuts under extremely hot dry conditions when nuts are very small. In environments such as Far North Queensland use the lower rates in the range.
	Annual Ground Cherry (2-3 leaf), Apple of Peru (2-4 leaf), Milkweed (2-3 leaf)		600mL	
	Anoda Weed (2-4 leaf), Blue Heliotrope (2-3 leaf), Stagger Weed (2-3 leaf), Wandering Jew (2-3 leaf)		800mL	
	Bellvine (2-3 leaf), Common Morning Glory (2 leaf)		1L	
Non-Agricultural Situations, Around Sheds, Roadways, Paths	Annual Weed Control	All States	1.6 to 4L/ha or 200ml/100L	Spray to thoroughly wet weeds. Use the higher rate when weed growth is dense.  4Farmers Paraquat 250 Herbicide can be mixed with Diuron, Simazine, or Atrazine to give rapid weed knockdown and prolonged control.  If Capeweed or Erodium species are present add Reglone at 750mL to 1.5L/ha (125mL to 250mL/100L for high volume spraying).  If 4Farmers Paraquat 250 Herbicide rate is less than the ratio 500mL/100L add 100mL <b>suitable wetter per 100L of spray mix</b>
	Columbus Grass	NSW only	160mL/100L Plus 1L Flupropanate 745g/L (Spot Spraying) 2.3 to 4.5L/ha plus 12 to 22L Flupropanate 745g/L (Boom Spray)	If 4Farmers Paraquat 250 Herbicide rate is less than the ratio 500mL/100L <b>add 100mL suitable wetter per 100L of spray mix.</b>
Firebreaks	Reduce fire hazard by knocking down weed growth or to assist firebreak burn.	All States	1.6L to 4L	Apply mid-winter to early summer.  Use the higher rate for dense weed growth. After desiccation is complete the sprayed area may be burnt (approximately 7 – 10 days after spraying). 4Farmers Paraquat 250 Herbicide can be mixed with Diuron, Simazine or Atrazine to give rapid weed knockdown and prolonged control.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**THIS PRODUCT IS TOO HAZARDOUS TO BE USED IN THE HOME GARDEN, USE ONLY AS AN AGRICULTURAL OR HORTICULTURAL HERBICIDE.**

#### WITHHOLDING PERIOD

##### GRAZING:

DO NOT GRAZE OR CUT FOR STOCK FEED FOR 1 DAY AFTER APPLICATION.

DO NOT GRAZE HORSES FOR 7 DAYS AFTER APPLICATION.

REMOVE STOCK FROM TREATED AREA 3 DAYS BEFORE SLAUGHTER.

##### HARVESTING:

CHICKPEAS, FABA BEANS, FIELD PEAS AND LUPINS – DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION.

OTHER CROPS: NOT REQUIRED WHEN USED AS DIRECTED.



## GENERAL INSTRUCTIONS:

4Farmers Paraquat 250 Herbicide kills annual grasses and a number of broadleaf weeds (excluding capeweed) in specified conditions and should not be used for any other purpose. It acts quickly on green plant tissue and is inactivated in the soil. Weeds are most easily killed when actively growing and not under stress. Dusty conditions or heavy dew will impede product performance. This product has the potential to damage pasture legumes and reduce seed set. Medic species are more susceptible to damage than sub-clover species.

## RESISTANT WEEDS WARNING:

4Farmers Paraquat 250 Herbicide is a member of the BIPYRIDLYS group of herbicides. 4Farmers Paraquat 250 Herbicide has the INHIBITORS OF PHOTOSYNTHESIS AT PHOTOSYSTEM I mode of action. For weed resistance management 4Farmers Paraquat 250 Herbicide is a Group L herbicide.

Some naturally occurring weed biotypes resistant to 4Farmers Paraquat 250 Herbicide and other INHIBITORS OF PHOTOSYNTHESIS AT PHOTOSYSTEM I herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by 4Farmers Paraquat 250 Herbicide or other INHIBITORS OF PHOTOSYNTHESIS AT PHOTOSYSTEM I herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, 4Farmers Pty Limited accepts no liability for any losses that may result from the failure of 4Farmers Paraquat 250 Herbicide to control resistant weeds.

## MIXING

Use clean water, free of clay, silt and algae. Add required amount of 4Farmers Paraquat 250 Herbicide to water in spray tank while agitating to give even mixing.

When tank mixing with flowables, the following order should be strictly adhered to:

1. Fill tank with the required amount of water.
2. Add the flowable formulation and mix thoroughly.
3. Add the necessary amount of 4Farmers Paraquat 250 Herbicide and mix thoroughly.

## APPLICATION:

### FOR GROUND APPLICATION ONLY:

Do not spray if weeds are wilted by dry or cold weather or are covered by dust or soil. Best results are obtained when applied in dull weather or at the end of the day.

#### Broad acre spraying:

Apply through flat fan-jets adjusted to a height giving a double overlap at the top of the canopy. Use spraying pressures of 200-300kPa. Use water rate of 70L per hectare (50L in WA) or greater, when application equipment is appropriate. Ensure that water volume is able to achieve good leaf coverage of the target weeds.

#### Orchards, Vineyards, general weed control:

Use a water rate of up to 600L per Ha. Refer to safety directions and wash spray tank and equipment immediately after use with a general purpose cleaner.

#### Misting machine:

Banana plantations ONLY.

**Don't forget the 5mL per litre suitable wetter.** Run machine at half throttle to prevent the formation of fine droplets, which are hazardous. Adhere to safety directions and wear protective clothing, including waterproofs, impervious gauntlet length gloves, goggles and face mask and respirator covering nose and mouth and capable of filtering spray droplets. Respirators should comply with the requirements of AS1718 (Standards Association of Australia).

#### Compatibility:

4Farmers Paraquat 250 Herbicide is compatible with Trifluralin, flowable and granular formulations of Diuron, Atrazine and Simazine, and Chlorsulfuron. See "Mixing Instructions".

## PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DRIFT WARNING: DO NOT apply under meteorological conditions or from spray equipment which could be expected to cause spray to drift onto nearby susceptible plants, adjacent crops, crop lands or pastures. DO NOT apply on or near water which is used for irrigation purposes.

## PROTECTION OF LIVESTOCK:

Keep domestic pets and poultry away from treated areas. This formulation should not be applied on or near water which is used for livestock watering.

## PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND THE ENVIRONMENT

Do not contaminate dams, rivers, drains or streams with chemical or used container. This formulation should not be applied on or near water, which is used for human consumption, livestock watering or irrigation purposes or water used for commercial or recreational fishing.

## STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. Do not store for prolonged periods in direct sunlight.

Do not repack or use container for any other purpose. Triple or preferably pressure rinse empty containers adding the rinsate to the spray tank. Do not dispose undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or





designated collection point. If not recycling, break, crush or puncture and bury containers in a local authority landfill. If not available bury the containers below 500mm in a disposal pit specifically set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

**For refillable containers:**

Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

**SAFETY DIRECTIONS**

Very dangerous, particularly the concentrate

Product is poisonous if swallowed. Will irritate the nose, throat and skin.

Attacks eyes, protect eyes while using. Avoid contact with eyes, skin and clothing.

When opening the container and preparing the product for use, wear elbow length PVC gloves and face shield or goggles. If product on skin, immediately wash area with soap and water. If clothing becomes contaminated with product, remove clothing immediately. If product in eyes, wash it out immediately with water. Avoid contact with spray mist. Do not inhale spray mist. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash contaminated clothing, gloves and face shield or goggles.

**Spray Application:**

DO NOT work in spray mist.

DO NOT continue to use if skin irritation or nosebleed occurs. This may be caused by exposure to spray mist as the result of incorrect use of equipment or adverse climatic conditions. Stop and review handling and spraying techniques before further spraying. If symptoms persist seek medical advice.

Where there is a risk of exposure to spray mist wear waterproof footwear and waterproof protective clothing, impervious gauntlet length gloves (rubber or PVC), goggles and a face mask and respirator covering nose and mouth and capable of filtering spray droplets. A high efficiency type particulate respirator is recommended, but in any event use a respirator which complies with the requirements of AS1716 (Standards Association of Australia). Further advice on safety equipment should be obtained from a safety equipment manufacturer.

Avoid contacting vegetation wet with spray, but if necessary to do so wear waterproof footwear and waterproof protective clothing and gloves.

**FIRST AID:**

If poisoning occurs, get to a doctor or hospital quickly. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

**MATERIAL SAFETY DATA SHEET:**

Additional information is listed in the Material Safety Data Sheet.

**CONDITIONS OF SALE:**

The use of 4 Farmers Paraquat 250 Herbicide, being beyond the control of the manufacturer, no warranty expressed or implied is given by 4 Farmers Pty. Ltd. regarding its suitability, fitness or efficacy for any purpose for which it is used by the buyer, whether in accordance with the directions or not and 4 Farmers Pty. Ltd. accepts no responsibility for any consequences whatsoever resulting from the use of this product.

**4FARMERS PTY LTD**  
**A.C.N. 067 443 485**  
**643 Murray Street**  
**WEST PERTH WESTERN AUSTRALIA 6005**

**Phone: (08) 9356 3445**  
**Fax: (08) 9356 3447**

Toxic 6 diamond

UN No. 3016

Pesticides, Bipyridilium Liquid toxic (contains paraquat).

Hazchem Code 2X

In an Emergency Dial 000 Police or Fire Brigade.