

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Crop Care

STOMP[®] Xtra

Herbicide

ACTIVE CONSTITUENT: 455 g/L PENDIMETHALIN

GROUP **D** HERBICIDE

For the control of annual ryegrass and wireweed in wheat, barley and peas and annual grasses and certain broadleaved weeds in various crops as shown in the DIRECTIONS FOR USE table.

IMPORTANT: READ THE ATTACHED BOOKLET BEFORE USING THIS PRODUCT

**Contents: 20 Litres
110 Litres
Bulk**

Crop Care Australasia Pty Ltd, 77 Tingira Street Pinkenba, Qld, 4008

® Registered trademark of BASF

DIRECTIONS FOR USE

FOR THE FULL DIRECTIONS FOR USE, READ THE ATTACHED BOOKLET BEFORE USING THIS PRODUCT.

STORAGE AND DISPOSAL

20 L ONLY

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should NOT be burnt.

BULK ONLY

Store in the closed, original container in a cool, well-ventilated area. Do NOT store for prolonged periods in direct sunlight.

Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

Do NOT dispose of undiluted chemicals on-site. This container remains the property of Crop Care Australasia Pty Ltd

110 L ONLY

Store in the closed, original container in a cool, well-ventilated area.

Do NOT store for prolonged periods in direct sunlight. Empty container by pumping through the dry-break Micro Matic connection system. Do NOT attempt to unscrew the Micro Matic valve or breach the locked filling point. Do NOT contaminate the container with water or other foreign material. Ensure that the Micro Matic coupler, pump, meter and hoses are disconnected, triple rinsed with clean water and drained after each use. When empty, or contents no longer required, return the container to the point of purchase. Do NOT dispose of undiluted chemicals on-site. This container remains the property of Crop Care Australasia Pty Ltd

SAFETY DIRECTIONS

Will irritate the eyes. Avoid contact with eyes. Wash hands after use. When preparing and using the prepared spray wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length PVC gloves. After each day's use, wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data sheet which can be obtained from your supplier.

CONDITIONS OF SALE

Crop Care Australasia Pty Ltd will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply or use of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Crop Care Australasia Pty Ltd is limited to the replacement of the goods or (at the option of Crop Care Australasia Pty Ltd) the refund of the price paid and is conditional upon a claim being made in writing and where possible sufficient part of the goods to enable proper examination being returned to Crop Care Australasia Pty Ltd within thirty days of delivery.

In a Transport Emergency Dial 000 Police or Fire Brigade	SPECIALIST ADVICE IN AN EMERGENCY ONLY 1800 033 498 ALL HOURS- AUSTRALIA WIDE
--	---



Crop Care Australasia Pty Ltd
ACN 061 362 347
77 Tingira Street
PINKENBA Qld 4008
Ph : (07) 3867 9100

Insert Barcode
Label code and
drumMUSTER logo
as required

APVMA Approval No.: 58924/20L/0305
58924/110L/0305
58924/BULK/0305

Batch No:

Date of Manufacture:

Print code: 06-04/ /0762/20L
06-04/ /0762/110L
06-04/ /0762/BULK

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Crop Care

STOMP[®] Xtra

Herbicide

ACTIVE CONSTITUENT: 455 g/L PENDIMETHALIN

GROUP D HERBICIDE

For the control of annual grasses and certain broadleafed weeds in various crops as shown in the DIRECTIONS FOR USE table.

THIS BOOKLET IS PART OF THE LABEL.

Crop Care Australasia Pty Ltd, 77 Tingira Street, Pinkenba Qld 4008

APVMA Approval No.: 58924/0305

® Registered trademark of BASF

Print code: 06-04/ /0762

DIRECTIONS FOR USE

RESTRAINTS:

Do NOT sow sensitive summer crops such as sorghum and millets within 12 months of application of STOMP Xtra.

Do NOT apply to a poorly prepared seedbed.

Do NOT use STOMP Xtra mixed with atrazine on heavy clay soils.

Do NOT apply STOMP Xtra where waterlogging is likely to occur.

Do NOT apply STOMP Xtra by aerial application to sugar cane.

Crop	Weeds Controlled	State	Rate/ Treated ha	Critical Comments
1. PRE-PLANT INCORPORATED TREATMENT (DOUBLE INCORPORATION)				
Cotton	See Weed Table A	Qld, NSW only	2.2 L/ha	<p>Incorporate into the soil within 24 hours of application. Use a combine with trailing harrows and chain or offset or tandem discs or Lilliston cultivators, set to work to a depth of 2-5 cm. Subsequent planting operations in irrigated cotton may remove the STOMP Xtra treated soil from hills leaving an untreated band over the row, requiring a further application of STOMP Xtra or other herbicides in that band. When the seedbed is rough, a second incorporation is necessary. STOMP Xtra may be applied by aerial or ground spraying. In the Macquarie Valley area, only apply by air when ground is too wet for ground application. (Refer note on "Incorporation" (a)).</p> <p>Use the higher rate on heavy textured soils or those high in organic matter and the lower rate on light to medium textured soils.</p>
Cow peas, mung beans, navy beans, peanuts, pigeon peas and soybeans	See Weed Table A	Qld, NSW, ACT, NT only	1.8 to 2.2 L/ha	
Sunflowers	See Weed Table A	Qld, NSW, ACT, Vic, SA, Tas only		
2. POST-PLANT PRE-EMERGENCE (AND POST EMERGENCE, ONION ONLY) SURFACE TREATMENT (OVERALL OR BAND SPRAYING): NON-MECHANICAL INCORPORATED				
Cotton	See Weed Table B	Qld, NSW only	3.3 L/ha	<p>COTTON: Where incorporation prior to sowing is impractical and where the seedbed tilth is fine and free of large stones and trash, STOMP Xtra should be applied within 48 hours after sowing.</p> <p>SUNFLOWERS: Where incorporation prior to sowing is impractical and where the seedbed tilth is fine and free of large stones and trash, STOMP Xtra should be applied after sowing and before the emergence of the crop and weeds. Where fat hen or blackberry nightshade is expected to be a major problem, pre-plant incorporation (No. 1 application method), is preferred. (Refer section on "Incorporation" (a) and (b)).</p>
Sunflowers		Qld, NSW, ACT, SA only		
Broad beans	Poppies, wireweed (<i>Polygonum aviculare</i>)	Tas only	2.9 L/ha	<p>MINIMUM PLANTING DEPTH</p> <p>Carrots: 15 mm</p> <p>Processing peas, Broad beans and French beans: 30 mm</p>
Carrots	See Weed Table D	Qld, NSW,	1.5 to 2.2 L/ha	<p>Seed should be sown with press wheels or rollers behind the planter in order to ensure coverage of the seed and compaction of the seed bed.</p> <p>Apply STOMP Xtra to a fine firm seedbed free of ridges, clods and trash within 2 days after sowing. Use the higher rates on heavier textured soils or those with a higher organic matter content.</p> <p>Incorporate with 12 to 25 mm of spray irrigation or rainfall applied within one day of application for optimum performance.</p> <p>DO NOT disturb the soil by rolling or harrowing after STOMP Xtra has been applied. DO NOT apply where heavy rainfall or irrigation is likely to lead to water-logging.</p> <p>DO NOT use STOMP Xtra on soils with an organic matter content above 6% as inconsistent weed control will occur.</p> <p>FRENCH BEANS: For Qld, use in the Lockyer and Fassifern valleys only.</p> <p>DO NOT apply to French beans sown during the autumn, winter or early spring or which are likely to be stressed by cold weather.</p>
	Weed Table D plus Sowthistle (<i>Sonchus oleraceus</i>), wild radish (<i>Raphanus raphanistrum</i>)	ACT, Vic, SA, Tas, WA only	1.5 L/ha plus linuron at 1.0 kg ac/ha	
French beans	See Weed Table D	Qld, Tas only	2.2 to 2.9 L/ha	<p>DO NOT use STOMP Xtra on soils with an organic matter content above 6% as inconsistent weed control will occur.</p> <p>FRENCH BEANS: For Qld, use in the Lockyer and Fassifern valleys only.</p> <p>DO NOT apply to French beans sown during the autumn, winter or early spring or which are likely to be stressed by cold weather.</p>
Processing peas		Qld, Vic only	1.5 to 2.9 L/ha	
		Tas only	2.9 L/ha	

Crop	Weeds Controlled	State	Rate/ Treated ha	Critical Comments
2. POST-PLANT PRE-EMERGENCE (OR POST EMERGENCE, ONION ONLY) SURFACE TREATMENT (OVERALL OR BAND SPRAYING): NON-MECHANICAL INCORPORATED (continued)				
Maize	See Weed Table B	Qld, NSW, ACT, NT only	3.3 L/ha	Where grasses are expected to be the major problem, this treatment is recommended - STOMP Xtra alone. Apply STOMP Xtra after planting and before emergence of crop and weeds. Do NOT incorporate this treatment. Avoid application in dry weather; this may cause pruning of aerial roots and lodging. (Refer sections on "Incorporation" (b) and (c)).
	See Weed Table C		2.2 L/ha plus atrazine at 1.0 kg ac/ha	Where thornapple, Noogoora burr, Bathurst burr, caltrop, paddy melon or mintweed are expected to be the major problems, STOMP Xtra should be applied in a tank mix combination with atrazine after planting and before the emergence of crop and weeds. Do NOT incorporate this mixture. Follow directions on atrazine label carefully. This treatment should not lead to soil residue problems with atrazine as the application rate is low. Do NOT use STOMP Xtra mixed with atrazine on heavy clay soils as weed control may not be adequate. (Refer section on "Incorporation" (b) and (c)).
Onions	Hogweed (wireweed) (<i>Polygonum aviculare</i>)	Tas only	550 mL to 730 mL	<p>Apply in conjunction with other herbicides as advised by a qualified crop adviser to ensure that an adequate level of weed control is achieved in adverse conditions.</p> <p>Pre-emergence: Do NOT use on sandy soils. MINIMUM PLANTING DEPTH: 15 mm</p> <p>Apply from immediately after sowing until just prior to emergence. Apply STOMP Xtra to a fine, firm seedbed free of clods and trash. Use the higher rates on heavy textured soils or those with a high organic matter content. For optimum performance, incorporate with no more than 12 mm of spray irrigation within one day of application. DO NOT exceed 550 mL /ha if soil conditions are wet and cold.</p>
			730 mL to 1.45 L	<p>Post-emergence: Apply STOMP Xtra from the first true leaf until the 3 leaf stage. Repeat applications may be made after the 1 leaf up to the 3 leaf stage providing total usage of STOMP Xtra on crop does not exceed 2.2 L/ha. Avoid applying STOMP Xtra to areas where water logging is likely to occur.</p>
		NSW, ACT, VIC, SA and WA only	<ul style="list-style-type: none"> ♦ Apply in conjunction with other herbicides as advised by a qualified crop adviser to ensure that an adequate level of weed control is achieved in adverse conditions. ♦ Depth of sowing is critical. Onions must be sown to a depth of at least 15mm with press wheels or rollers behind planter to ensure coverage and compaction. ♦ DO NOT disturb the soil by rolling or harrowing after application. ♦ For optimum performance, incorporate with spray irrigation within a day of planting. However, DO NOT water heavily (no more than 12 mm irrigation) for at least 5 days after application. ♦ AVOID application to areas where water logging is likely to occur. ♦ DO NOT apply if heavy rain is likely within a week of application, as crop damage may result. ♦ DO NOT exceed 550 mL /ha if soil conditions are wet and cold. ♦ DO NOT apply more than 3 times per season. ♦ DO NOT apply more than 2.2 L/ha per season. 	
			550 mL to 730 mL (see Critical comments)	<p>Light/ sandy soils DO NOT apply pre-emergence. Application to moist soil is best. Apply 550-730 mL/ha post emergence at the 4-5 leaf stage, with 1-2 further applications of 550-730 mL/ha as needed throughout the season (approx 3-4 weeks apart)</p>
			730mL to 1.45 L (see Critical comments)	<p>Heavy/high organic matter soils Application to moist soil is best. Apply 730-1.45 L/ha pre-emergence within 2 days of planting. Apply to a fine firm seed bed free of ridges, clods and trash. For furrow irrigated crops, application to a dry seed bed, followed by irrigation within 5 days is optimal. An application of 730mL-1.1 L/ha post emergence may be made at the 2 leaf stage. (Application at this stage may cause yellowing of onions).</p>

Crop	Weeds Controlled	State	Rate/ Treated ha	Critical Comments
3. POST-PLANT PRE-EMERGENCE TREATMENT (INCORPORATED OR NON-INCORPORATED)				
Sugarcane	Awnless barnyard grass (<i>Echinochloa colona</i>), crowsfoot grass (<i>Eleusine indica</i>), green summer grass (<i>Brachiaria subquadrifera</i>), guinea grass (<i>Panicum maximum</i> cv. <i>Hamil</i> and cv. <i>Guinea</i>), summer grass (<i>Digitaria ciliaris</i>)	Qld, NSW, WA only	2.2 to 3.3 L/ha	STOMP Xtra should be applied as soon as possible after planting and before sugarcane and weed emergence. Consolidation (cutaway operation) of the planting furrow prior to application to prevent soil movement into the base of the drill is recommended. Poor grass control is expected in soils which contain a high percentage of organic matter. (Refer section on "Incorporation" (a)). Use the higher rate where longer weed control is required or where incorporation by irrigation or rainfall is likely to be delayed more than 5 days. INCORPORATED: Under dry conditions or where some cultivation is expected after application STOMP Xtra should be thoroughly incorporated by finger rakes to a shallow depth. In subsequent tillage operations, cultivation depth should not exceed the depth of incorporation. WATER INCORPORATION: Incorporate using 12 to 25 mm of spray irrigation or rainfall. Incorporate as soon as possible and within 3-5 days after application. The area should not be disturbed by cultivation for the expected duration of weed control where this method of application is used. If excessive irrigation or rainfall occurs after the application of STOMP Xtra, resulting in the movement of soil from the top to the bottom of the planting drill, weed control will be reduced. Flood irrigation may not incorporate STOMP Xtra adequately if soils in the treatment area are variable or difficult to wet.
	Blue top (blue billygoat weed) (<i>Ageratum houstonianum</i>)		2.2 L/ha plus atrazine at 1.5 kg ac/ha or diuron at 1.5 kg ac/ha	For other broadleaved weeds higher rates of atrazine or diuron can be tank mixed with STOMP Xtra - see the relevant atrazine or diuron labels for details.
	Barnyard grass (<i>E. crus-galli</i>), blackberry nightshade (<i>Solanum nigrum</i>), blue top (blue billygoat weed) (<i>Ageratum houstonianum</i>), common sida (<i>Sida rhombifolia</i>), giant/black pigweed (<i>Trianthema portulacastrum</i>), glossy nightshade (<i>Solanum americanum</i>), green amaranth (<i>Amaranthus viridis</i>), Ipomoea spp., liverseed grass (<i>Urochloa panicoides</i>), milkweed (Mexican fire plant) (<i>Euphorbia heterophylla</i>), pigweed (<i>Portulaca oleracea</i>)		2.2 L/ha plus 400 mL/ha Flame [®] Herbicide	Incorporate into the soil either by overhead irrigation, rainfall or mechanical incorporation, to a minimum depth of 5 cm. Incorporate as soon as possible and within 3-5 days after application. Read all directions for use in sugarcane on the Flame Herbicide label.

Crop	Weeds Controlled	State	Rate/ Treated ha	Critical Comments
4. PRE-TRANSPLANT TREATMENT				
Transplanted broccoli, cabbage, cauliflower and processing tomatoes	See Weed Table D	All states	1.5 to 2.2 L/ha	Apply STOMP Xtra to a firm seedbed free of ridges, clods and trash between 7 and 2 days before transplanting. Use the higher rates on heavier textured soils or those with a higher organic matter content. Incorporate with 12 to 25 mm of spray irrigation or rainfall applied within one day of application for optimum performance. Mechanical disturbance to the seed bed at transplanting should be minimised to achieve optimum performance. DO NOT apply after transplanting as damage will occur. DO NOT apply where water-logging is likely to occur after transplanting or crop stunting will result. DO NOT use STOMP Xtra on soils with an organic matter content above 6% as inconsistent weed control will occur. Ramrod* may be tank mixed with STOMP Xtra prior to transplanting or applied separately after transplanting.
	Weed Table D plus dwarf amaranth (<i>Amaranthus macrocarpus</i>), green amaranth (<i>Amaranthus viridis</i>), annual nettles (<i>Urtica spp.</i>), Indian hedge mustard (<i>Sisymbrium orientale</i>), potato weed (<i>Galinsoga parviflora</i>), sowthistle, turnip weed (<i>Rapistrum rugosum</i>), wild radish, wild turnip (<i>Brassica tournefortii</i>), winter grass (<i>Poa annua</i>)		1.5 to 1.8 L/ha plus 9 L/ha Ramrod Flowable	
Transplanted lettuce	See Weed Table D		1.5 to 2.9 L/ha	
5. PERENNIAL CROPS				
Avocados, bananas, citrus, deciduous fruits (not grapevines), lychees, macadamia nut, mangoes, nuts, olives	Dwarf amaranth, green amaranth, annual ryegrass, asthma plant (<i>Euphorbia hirta</i>), barnyard grass, chickweed (<i>Stellaria media</i>), crowsfoot grass, deadnettle (<i>Lamium amplexicaule</i>), fat hen (<i>Chenopodium album</i>), pigeon grass, pigweed (<i>Portulaca oleracea</i>), prickly lettuce (<i>Lactuca serriola</i>), sowthistle, summer grass, winter grass, wireweed	All states	6.5 to 8.7 L/ha	STOMP Xtra alone or STOMP Xtra plus simazine is likely to affect the emergence and growth of green manure crops such as oats or beans between rows of trees, if they are sown into the treated band within 12 months of application. Use the higher rate on medium to heavy textured soils and the lower rate on light textured soils. Use a directed spray avoiding spray contact with green bark, fruit and foliage. If applied to freshly transplanted trees, soils should be compacted prior to application of STOMP Xtra to avoid contact with roots. Crop damage and/or reduced weed control may result if STOMP Xtra is not incorporated successfully. If insufficient rainfall follows within 10 days after application, the product must be incorporated over the whole surface area treated through the use of overhead sprinklers or by mechanical means, using appropriate equipment to avoid root damage. Drip irrigation is inadequate for this purpose. A minimum of 10 - 15mm of rain or sprinkler irrigation is normally required to incorporate STOMP Xtra into the soil. For best results, soil surface should be free of weeds, surface litter and clods at the time of application. If small weeds are present at the time of application, STOMP Xtra should be tank mixed with a knockdown herbicide at the recommended rate.
	As above plus prairie grass (<i>Bromus unioioides</i>), spotted medic (<i>Medicago arabica</i>)		8.7 L/ha	
Citrus, deciduous fruits (not grapevines) and nuts	As above plus cobbler's pegs (<i>Bidens pilosa</i>), curious weed (<i>Schkuhria pinnata</i>), stinking Roger (<i>Tagetes minuta</i>)		6.5 L/ha plus simazine at 800 g ai/ha	
As above plus caltrop (<i>Tribulus terrestris</i>)	8.7 L/ha plus simazine at 1 kg ai/ha			

Crop	Weeds Controlled	State	Rate/ Treated ha	Critical Comments
5. PERENNIAL CROPS (CONT.)				
Eucalypt forestry plantations	Amaranths (<i>Amaranthus spp.</i>), annual rye grass, asthma plant (<i>Euphorbia hirta</i>), barnyard grass (<i>Echinachloa crus-galli</i>), chickweed (<i>Stellaria media</i>), crowfoot grass (<i>Eluesine indica</i>), deadnettle (<i>Lamium amplexicaule</i>), fat hen (<i>Chenopodium album</i>), pigeon grass (<i>Setaria spp.</i>), pigweed (<i>Portulaca oleracea</i>), prickly lettuce (<i>Lactuca serriola</i>), stagger weed (<i>Stachys arvensis</i>), sow thistle (<i>Sonchus oleraceus</i>), summer grass (<i>Digitaria ciliaris</i>), winter grass (<i>Poa annua</i>), wireweed (<i>Polygonum aviculare</i>)	All States	6.5 L to 8.7 L/ha	PRE-PLANTING: Apply 7 to 2 days prior to transplanting. Use the higher rate on heavier textured soils or soils with higher organic matter. Mechanical disturbance should be minimised during planting to optimise performance. POST-PLANTING: Apply immediately following planting, before emergence of weeds. Repeat applications may be made as directed sprays to provide pre-emergence control of weeds. Emerged weeds should be controlled with carefully directed applications of a non-selective herbicide or over-the-top application of a selective herbicide. If applied to freshly transplanted trees, soils should be compacted prior to application to avoid contact with roots. Weed control may be reduced if rainfall or irrigation does not occur within 10 days of application. Flood irrigation may reduce control.
Teatree (<i>Melaleuca alternifolia</i>)	Weed Table B plus dwarf amaranth, green amaranth, annual ryegrass, barnyard grass, chickweed (<i>Stellaria media</i>), crowfoot grass, deadnettle (<i>Lamium amplexicaule</i>), fat hen, pigeon grass, pigweed, prickly lettuce (<i>Lactuca serriola</i>), sowthistle, summer grass, winter grass, wireweed	NSW, ACT, Qld, SA, NT only	3.3 to 6.5 L/ha	DO NOT tank mix with simazine as crop damage may result. For best results the soil surface should be free of weeds, surface litter and clods at the time of application. PRE-TRANSPLANTING: Apply 7 to 2 days prior to transplanting seedlings. Use the higher rates on heavier textured soils or soils with higher organic matter. Mechanical disturbance should be minimised during planting to optimise performance. Avoid contact of treated soil with seedling roots. POST-PLANTING: Apply immediately following planting and pre-emergence of weeds. If applied to freshly transplanted trees, soils should be compacted prior to application of STOMP Xtra to avoid contact with roots. Weed control may be reduced if rainfall or irrigation does not occur within 10 days of application. Flood irrigation may reduce control.
6. IRRIGATION CHANNELS				
Cotton irrigation banks, channels and drains	Barnyard grass, silvertop grass (brown beetle grass)	Qld, NSW only	3.3 to 6.5 L/ha	Use the lower rate when short term control is required. Apply in late winter or early spring after regarding or clearing channels. The soil should be loose and free of large clods. If 25-50 mm of rainfall has not fallen within 14 days of application the channel should be filled with water and allowed to stand for 1 day. The water in the channel should then be drained off and used to pre-irrigate cotton fields. DO NOT use water in channel to irrigate or pre-irrigate susceptible crops.
			4.4 L/ha plus diuron at 8.1 kg ac/ha	Use this mixture where long term control is required or a broader weed spectrum is present. See diuron label for details. Water in the channel should NOT be used to irrigate or pre-irrigate susceptible crops.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED

A. IN COTTON, COW PEAS, MUNG BEANS, NAVY BEANS, PEANUTS, PIGEON PEAS, SOYBEANS, SUNFLOWERS - with pre-plant incorporated treatment (double incorporation)

Barnyard grass Button grass Common heliotrope	<i>Echinochloa spp.</i> <i>Dactyloctenium radulans</i> <i>Heliotropium europeum</i>
Crested goosefoot Crowsfoot grass Dwarf amaranth	<i>Chenopodium cristatum</i> <i>Eleusine indica</i> <i>Amaranthus macrocarpus</i>
Early spring grass Fat hen Green amaranth Liverseed grass Mexican clover (white eye) Mossman River grass	<i>Eriochloa spp.</i> <i>Chenopodium album</i> <i>Amaranthus viridis</i> <i>Urochloa panicoides</i> <i>Richardia brasiliensis</i> <i>Cenchrus echinatus</i>
Native millet Pale pigeon grass Paspalidium (Brigalow grass) Pepper grass Pigweed Queensland blue grass	<i>Panicum decompositum</i> <i>Setaria glauca</i> <i>Paspalidium spp.</i> <i>Panicum whitei</i> <i>Portulaca oleracea</i> <i>Dichanthium sericeum</i>
Red Flinders grass Scarlet pimpernel Small burr grass Stink grass Weeping love grass	<i>Iseilema vaginiflorum</i> <i>Anagallis arvensis</i> <i>Tragus australianus</i> <i>Eragrostis cilianensis</i> <i>Eragrostis parviflora</i>
Wireweed	<i>Polygonum aviculare</i>

Useful suppression of the following weeds is also obtained:

Blackberry nightshade Caltrop Common verbena Mintweed Peppergrass Stagger weed	<i>Solanum nigrum</i> <i>Tribulus terrestris</i> <i>Verbena officinalis</i> <i>Salvia reflexa</i> <i>Lepidium spp.</i> <i>Stachys arvensis</i>
---	---

B. IN COTTON, SUNFLOWERS, MAIZE and TEATREE - with post-plant pre-emergence surface treatment (STOMP Xtra alone)

Barnyard grass Button grass	<i>Echinochloa spp.</i> <i>Dactyloctenium radulans</i>
Common heliotrope Crested goosefoot Crowsfoot grass Dwarf amaranth	<i>Heliotropium europeum</i> <i>Chenopodium cristatum</i> <i>Eleusine indica</i> <i>Amaranthus macrocarpus</i>
Early spring grass Green amaranth Liverseed grass Mexican clover (white eye) Mossman River grass Native millet	<i>Eriochloa spp.</i> <i>Amaranthus viridis</i> <i>Urochloa panicoides</i> <i>Richardia brasiliensis</i> <i>Cenchrus echinatus</i> <i>Panicum decompositum</i>
Pale pigeon grass Paspalidium (Brigalow grass) Pepper grass Pigweed Queensland blue grass	<i>Setaria glauca</i> <i>Paspalidium spp.</i> <i>Panicum whitei</i> <i>Portulaca oleracea</i> <i>Dichanthium sericeum</i>
Red Flinders grass Scarlet pimpernel Small burr grass Stink grass Weeping love grass	<i>Iseilema vaginiflorum</i> <i>Anagallis arvensis</i> <i>Tragus australianus</i> <i>Eragrostis cilianensis</i> <i>Eragrostis parviflora</i>
Wireweed	<i>Polygonum aviculare</i>

Useful suppression of the following weeds is also obtained:

Caltrop Common sowthistle Common verbena Mintweed Peppergrass Stagger weed	<i>Tribulus terrestris</i> <i>Sonchus oleraceus</i> <i>Verbena officinalis</i> <i>Salvia reflexa</i> <i>Lepidium spp.</i> <i>Stachys arvensis</i>
---	--

C. IN MAIZE - with post-plant pre-emergence surface treatment of STOMP Xtra plus tank mix with atrazine.

The following weeds are controlled in addition to those controlled by STOMP Xtra alone:

Bathurst burr Caltrop Mintweed Noogoora burr Paddy melon Thornapple	<i>Xanthium spinosum</i> <i>Tribulus terrestris</i> <i>Salvia reflexa</i> <i>Xanthium pungens</i> <i>Cucumis spp.</i> <i>Datura spp.</i>
--	---

The following weeds are not controlled by STOMP Xtra or the mixture with atrazine:

Bindweed Wandering jew Nut grass	<i>Convolvulus spp.</i> <i>Commelina spp.</i> <i>Cyperus rotundus</i>
--	---

D. IN CARROTS, PROCESSING PEAS, FRENCH BEANS AND TRANSPLANTED CABBAGE, CAULIFLOWER, BROCCOLI, LETTUCE, PROCESSING TOMATOES - with pre-emergence surface treatment after planting or before transplanting as directed and incorporated by spray irrigation.

The following weeds are controlled:

Awnless barnyard grass Bittercress Bladder ketmia Chickweed Deadnettle Fat hen Green fat hen Pigweed Prickly lettuce Shepherd's purse Wireweed	<i>Echinochloa colona</i> <i>Coronopus didymus</i> <i>Hibiscus trionum</i> <i>Stellaria media</i> <i>Lamium amplexicaule</i> <i>Chenopodium album</i> <i>Chenopodium murale</i> <i>Portulaca oleracea</i> <i>Lactuca serriola</i> <i>Capsella bursa-pastoris</i> <i>Polygonum aviculare</i>
--	---

Useful suppression only of the following weeds at the lower rates. Higher rates or tank mixtures are required for complete control (see Directions for Use table).

Annual nettles Blackberry nightshade Common sowthistle Fumitory Indian hedge mustard Turnip weed Wild radish Winter grass	<i>Urtica spp.</i> <i>Solanum nigrum</i> <i>Sonchus oleraceus</i> <i>Fumaria spp.</i> <i>Sisymbrium orientale</i> <i>Rapistrum rugosum</i> <i>Raphanus raphanistrum</i> <i>Poa annua</i>
--	---

GENERAL INSTRUCTIONS

STOMP Xtra Herbicide is a selective herbicide for the control of most annual grasses and certain broadleaved weeds. STOMP Xtra controls weeds by inhibiting seedling development; it will not control established weeds. For best results, seedbeds should be free of weeds, trash and clods at the time of application. Soils containing a high percentage of organic matter can result in poor control. It is recommended that application to soils containing a high percentage of organic matter (greater than 6%) be avoided.

Crop injury may result if cold, wet weather follows planting or if STOMP Xtra is applied to a poorly prepared seedbed. With the return of favourable growing conditions (warm weather), the crop will usually recover and resume normal growth.

Caution: prolonged periods of rainfall and/or excessive soil moisture at or following application may delay breakdown of STOMP Xtra capsules, resulting in reduced weed control. STOMP Xtra may stain (yellow) any objects it contacts. Keep all objects etc. that are not to be stained well clear of the product and spray.

MIXING

Shake, invert or roll container several times before use. Put half the required volume of water in spray tank and start agitation. When using STOMP Xtra alone, add STOMP Xtra to partially filled tank and then add the remainder of the water. Mix thoroughly.

When using STOMP Xtra in tank mixes with products such as atrazine or diuron, mix these products as indicated on the relevant label and add the STOMP Xtra last, then complete filling of the tank. Maintain good agitation at all times until spraying is completed. If the spray mixture is allowed to settle, thorough agitation is essential to re-suspend the mixture before spraying is resumed. Note that when tank mixing STOMP Xtra with paraquat and glyphosate herbicides, the STOMP Xtra should be thoroughly mixed in the spray tank first, before adding a paraquat or glyphosate product.

APPLICATION METHOD

Ground Application: Use conventional sprayers with either mechanical or by-pass agitation. Flat fan nozzles should be used. Spray equipment should be correctly calibrated to ensure proper application. If applying STOMP Xtra in tank mix combination with atrazine or diuron, nozzle screens must be no finer than 50 mesh. Apply the recommended quantity of STOMP Xtra in 50-200 litres of water per hectare. Where pre-emergence surface treatment is used, the higher rates of water are recommended.

Aerial Application:

Pre-plant soybeans, cotton, sunflowers, peanuts, navy beans, cow peas, mung beans and pigeon peas only.

Apply the recommended quantity of STOMP Xtra in 25-60 litres of water per hectare. It is essential that the equipment is correctly calibrated and an even spray pattern is obtained. It is essential that markers be used and an accurate swath width is maintained. DO NOT apply if wind speed is greater than 16 km/hour. Always spray with a cross wind in accordance with recommended flying practice.

DO NOT apply STOMP Xtra Herbicide by aerial spraying in eucalypt forestry situations.

INCORPORATION:

- (a) STOMP Xtra works best if thoroughly mixed with soil either mechanically or by irrigation or rainfall. The aim of incorporation is to produce an even band of herbicide to intercept germinating weed seeds.
- (b) Post-plant pre-emergence surface applications of STOMP Xtra alone, or as a tank mixture, are most effective in controlling weeds when adequate rainfall occurs or irrigation is applied as soon as possible and within 3-5 days after treatment to achieve incorporation. If no rain or irrigation is received within 3-5 days, light mechanical incorporation is required for optimum weed control, except when used on maize and carrots or on sugarcane, if water incorporation is used.
- (c) Maize is tolerant to STOMP Xtra either alone or in mixtures with atrazine if the seed germinates below the chemically treated band. Maize, carrots and rice are NOT tolerant to STOMP Xtra if it is in direct contact with the seed.

WEATHER CONDITIONS AT PLANTING

Crop injury may result if cold weather follows planting. With the return of favourable growing conditions (warm weather), crops usually recover and resume normal growth.

RESISTANT WEEDS WARNING



STOMP Xtra Herbicide is a member of the dinitroaniline group of herbicides. STOMP has the inhibition of tubulin formation mode of action. For weed resistance management, STOMP is a Group D herbicide. Some naturally-occurring weed biotypes resistant to STOMP and other Group D herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by STOMP or other Group D herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Crop Care Australasia Pty Ltd accepts no liability for any losses that may result from the failure of STOMP to control resistant weeds.

COMPATIBILITY

Physical compatibility of STOMP Xtra has been tested with the following herbicides: atrazine, diuron, simazine, prometryn, fluometuron, fluometuron plus prometryn, ametryn, paraquat, propanil, linuron, glyphosate, Ramrod* Flowable, Logran+/Nugran+ and Avadex*Xtra (trilalate). If tank mixing, observe the directions under mixing.

FOLLOWING CROPS

In the event of crop failure the soil should be cultivated to a minimum depth of 15 cm to ensure any residues are evenly dispersed throughout the soil.

If a maize crop fails due to weather conditions or some other reason, maize can be replanted the same year without adverse effects but seeding depth must be below the retilled area.

The following crops may be sown after a minimum interval of 2 months after application of STOMP Xtra:

carrots, parsnips, parsley, celery, potatoes, peas, French beans, transplanted brassicas, transplanted lettuce, transplanted tomatoes, transplanted capsicums.

The following crops may be sown after a minimum interval of 5 months after application of STOMP Xtra:

turnips, lettuce, radish, Brussels sprouts, cabbage, cauliflower, broccoli, onions, leeks, sweetcorn, pumpkins, squash, melons, cucumbers.

The following crops may be sown after a minimum interval of 12 months after application of STOMP Xtra:

redbeet (beetroot), spinach, silverbeet, poppy.

PRECAUTION

DO NOT use human flaggers unless they are protected by engineering controls such as enclosed cabs.

Re-entry period

DO NOT allow entry into treated areas for 12 hours. When prior entry is necessary, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under meteorological conditions or from spraying equipment which could be expected to cause spray to drift onto nearby susceptible plants, adjacent crops, crop lands or pastures.

**PROTECTION OF WILDLIFE, FISH,
CRUSTACEANS AND ENVIRONMENT**

Dangerous to fish and other aquatic life.
DO NOT contaminate dams, rivers, drains or streams with chemical or used containers.
DO NOT apply or allow spray to drift onto adjacent aquatic areas (ponds, streams, lakes, rivers and waterways). Allow sufficient buffer distances between downwind water bodies and the sprayed area.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight.

DO NOT dispose of undiluted chemicals on-site.

The method of disposal of the container depends on the container type. Read the "Storage and Disposal" instructions on the label that is attached to the container.

SAFETY DIRECTIONS

Will irritate the eyes. Avoid contact with eyes. Wash hands after use. When preparing and using the prepared spray wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length PVC gloves. After each day's use, wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data sheet which can be obtained from your supplier.

CONDITIONS OF SALE

Crop Care Australasia Pty Ltd will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply or use of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Crop Care Australasia Pty Ltd is limited to the replacement of the goods or (at the option of Crop Care Australasia Pty Ltd) the refund of the price paid and is conditional upon a claim being made in writing and where possible sufficient part of the goods to enable proper examination being returned to Crop Care Australasia Pty Ltd within thirty days of delivery.

APVMA Approval No. :58924/0305

* Registered trademark of Nufarm Technologies USA Pty Ltd
+ Other Trademarks



Crop Care Australasia Pty Ltd
ACN 061 362 347
77 Tingira Street
PINKENBA Qld 4008
Ph : (07) 3867 9100

In a Transport Emergency Dial 000 Police or Fire Brigade	SPECIALIST ADVICE IN AN EMERGENCY ONLY 1800 033 498 ALL HOURS- AUSTRALIA WIDE
--	---