

**POISON**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

**EverGol<sup>®</sup> Xtend**

**SEED TREATMENT**

**Active Constituent: 154 g/L PENFLUFEN**  
**154 g/L TRIFLOXYSTROBIN**

**GROUP 7 11 FUNGICIDE**

**For the control of damping off and hypocotyl rot caused by *Rhizoctonia solani* and *Fusarium* spp. in various crops as specified in the DIRECTIONS FOR USE table**

## GENERAL INSTRUCTIONS

### Fungicide Resistance Warning

EverGol Xtend is a combination of carboxinide and strobilurin fungicide. For fungicide resistance management EverGol Xtend is both a Group 7 and Group 11 fungicide. Some naturally occurring individual fungi resistant to EverGol Xtend and other Group 7 and/or Group 11 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by this product and other Group 7 and/or Group 11 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Bayer CropScience Pty Ltd accepts no liability for any losses that result from failure of EverGol Xtend to control resistant fungi.

### Application

Prior to pouring, shake the container vigorously, then add the required quantity of EverGol Xtend to sufficient water to give even coverage of the seed to be treated. Ensure that a high enough slurry volume is used to gain even coverage on each seed. Place seed in seed treatment equipment and operate until even coverage of seed is obtained. The quantity of water used for mixing will vary depending on type of equipment and type of seed, refer to Bayer for specific treatment information. Good coverage of seed can be obtained with lower slurry volumes when professional seed treatment equipment is used. The slurry mixture should be gently stirred regularly.

### Seed Quality

EverGol Xtend seed treatment should not be used on seed with more than 8% moisture content, or on sprung, sprouted, damaged or severely pinched seed, or seed of poor viability. If in doubt, have a germination test carried out on the seed before treatment to ensure that it is of acceptable standard.

Treating with EverGol Xtend as directed will raise the moisture level of the seed by up to 1%, depending on weather conditions at treatment. The use of EverGol Xtend mixed with water at recommended rates will have no effect on the storage life of treated sound seed.

### Storage of Treated Seed

Bayer recommends that treated seed be sown in the year of treatment. If the seed cannot be sown in the year of treatment it should be stored in a dry, well ventilated place. Although EverGol Xtend has no effect on the viability of treated seed, subsequent germination can be adversely affected by insect pest damage and poor storage conditions such as high moisture combined with high temperatures. It is recommended that a germination test be carried out prior to using stored seed to ensure it is still fit for purpose. No liability can be accepted for the performance of stored treated seed. For further information on storing treated seed contact the Bayer CropScience Technical Enquiry Hotline 1800 804479 or your local Bayer CropScience representative.

### Compatibility

EverGol Xtend is compatible with Jockey Stayer and Gaucho 600 FS when applied to canola and compatible with Gaucho 600 FS when applied to forage brassicas.

EverGol Xtend is compatible with Gaucho 600 FS when applied to grass and broadleaf pasture and with Apron XL when applied to lucerne and sub-clover pasture.

For the latest compatibility recommendations contact the Bayer CropScience Technical Enquiry Hotline 1800 804479 or your local Bayer CropScience representative.

### Labelling of Treated Seed

Label treated seed with the appropriate precautions using printed sacks, labels or bag tags.

**Cleaning Up**

Equipment should be thoroughly cleaned with water. In the event of changing from this water based flowable formulation to a solvent-based liquid formulation, it is essential to rinse equipment with methylated spirits after thoroughly cleaning with water. If changing from a solvent based liquid to this water based flowable, thoroughly clean with methylated spirits then rinse with water. Failure to do so will result in clogging of equipment.

**PRECAUTION**

Do not use treated seed for human or animal consumption. Do not allow treated seed to contaminate seed intended for human or animal consumption. If possible, use a separate auger for treating seed or moving treated seed. If using the same auger for treating seed or moving treated seed and moving grain for human consumption, remove all EverGol Xtend residues from the auger to avoid contaminating untreated seed with seed treatment residues.

**Re-handling**

Do not allow re-handling of treated seed unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and impervious footwear and chemical resistant gloves. Clothing must be laundered after each day's use.

**PROTECTION OF LIVESTOCK**

Seed treated with this product must not be used for animal consumption or poultry feed or mixed with animal feed. DO NOT allow seed treated with this product to contaminate seed intended for animal consumption.

**PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

Very toxic to aquatic life. DO NOT contaminate wetlands or watercourses with this product, used containers or bags which have held treated seeds. DO NOT feed treated seed or otherwise expose to wild or domestic birds. Any spillages of treated seed, however minor, must be cleaned up immediately, preferably by recovery and re-use. If disposal is required, ensure treated seeds are thoroughly buried and not accessible to birds and other wildlife.

**STORAGE AND DISPOSAL****Product and Original Container**

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Do not store near foodstuffs or animal feed.

***Non returnable containers only***

Triple rinse containers before disposal. Add rinsings to slurry in auger/mixer, or dispose of rinsings according to State/Territory legislative requirements. DO NOT dispose of undiluted chemicals on site. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product. Do not re-use empty container for any other purpose.

***Returnable containers only***

If tamper evident seals are broken prior to initial use then the integrity of the contents cannot be assured. Empty container by pumping through dry-break connection system. Do not attempt to breach the valve system or the filling point, or contaminate the container with water or other products.

Ensure that the coupler, pump, meter and hoses are disconnected, triple rinsed and drained after each use. When empty, or contents no longer required, return the container to the point of purchase. This container remains the property of Bayer CropScience Pty Ltd.

**Treated Seed and Containers of Treated Seed**

When treated seed is stored it should be kept apart from other grain and the bags or containers should be clearly marked to indicate the contents have been treated with this product. DO NOT use treated seed for human or animal consumption. Do not allow treated seed to contaminate seed intended for human or animal consumption. Bags which have held treated seed are not to be used for any other purpose.

**Application Equipment**

Rinse all equipment with clean water immediately after use and dispose of rinsings according to State/Territory legislative requirements.

**SAFETY DIRECTIONS**

Will irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. When opening the container, mixing and loading and using the product wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow length chemical resistant gloves and face shield or goggles and impervious footwear. If product on skin, immediately wash area with soap and water and in eyes, wash it out immediately with water. Wash hands after use. After each day's use wash gloves and face shield or goggles and contaminated clothing.

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone: Australia 13 11 26.

**SAFETY DATA SHEET**

Additional information is listed in the Safety Data Sheet, which can be obtained from [www.bayercropscience.com.au](http://www.bayercropscience.com.au).

**EXCLUSION OF LIABILITY**

This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty. Ltd. accepts no liability or responsibility for loss or damage arising from failure to follow such directions and instructions.

EverGol® Jockey® Stayer® and Gaucho® are Registered Trademarks of the Bayer Group

APVMA Approval No.: 66882/59071

FOR 24 HOUR SPECIALIST ADVICE  
IN EMERGENCY ONLY  
PHONE 1800 033 111

**DIRECTIONS FOR USE**

<b>CROP</b>	<b>DISEASE</b>	<b>RATE</b>	<b>CRITICAL COMMENTS</b>
Canola, Forage brassicas	Damping off and hypocotyl rot caused by <i>Rhizoctonia solani</i>  Fusarium damping off and root rot ( <i>Fusarium</i> spp.)	65 mL/100 kg seed	EverGol Xtend is applied to seed prior to sowing. Ensure even coverage of seed.  Control of <i>Rhizoctonia</i> and <i>Fusarium</i> by EverGol Xtend is characterised by improved plant emergence and survival, a reduction in hypocotyl damage and an increase in above ground biomass.
Broadleaf pasture	Rhizoctonia damping off and root rot ( <i>Rhizoctonia solani</i> )  Fusarium damping off and root rot ( <i>Fusarium</i> spp.)	35 mL /100 kg seed	EverGol Xtend will not control damping off caused by <i>Pythium</i>
Grass pasture	Rhizoctonia damping off and root rot ( <i>Rhizoctonia solani</i> )		

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS  
AUTHORISED UNDER APPROPRIATE LEGISLATION**

**WITHHOLDING PERIODS**

**Harvest: NOT REQUIRED WHEN USED AS DIRECTED**

**Grazing: Grass pastures: DO NOT GRAZE PLANTS GROWN FROM TREATED SEED, OR CUT FOR STOCK  
FOOD WITHIN 4 WEEKS OF SOWING**

**Canola, forage brassicas, broadleaf and mixed broadleaf/grass pastures: DO NOT GRAZE  
PLANTS GROWN FROM TREATED SEED, OR CUT FOR STOCK FOOD WITHIN 6 WEEKS OF  
SOWING**