

**POISON**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

**KENSO**  
*agcare*

# Ken-Zon

Herbicide

ACTIVE CONSTITUENT : 300 g/L TRICLOPYR present as butoxyethyl ester  
100 g/L PICLORAM present as hexyloxypropylamine salt

GROUP	I	HERBICIDE
-------	---	-----------

For control of a range of environmental and noxious **woody** and herbaceous weeds as specified in the Directions for Use table.

CONTENTS:  
20 Litres

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT**



**KENSO**  
*agcare*

Kenso Corporation (M) Sdn Bhd 15a/243 Bradman Street, Acacia Ridge 4110 Ph: (07) 3711 5900

NRA Approval No: 56685/0703

**STORAGE AND DISPOSAL**

Store in the closed, original container in a cool, well ventilated area. **DO NOT** store for prolonged periods in direct sunlight. **DO NOT** store near food, feedstuffs, fertilizers or seed.

Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. **DO NOT** dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

**SMALL SPILL MANAGEMENT**

Wear protective equipment (See Safety Directions). Apply absorbent material such as earth, sand, clay granules or cat litter to the spill. Sweep up material for disposal when absorption is completed and contain in a refuse vessel for disposal. (See Storage and Disposal section). If necessary, wash the spill area with an alkali detergent and water and absorb as above the wash liquid for disposal.

**SAFETY DIRECTIONS**

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When preparing the spray, wear cotton overalls buttoned to the neck and wrists, a washable hat, elbow-length PVC gloves and a face shield or goggles.

If the product is in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

**FIRST AID**

If poisoning occurs contact a doctor or Poisons Information Centre (Ph.: 131126). If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

**MATERIAL SAFETY DATA SHEET**

Additional information is listed in the Material Safety Data Sheet.

**NOTICE**

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of the product contrary to label instructions or under off-label permits not endorsed by Kenso Corporation (M) Sdn Bhd, or under abnormal conditions.

In a Transport Emergency Dial <b>000</b> Police or Fire Brigade
--

**NRA Approval No. :56685/0703**

Batch No. :

Date of Manufacture :

# POISON

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



## Ken-Zon

Herbicide

**Active Constituent : 300 g/L TRICLOPYR present as butoxyethyl ester  
100 g/L PICLORAM present as  
hexyloxypropylamine salt**



For control of a range of environmental and noxious woody and herbaceous weeds as specified in the Directions for Use table.

**READ THIS LEAFLET BEFORE USE**

**SHAKE WELL BEFORE USE**

**NRA Approval No : 56685/0703**



**Kenso Corporation (M) Sdn Bhd**  
15a/243, Bradman Street, Acacia Ridge, 4110  
Phone 07 3711 5900

**DIRECTIONS FOR USE****RESTRAINTS**

**DO NOT** apply to weeds which may be stressed (not actively growing) due to prolonged periods of extreme cold, moisture stress (water-logged or drought affected), poor nutrition, presence of disease, damage or previous herbicide treatment, as reduced levels of control may result.

**DO NOT** spray if rain is likely within one hour or if foliage is wet from rain or dew. .

**DO NOT** burn off, cut or clear blackberry or other woody weeds for at least six months after spraying.

**DO NOT** apply by aerial application in wind in excess of 15 km/hr and air temperatures above 35°C.

In areas prone to flooding treatment should commence after any annual flooding as such areas flooded within 9 months following application may have reduced results.

**1. WOODY WEED SITUATIONS****Table A : High Volume Spraying**

See **GENERAL INSTRUCTIONS – APPLICATION** section for application method details.

<b>AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY.</b>				
<b>WEEDS CONTROLLED</b>	<b>WEED GROWTH STAGE</b>	<b>STATE</b>	<b>RATE /100 L Water</b>	<b>CRITICAL COMMENTS</b>
African boxthorn	Less than 2 m tall	All States	500 mL	Apply when bushes have good leaf cover, growth and <b>no leaf fall</b> .
<i>Angophora</i> Spp.	1 to 3 m tall		350 mL	
Australian blackthorn	Less than 2 m tall		500 mL	Apply from late spring to early autumn.
<i>Banksia</i> Spp.	1 to 3 m tall		350 mL	
Biddy bush (Chinese shrub) (Sifton bush)	Autumn when actively growing	ACT and NSW only	500 mL	Add a 100% concentrate non-ionic surfactant (e.g. as Kenso Agcare Ken-Wett 1000) at 125 mL/100 L of water for best results.
Blackberry in association with: Docks Ragwort Smartweed Thistles	Late spring to autumn	All States	350 mL OR 500 mL	<b>Use the higher rate on plants that have been damaged by grazing stock or insects and on known difficult to kill blackberry.</b> Where herbicides other than Group I herbicides have been used, allow two seasons regrowth to occur before respraying with Ken-Zon.
Blue heliotrope	Flowering	NSW and Qld only	500 mL	Apply in a minimum spray volume of 1250 L/ha.
Brooms: Cape English Flax leaf Montpellier	Spring to mid-summer prior to pod formation Autumn to winter	All States	250 mL	Apply as a thorough foliage spray.
Camphor laurel	Less than 2 m tall		350 mL	
	Above 2 m tall		500 mL	
<i>Casuarina</i> Spp.	1 to 3 m tall		350 mL	
Chinese apple	Less than 2 m tall	Qld and WA only	350 mL	Add a 1 00% concentrate non-ionic surfactant (e.g. Kenso Agcare Ken-Wett 1000) at 100 mL/100 L of water for best results.
Cockspur thorn	Spring to autumn	ACT, NSW and Qld only		Apply as a thorough foliage spray.

Table A : High Volume Spraying

See GENERAL INSTRUCTIONS – APPLICATION section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY.				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE /100 L Water	CRITICAL COMMENTS
Common sensitive plant	Any time when actively growing	NT, Qld and WA only	200 mL	To avoid leaves closing during application, spray plants while moving forward. Add a 100% concentrate non-ionic surfactant (e.g. Kenso Agcare Ken-Wett 1000) at 100 mL/100 L of water for best results.
Crofton weed	Spring to autumn	ACT, NSW and Qld only	350 mL	Apply as a thorough foliage spray.
Eastern cotton bush ( <i>Maireana microphylla</i> )	Spring to autumn	NSW and Qld only	500 mL	Add Uptake Spraying Oil at 500 mL/100 L of water. Some bushes may require a follow-up spray to control regrowth.
<i>Eucalyptus</i> spp.	Seedling and regrowth from small lignotubers, 1 to 3 m tall	All States	350 mL OR 500 mL	<b>Apply the high rate where difficult to control species of <i>Eucalyptus</i> regrowth is present.</b> Addition of an adjuvant may improve results - contact Kenso Agcare for details.
Galenia	Fresh growth in spring to summer	NSW only	500 mL	Use 2000 L of spray mixture/ha.
Giant bramble	Spring to autumn	NT, Qld and WA only		Penetration of thick clumps may be difficult and respraying may be necessary. Add a 100% concentrate non-ionic surfactant (e.g. Kenso Agcare Ken-Wett 1000) at 100 mL/100 L of water for best results.
Gorse	1 to 1.5 m tall	All States	250 mL	Spring and Summer treatment only. Add a 100% concentrate non-ionic surfactant (e.g. Kenso Agcare Ken-Wett 1000) at 100 mL/100 L of water for best results.
	Over 1.5 m tall or autumn treatment		350 mL	
	Winter treatment		500 mL	
Groundsel bush ( <i>Baccharis halimifolia</i> )	1 to 1.5 m tall in spring to summer	All States	250 mL	Apply as a thorough foliar spray.
	Over 1.5 m tall or autumn treatment		350 mL	
Green cestrum	Late spring to early autumn	ACT, NSW and Qld only	500 mL	One application may give satisfactory control. Any subsequent regrowth and seedlings must be resprayed at approximately 1 metre high.
Hawthorn	Less than 2 m tall	All States		Apply from late spring to early autumn.
Horehound	Pre-flowering	NSW and Qld only	350 mL	Apply as a thorough foliar spray.
Japanese sunflower				

Table A : High Volume Spraying

See GENERAL INSTRUCTIONS – APPLICATION section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY.				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE /100 L Water	CRITICAL COMMENTS
Lantana ( <i>Lantana camara</i> ) ( <i>Lantana montevidensis</i> )	Up to 1 m tall in summer to autumn	All States	350 mL	Add one of the following adjuvants, when using 350 mL rate: Uptake* Spraying Oil @ 0.5% v/v. Pulse® Penetrant @ 0.1% v/v.
	1 to 2 m tall in summer to autumn		500 or 750 mL	Thoroughly wet foliage, stems and soil around the base of the plants. Use higher rate on known harder to kill varieties.
Lion tail ( <i>Leonatis nepetifolia</i> )	Pre-flowering	Qld only	200 mL	Apply as a thorough foliar spray. Add a 100% concentrate non-ionic surfactant (e.g. Kenso Agcare Ken-Wett 1000) at 100 mL/100 L of water for best results.
Limebush	Any time of year with good leaf cover and soil moisture	NT and Qld only	350 mL	Penetration of thick clumps may be difficult and respraying may be required. Addition of an adjuvant may improve results - contact Kenso Agcare for details.
Manuka	At flowering	Vic only	500 mL	For optimum results, add Pulse Penetrant at 200 mL/100 L of spray. Thoroughly wet foliage, stems and soil around the base of the plants.
Mesquite ( <i>Prosopis</i> spp.)	Seedling, full leaf and flowering before podding	NSW, NT, Qld and WA only	350 mL	<b>DO NOT</b> spray plants bearing pods. Add a 100% concentrate non-ionic surfactant (e.g. Kenso Agcare Ken-Wett 1000) at 100 mL/100 L of water for best results.
<i>Prosopis velutina</i>		Qld only	670 mL	
Mistflower	Spring to Autumn	ACT, NSW and Qld only		Apply as a thorough foliar spray.
Mother-of-millions	Flowering	NSW and Qld only	500 mL	Add a 100% concentrate non-ionic surfactant (e.g. Kenso Agcare Ken-Wett 1000) at 100 mL/100 L of water.
Paddy's lucerne.	Active growth	NSW only		Plants that have been continually slashed or grazed over many seasons may be difficult to control and regrowth may occur.
Parkinsonia	Under 2 m tall	NT, Qld and WA only	350 mL	Add Uptake Spraying Oil @ 500 mL/100 L water. Avoid spraying under dry conditions when plants are stressed or bearing pods. Thoroughly wet foliage.
Prickly pear (common) Smooth tree pear	Active phyllode growth	All States	500 mL	Apply as a thorough foliage spray. Regrowth may occur, so a follow-up application may be necessary.

Table A : High Volume Spraying

See GENERAL INSTRUCTIONS – APPLICATION section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY.				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE /100 L Water	CRITICAL COMMENTS
Rubber vine (Not infected with rust)	Up to 1.5 m tall at flowering	NT, Qld and WA only	350 mL	Spray all leaves and stems just to the point of runoff and thoroughly spray the base of the plant. With faroer, more dense stands, regrowth may occur. Subsequent control of any regrowth should be done by basal bark spraying.
	Dense stands greater than 1.5 m tall at flowering		500 mL	
Siam weed	Active growth	Qld and WA only	350 mL	Add a 100% concentrate non-ionic surfactant (e.g. Kenso Agcare Ken-Wett 1000) at 100 mL/100 L of water for best results.
Sicklepod	Up to flowering	NT, Qld and WA only	200 mL	<b>DO NOT</b> apply to podding plants. Add a 100% concentrate non-ionic surfactant (e.g. Kenso Agcare Ken-Wett 1000) at 100 mL/100 L of water for best results.
St John's wort	From flowering to early seed set	All states	500 mL	Late spring to early summer.
Sweet briar	Up to 1.5 m tall	All States	350 mL	Add metsulfuron-methyl at 10 g/100 L water to obtain more reliable results with the lower rate of Ken-Zon.
			500 mL	Full leaf to ripe fruit prior to leaf fall. Thorough wetting including the crown is recommended.
Tobacco weed	Actively growing plants	NT, Qld and WA only	300 mL	Add a 100% concentrate non-ionic surfactant (e.g. Kenso Agcare Ken-Wett 1000) at 100 mL/100 L of water for best results.
Wattle ( <i>Acacia</i> spp.) (except corkwood wattle)	1 to 3 m tall	All States	350 mL	
Wild rosemary ( <i>Cassinia laevis</i> )	Active growth, 0.5 to 1.0 m tall	Qld only	350 to 500 mL	Use lower rate on seedlings 0.5 m tall. Apply as a thorough foliar spray.
Wild tobacco tree	Spring to autumn up to 2 m tall	ACT, NSW and Qld only	350 mL	

Table B : Aerial Application

See GENERAL INSTRUCTIONS - APPLICATION section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, PASTURES AND RIGHTS-Of-WAY.				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE /ha	CRITICAL COMMENTS
Blackberry	Summer to autumn	NSW, Qld, SA, Vic and WA only	10 L	Where herbicides other than Group I herbicides have been used, allow two seasons regrowth to occur before respraying with Ken-Zon. <b>WARNING:</b> <i>Eucalyptus</i> species up to 4 m may be killed if sprayed during this treatment. Mature trees which are 15 to 20 m tall may be partially defoliated but are likely to recover.
Gorse		Tas only		
Cockspur thorn Crofton weed Lantana Mistflower	Late autumn	NSW, NT and Qld only (helicopter only)	1.5 L plus 7.5 L Ken-Amine 500	Spray with calibrated equipment using the half overlap opposite pass technique applying a minimum spray volume of 150 L/ha. <b>Follow-up respraying will be required.</b>
Lantana			10 L	
Rubber vine (Not infected with rust)	When flowering	NT and Qld only (helicopter only)	3 L to 5 L	Use rates will depend upon the density and height of the rubber vine stand. The higher rate should be used on dense stands, however, complete coverage and penetration may be difficult. <b>Follow-up respraying will be required.</b> Any regrowth should be sprayed with a suitable basal bark herbicide.
St John's wort	Flowering to early seed set (Nov-Jan)	NSW only	4 L	Helicopter application only. Follow-up spraying will be required in the following season.
AGRICULTURAL NON-CROP AREAS ON FLOOD PLAINS.				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE /ha	CRITICAL COMMENTS
Parkinsonia	Seedlings 1-2 m tall, or 12 - 24 months old	Qld and NT only (helicopter only)	3L	Add Uptake Spraying Oil at 1 L/ha.

Table C: Controlled Droplet Application (C.D.A.)

See GENERAL INSTRUCTIONS - APPLICATION section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-Of-WAY.				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE /ha	CRITICAL COMMENTS
Blackberry in association with: Docks Ragwort St John's wort Thistles	Summer to autumn	All States	Apply undiluted	One application may give satisfactory control but subsequent regrowth and seedlings should be re-sprayed after hardening off. Where herbicides other than Group I herbicides have been used, allow two seasons regrowth to occur before respraying with Ken-Zon.



**Table D : Low Volume High Concentrate Application Techniques  
(Gas Powered Gun, Sprinkler Sprayer)**

See **GENERAL INSTRUCTIONS. APPLICATION** section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY.					
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE /10 L Water	CRITICAL COMMENTS	
Blackberry	Late spring to autumn	ACT, NSW, Qld, SA, Tas and WA only	335 mL	Apply to actively growing bushes which are able to be sprayed on all sides. For larger bushes, the high volume application technique is recommended.	
Camphor laurel Cockspur thorn Crofton weed	Less than 1.5 m high	ACT, NSW and Qld only	500 mL		
<i>Eucalyptus</i> species	Seedlings up to 2 m tall	All States	335 mL		
Mistflower	Less than 1.5 m high	ACT, NSW and Qld only	500 mL		
Sweet briar	1.5 m tall, full leaf to ripe fruit	NSW only			<b>Gas Powered Gun only:</b> Apply to actively growing bushes not more than 1.5 m tall that have not more than 5 stems from the crown.
St John's wort	During flowering to early seed set	NSW, Tas and Vic only			<b>Gas Powered Gun only:</b> One application should provide control. Minor regrowth and seedlings may be retreated the following summer.
Wild tobacco tree	Less than 1.5 m high	ACT, NSW and Qld only			Apply to actively growing bushes which are able to be sprayed on all sides. For larger bushes, the high volume application technique is recommended.

**Table E: Boom Application**

See **GENERAL INSTRUCTIONS. APPLICATION** section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, PASTURES AND RIGHTS-OF-WAY.				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE /ha	CRITICAL COMMENTS
Galenia	Fresh growth during spring to summer	NSW only	5L	Rough mine sites will require adequate spray equipment such as boomless nozzles for effective coverage.
Sicklepod	Up to flowering	NT and Qld only	3L	<b>DO NOT</b> apply to podding plants. Add a 100% concentrate non-ionic surfactant (e.g. Kenso Agcare Ken-Wett 1000) at 100 mL/100 L of water.
St John's wort	Flowering to early seed set (Nov-Jan)	NSW only	2 to 4 L	Use the higher rate on dense infestations and when longer residual control is required. Follow-up respraying will be required in the following season.

## 2. FALLOW SITUATIONS

**Table A: Boom Application**

See **GENERAL INSTRUCTIONS - APPLICATION** section for application method details.

FALLOW				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE /ha	CRITICAL COMMENTS
Blackberry nightshade - <b>Suppression only</b>	10 to 25 cm tall, prior to flowering	NSW and Qld only	200 to 400 mL + 1.2 L Ken-Up 450 CT + adjuvant	<p><b>FOR USE BY GROUND EQUIPMENT ONLY.</b></p> <p>Plants must be actively growing.</p> <p>Use the lower rate on the smaller weeds, as specified in the weed growth stage (or up to 5 cm diameter for <i>Polymeria pusilla</i>).</p> <p>Refer to Ken-Up 450 CT label for use of adjuvant.</p> <p><b>DO NOT</b> plant susceptible crops for up to nine months after application, as specified in General Instructions - Minimum Recropping Periods – Black Cracking Clay Soils, NNSW &amp; Qld.</p> <p>Dry conditions after application will increase the recropping interval.</p>
Camel melon Prickly paddy melon Cucumber melon ( <i>Cucumis melo</i> )	From 2 leaf to 50cm diameter			
Common sowthistle	From 8 leaf to flowering			
Cow vine	From 2 to 5 leaf up to 15 cm diameter, prior to flowering			
Lucerne (established)	Active growth, 15 to 25 cm high, during spring			
<i>Polymeria pusilla</i>	2 to 12 leaf up to 20 cm diameter, prior to flowering		300 to 500 mL + 1.2 L Ken-Up 450 CT + adjuvant	
			200 to 400 mL + 1.2 L Ken-Up 450 CT + adjuvant	

**Table B : Blanket Wiper Application**

See **GENERAL INSTRUCTIONS - APPLICATION** section for application method details.

FALLOW				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE	CRITICAL COMMENTS
Bitterbark ( <i>Alstonia constricta</i> )	From summer to end of autumn	Qld	1:4 (1 part Ken-Zon to 4 parts water)	<p>For use with blanket wipers only.</p> <p>For best results apply in autumn to tall (&gt; 60 cm) plants using two opposite directional passes (up and back).</p> <p>Follow up "missed" plants with a spot spray application. These will be obvious after 6 weeks.</p> <p>Blanket wiper applications can be made in summer when plants are smaller but follow up spot spraying may be necessary.</p> <p>Do not disturb (cultivate) the treated patches for at least 3 months. Best long term control is achieved when patches are left undisturbed for as long as possible after treatment (at least 6 months).</p> <p><b>Spot spraying "missed" plants:</b> thoroughly wet all stems and leaves without producing any solution runoff. Avoid any spray reaching the soil surface.</p>
			2 % solution for spot spray (eg. 100 mL Ken-Zon in 5 L water)	

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**IN TASMANIA FOR BLACKBERRY**

DO NOT treat bushes carrying mature or near mature fruit.

**FOR NATIVE VEGETATION**

Use of Ken-Zon on native vegetation must be done in accordance with STATE and/or LOCAL legislation.

**WITHHOLDING PERIOD**

NOT REQUIRED WHEN USED AS DIRECTED.

**GENERAL INSTRUCTIONS**

**MINIMUM RECROPPING PERIODS - Black Cracking Clay Soils, NNSW & Qld.**

**A: Boom Application**

Plant-back periods for crops following the application of Ken-Zon for rates up to 600 mL/ha.				
RATE mL/HA	200	300	400	600
CROP	Months			
Wheat	2	2	4	4
Barley	2	2	4	4
Canola	2	4	4	4
Faba bean	4	4	6	6
Chickpea	4	6	6	6
Lucerne	6	9	9	9

These plantback periods are based on a normal rainfall pattern. During drought conditions (or when the rainfall is less than 100 mm for a period of 4 months or greater) the plantback period may be significantly longer.

**Table B : Blanket Wiper Application**

Plant-back periods for crops following Blanket wiper application	
CROP	Months
Broadleaf crops	18
Lucerne	6
	This will allow any potential soil residues to dissipate. if any, and allow effective control.

**Note:** Before using Ken-Zon in tank mixes with other herbicides, check the plant-back information on all product labels. The most residual product, i.e. the product with the longest plant-back period, will determine the time between spraying and planting.

**COMPATIBILITY**

Follow any regional restrictions, and all directions and restrictions on the label, of any chemical mixed with Ken-Zon(e.g. 2,4-D amine).

Ken-Zon is compatible with the following herbicides:

2,4-D amine	DP <sup>®</sup> 600	metsulfuron-methyl
Ken-Up 450 CT	Roundup CT Xtra	Starane* 200

Ken-Zon is compatible with the following adjuvants, as per Directions for Use:

Kenso Agcare Ken-Wett 1000	Pulse <sup>®</sup> Penetrant	Uptake <sup>®</sup> Spraying Oil
----------------------------	------------------------------	----------------------------------

**MIXING****Mix only with water.**

Half fill the spray unit with water, and add the required amount of Ken-Zon. Add the remaining water with the agitator running. If required, then add spray oils or wetters (surfactants). Maintain mechanical or by-pass agitation in the spray tank during spraying. Only mix sufficient solution for immediate daily use and avoid storing.

**APPLICATION****1. WOODY WEED SITUATIONS**

**Weeds need to be actively growing for herbicides to have optimum effect. Delay treatment until all regrowth has had time to grow to approximately 1 metre in situations which have been bulldozed, slashed, burnt, ploughed or areas having a previous chemical treatment.**

**A. High Volume Spraying**

- Thorough coverage of foliage to the point of run-off is essential, however, avoid excess spraying which is wasteful of chemical.

**Hand Gun**

- Apply the recommended mix to give full coverage of leaves and stems through a No.6 to 8 tip at 700 to 1500 kPa (400 to 500 kPa for St John's wort).
- A spray volume of 3000 to 4000 L per infested hectare of 1 to 2 metre high blackberry (30 to 40 L/100 m<sup>2</sup>) should be used.
- Use 2000 L of spray mixture/ha of galenia infestation (Le. 20 L/100 m<sup>2</sup> infested area).

**Knapsack**

- Apply the recommended spray mix to give full coverage of leaves and stems. The final volume of application should be similar to hand gun.
- A spray volume of 3 to 4 L/10 m<sup>2</sup> infested area should be used.
- A spray volume of 2 L/10 m<sup>2</sup> should be, used for an area infested with galenia.

**B. Aerial Application**

- Apply in 200 L of water/ha using an aircraft to apply 100 L per pass on a double overlap pattern using nozzle configurations to produce droplets of 250 to 350 micron diameter.
- The potential for damage from drift can be greatly reduced by avoiding unsuitable spraying conditions and using spray pressure and nozzles to minimise the production of small droplets.
- **DO NOT** spray when wind exceeds 15 km/hr and/or air temperature reaches 35°C.

**C. Controlled Droplet Application (C.D.A.)**

- Results similar to high volume spraying can be obtained using Micron Herbi<sup>®</sup> or similar equipment. Select a nozzle to give a flow rate of 2 mL/sec and sweeping action of approximately 1 m/sec to ensure a droplet density of 20/cm<sup>2</sup>. Use a marking agent, as recommended by the equipment manufacturers, to check spray coverage. Also, consult directions provided with C.D.A. unit.

**D. Low Volume High Concentrate Application Techniques**

- Good control will be achieved, similar to high volume application, where bush size enables good coverage of entire bush. Use a marking agent, as recommended by the equipment manufacturers, to check spray coverage.
- **Gas powered gun:** Apply 50 mL shots to obtain uniform coverage of 4 to 5 m<sup>2</sup> of surface area of bush. This relates to 20 droplets/cm<sup>2</sup> of leaf surface.
- **Sprinkler sprayer:** This technique involves using a micro sprinkler that is connected to a hollow fibre glass rod attached to a pneumatic knapsack sprayer. Use at low pressures (50 to 200 kPa) and apply with a slow sweeping action over the top of the plants ensuring even coverage on the leaves.

**E. Boom Application**

- Application in a minimum spray volume of 200 L/ha for galenia and St. John's wort and 600 L of water/ha for sicklepod. Flat fan nozzles are recommended, using pressure in the range of 200-300 kPa. Boom height must be set to ensure double overlap of nozzle patterns.

## 2. FALLOW SITUATIONS

### A. Boom Application

- Application of Ken-Zon in a minimum spray volume of 50 L/ha is recommended. Flat fan nozzles are recommended, using pressure in the range of 200-300 kPa. Boom height must be set to ensure double overlap of nozzle patterns.

### B. Blanket Wiper Application

- Blanket needs to be made from durable and wettable material with a rigid backing.
- Blanket should be rigidly mounted behind motorised vehicle (eg. tractor, 4-wheel drive vehicle) and set low but never touching the ground. The chemical solution should be fed to the blanket at a flow rate sufficient to keep the blanket wet but not dripping. In thick patches the blanket may require more frequent solution recharge (rewetting).
- Ideally, a scraper bar should be mounted in front of the blanket in order to scrape or damage the bark (but not sever the stems) prior to the blanket wiping the stems. This scraper may be mounted at the front of the vehicle.
- Two passes (in opposite direction) with the blanket increases the contact with the plant. Ground speeds of 10 -15 kph are ideal for blanket wiping application.

## CLEANING SPRAY EQUIPMENT

### Rinsing

- After using Ken-Zon, empty the spray unit completely and drain the whole system. Thoroughly wash inside the unit using a pressure hose. Drain the spray unit, and clean any filters in the tank, pump, lines, hoses and nozzles.
- After cleaning the spray unit as above, quarter fill with clean water and circulate through the pump, lines, hoses and nozzles. Drain and repeat the rinsing procedure twice.

### Decontamination

- Before spraying cotton and other sensitive crops with equipment that has been used to apply Ken-Zon, see PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS section.
- Wash the tank and rinse the system as above. Then quarter fill the tank and add an alkali detergent (e.g. liquid SURF<sup>®</sup>, OMO<sup>®</sup>, OMOMATIC<sup>®</sup>, DRIVE<sup>®</sup> at 500 mL/100 L of water or the powder equivalent at 500 g/100 L of water) and circulate throughout the system for at least 15 minutes.
- Drain the whole system. Remove filters and nozzles and clean them separately. Finally flush the system with clean water and allow to drain.

**Rinse water should be discharged onto a designated disposal area or, if this is unavailable, onto unused land away from desirable plants and watercourses.**

## RESISTANT WEEDS WARNING

GROUP	I	HERBICIDE
-------	---	-----------

Ken-Zon Herbicide is a member of the pyridines group of herbicides. The product has the disrupters of plant cell growth mode of action. For weed resistance management, the product is a Group I herbicide.

Some naturally-occurring weed biotypes resistant to the product and other disrupters of plant cell growth herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by the product or other disrupters of plant cell growth herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Kenso Corporation (M) Sdn Bhd accepts no liability for any losses that may result from the failure of the product to control resistant weeds. Strategies to minimise the risk of herbicide resistance are available. Contact your farm chemical supplier, consultant, local Department of Agriculture, or local Kenso Corporation (M) Sdn Bhd representative.

## PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

- Crops susceptible to Ken-Zon include, but are not limited to: peas, lupins, lucerne, navy beans, peanuts, soybeans and other legumes, cotton, flowers, fruit, hops, ornamentals, shade trees and *Pinus* spp., potatoes, safflower, sugar beet, sunflowers, tobacco, tomatoes, vegetables and vines.
- Ken-Zon is damaging to susceptible crops during both growing and dormant periods. Grasses are normally unaffected and establish quickly after treatment.
- Picloram, one of the active constituents in this product can remain active in the soil for extended periods depending on soil type and application rate, rainfall, temperature, humidity, soil moisture and soil organic matter.

- **DO NOT** apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands, pastures, waterways or native vegetation.
- **DO NOT** allow physical spray drift onto waterways, native vegetation and susceptible crops.
- **DO NOT** apply close to, or in areas, containing roots of desirable vegetation, where treated soil may be washed onto areas growing (or areas to be planted with) desirable plants.
- **DO NOT** apply on sites where surface water from heavy rain can be expected to run off to areas containing, or to be planted with susceptible crops or plants. **DO NOT** move soil, which may have been treated to areas where desirable plants are to be grown.

#### **PROTECTION OF LIVESTOCK**

- Poisonous plants may become more palatable after spraying and stock should be kept away from these plants until they have died down.
- Many plants remain poisonous after death, and stock should not be allowed access it, as there is a likelihood that they may graze the dead material. Such material should be burnt if possible.

#### **PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

- **DO NOT** contaminate streams, rivers or waterways with the chemical or used containers.

#### **STORAGE AND DISPOSAL**

Store in the closed, original container in a cool, well ventilated area. **DO NOT** store for prolonged periods in direct sunlight. **DO NOT** store near food, feedstuffs, fertilizers or seed.

Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. **DO NOT** dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

#### **SMALL SPILL MANAGEMENT**

Wear protective equipment (See Safety Directions). Apply absorbent material such as earth, sand, clay granules or cat litter to the spill. Sweep up material for disposal when absorption is completed and contain in a refuse vessel for disposal. (See Storage and Disposal section). If necessary, wash the spill area with an alkali detergent and water and absorb as above the wash liquid for disposal.

#### **SAFETY DIRECTIONS**

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When preparing the spray, wear cotton overalls buttoned to the neck and wrists, a washable hat, elbow-length PVC gloves and a face shield or goggles.

If the product is in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

#### **FIRST AID**

If poisoning occurs contact a doctor or Poisons Information Centre (Ph.: 131126). If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

#### **MATERIAL SAFETY DATA SHEET**

Additional information is listed in the Material Safety Data Sheet.

#### **NOTICE**

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of the product contrary to label instructions or under off-label permits not endorsed by Kenso Corporation (M) Sdn Bhd, or under abnormal conditions.

In a Transport  
Emergency Dial  
**000**  
Police or Fire Brigade



**Kenso Corporation (M) Sdn Bhd**  
15a/243, Bradman Street, Acacia Ridge, 4110  
Phone 07 3711 5900

**NRA Approval No. : 56685/0703**