

## **POISON**

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

# **GALLOP**

Herbicide

**ACTIVE CONSTITUENTS: 300 g/L TRICLOPYR present as the butoxyethyl ester.  
100 g/L PICLORAM present as hexyloxypropylamine salt.**

**GROUP II HERBICIDE**

For the control of environmental and noxious woody and herbaceous weeds  
as specified in the Directions for Use Table.

**IMPORTANT: READ THE ATTACHED LEAFLET  
BEFORE USING THIS PRODUCT.  
SHAKE WELL BEFORE USE.**

Contents: 1 Litre  
5 Litres  
10 Litres  
20 Litres

Crop Care Australasia Pty Ltd, Unit 15/16 Metroplex Avenue, Murarrie QLD 4172.

**GENERAL INSTRUCTIONS****STORAGE AND DISPOSAL**

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should NOT be burnt.

**SMALL SPILL MANAGEMENT**

Wear protective equipment (See Safety Directions). Apply absorbent material such as earth, sand, clay granules or cat litter to the spill. Sweep up material for disposal when absorption is completed and contain in a refuse vessel for disposal. (See Storage and Disposal section). If necessary, wash the spill area with an alkali detergent and water and absorb as above the wash liquid for disposal.

**SAFETY DIRECTIONS**

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When preparing spray wear cotton overalls buttoned to the neck and wrists, a washable hat, elbow-length PVC gloves and a face shield or goggles. If product is in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

**MATERIAL SAFETY DATA SHEET**

Additional information is listed in the Material Safety Data Sheet which can be obtained from your supplier.

**CONDITIONS OF SALE**

Crop Care Australasia Pty Ltd will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply or use of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Crop Care Australasia Pty Ltd is limited to the replacement of the goods or (at the option of Crop Care Australasia Pty Ltd) the refund of the price paid and is conditional upon a claim being made in writing and where possible sufficient part of the goods to enable proper examination being returned to Crop Care Australasia Pty Ltd within thirty days of delivery.

In a Transport Emergency Dial <b>000</b> Police or Fire Brigade	SPECIALIST ADVICE IN AN EMERGENCY ONLY <b>1800 033 498</b> ALL HOURS- AUSTRALIA WIDE
--	---



Crop Care Australasia Pty Ltd  
ACN 061 362 347  
Unit 15/16 Metroplex Avenue  
Murarrie Qld 4172  
Ph : (07) 3909 2000

APVMA Approval No: 61664/1L/0407  
APVMA Approval No: 61664/5L/0407  
APVMA Approval No: 61664/10L/0407  
APVMA Approval No: 61664/20L/0407

**In case of emergency: Phone 1800 033 498**  
Ask for shift supervisor. Toll free 24 hours.

Batch No:

Date of Manufacturer:

Barcode

DrumMuster logo

## **POISON**

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

# **GALLOP**

Herbicide

**ACTIVE CONSTITUENTS: 300 g/L TRICLOPYR present as the butoxyethyl ester.  
100 g/L PICLORAM present as hexyloxypropylamine salt.**

**GROUP II HERBICIDE**

For the control of environmental and noxious woody and herbaceous weeds  
as specified in the Directions for Use Table.

**IMPORTANT: READ THOROUGHLY BEFORE USE.  
SHAKE WELL BEFORE USE.**

APVMA Approval No.: 61664/0407

Crop Care Australiasia Pty Ltd, Unit 15/16 Metroplex Avenue, Murarrie QLD 4172.

**DIRECTIONS FOR USE****RESTRAINTS:**

DO NOT apply to weeds which may be stressed (not actively growing) due to prolonged periods of extreme cold, moisture stress (water-logged or drought affected), poor nutrition, presence of disease, damage or previous herbicide treatment, as reduced levels of control may result.

DO NOT spray if rain is likely within one hour or if foliage is wet from rain or dew.

DO NOT burn off, cut or clear blackberry or other woody weeds for at least six months after spraying.

DO NOT apply by aerial application in wind in excess of 15km/hr and air temperatures above 35°C.

In areas prone to flooding, treatment should commence after any annual flooding as such areas flooded within 9 months following application may have reduced results.

**1. WOODY WEED SITUATIONS****Table A: High Volume Spraying**

See **GENERAL INSTRUCTIONS - APPLICATION** section for application method details.

<b>AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHT-OF-WAYS</b>				
<b>WEEDS CONTROLLED</b>	<b>WEED GROWTH STAGE</b>	<b>STATE</b>	<b>RATE /100L WATER</b>	<b>CRITICAL COMMENTS</b>
<b>African boxthorn</b>	Less than 2 m tall	All States	500 mL	Apply when bushes have good leaf cover, growth and no leaf fall.
<i>Angophora</i> spp.	1 to 3 m tall		350 mL	
<b>Australian blackthorn</b>	Less than 2 m tall		500 mL	Apply from late spring to early autumn.
<i>Banksia</i> spp.	1 to 3 m tall	350 mL		
<b>Biddy bush (Chinese shrub) (Sifton bush)</b>	Autumn when actively growing	ACT and NSW only	500 mL	Add Chemwet 1000 at 125 mL/100 L of water for best results.
<b>Blackberry in association with: Docks Ragwort Smartweed Thistles</b>	Late spring to autumn	All States	350 mL or 500 mL	Use the higher rate on plants that have been damaged by grazing stock or insects and on known difficult to kill blackberry. Where herbicides other than Group I herbicides have been used, allow two seasons regrowth to occur before respraying with Gallop.
<b>Blue heliotrope</b>	Flowering	NSW and Qld only	500 mL	Apply in a minimum spray volume of 1250 L/ha
<b>Brooms: Cape English Flax leaf Montpellier</b>	Spring to mid-summer prior to pod formation	All States	250 mL	Apply as a thorough foliage spray
	Autumn to winter		350 mL	
<b>Camphor laurel</b>	Less than 2 m tall	All States	500 mL	
	Above 2 m tall			
<i>Casuarina</i> spp.	1 to 3 m tall		350 mL	
<b>Chinee apple</b>	Less than 2 m tall	Qld and WA only	350 mL	Add Chemwet 1000 at 100 mL/100 L of water for best results.
<b>Cockspur thorn</b>	Spring to Autumn	ACT, NSW and Qld only		Apply as thorough foliage spray
<b>Common sensitive plant</b>	Any time when actively growing	NT, Qld and WA only	200 mL	To avoid leaves closing during application, spray plants while moving forward. Add Chemwet 1000 at 100 mL/100 L of water for best results.
<b>Crofton weed</b>	Spring to autumn	ACT, NSW and Qld only	350 mL	Apply as thorough foliage spray.

**DIRECTIONS FOR USE:** continued**Table A: High Volume Spraying**See **GENERAL INSTRUCTIONS - APPLICATION** section for application method details.

<b>AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHT-OF-WAYS</b>				
<b>WEEDS CONTROLLED</b>	<b>WEED GROWTH STAGE</b>	<b>STATE</b>	<b>RATE /100L WATER</b>	<b>CRITICAL COMMENTS</b>
<b>Eastern cotton bush</b> <i>(Maireana microphylla)</i>	Spring to autumn	NSW and Qld only	500 mL	Add Uptake* Oil at 500 mL/100 L of water. Some bushes may require a follow up spray to control regrowth.
<b>Eucalyptus spp.</b>	Seedling and regrowth from small lignotubers, 1 to 3 m tall	All States	350 mL or 500 mL	Apply the high rate where difficult to control species of Eucalyptus regrowth is present. Addition of an adjuvant may improve results- contact Crop Care Australasia Pty Ltd for details.
<b>Galenia</b>	Fresh growth in spring to summer	NSW only	500 mL	Use 200 L of spray mixture/ha
<b>Giant bramble</b>	Spring to autumn	NT, Qld and WA only		Penetration of thick clumps may be difficult and respraying may be necessary. Add Chemwet 1000 at 100 mL/100 L of water for best results.
<b>Gorse</b>	1 to 1.5 m tall	All States	250 mL	Spring and Summer treatment only. Add Chemwet 1000 at 100 mL/100 L of water for best results.
	Over 1.5 m tall or autumn treatment		350 mL	
	Winter treatment		500 mL	
<b>Groundsel bush</b> <i>(Baccharis halimifolia)</i>	1 - 1.5 m tall in spring to summer	All States	250 mL	Apply as a thorough foliar spray.
	Over 1.5 m tall or autumn treatment		350 mL	
<b>Green cestrum</b>	Late spring to early autumn	ACT, NSW and Qld only	500 mL	One application may give satisfactory control. Any subsequent regrowth and seedlings must be resprayed at approximately 1 metre high.
<b>Hawthorn</b>	Less than 2 m tall	All States		Apply from late spring to early autumn.
<b>Horehound</b>	Pre- flowering	NSW and Qld only	350 mL	Apply as a thorough foliar spray
<b>Japanese sunflower</b>				
<b>Lantana</b> <i>(Lantana camara)</i> <i>(Lantana montevidensis)</i>	Up to 1 m tall in summer to autumn	All States		Add one of the following adjuvants, when using 350 mL rate: A crop oil @ 0.5% v/v. Pulse^^ Penetrant @ 0.1% v/v.
	1 to 2 m tall in summer to autumn		500 or 750 mL	Thoroughly wet foliage, stems and soil around the base of the plants. Use higher rate on known harder to kill varieties.
<b>Lion Tail</b> <i>(Leonotis nepetifolia)</i>	Pre-flowering	Qld only	200 mL	Apply as a thorough foliar spray. Add Chemwet 1000 at 100 mL/100L of water for best results.

## DIRECTIONS FOR USE: continued

## Table A: High Volume Spraying

See GENERAL INSTRUCTIONS - APPLICATION section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHT-OF-WAYS				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE /100L WATER	CRITICAL COMMENTS
Limebush	Any time of the year with good leaf cover and soil moisture	NT and Qld only	350 mL	Penetration of thick clumps may be difficult and respraying may be required. Addition of an adjuvant may improve results - contact Crop Care Australasia Pty Ltd for details.
Manuka	At flowering	Vic only	500 mL	For optimum results, add Pulse Penetrant at 200 mL/100 L of spray. Thoroughly wet foliage, stems and soil around the base of the plants.
Mesquite ( <i>Prosopis</i> spp.)	Seedling, full leaf and flowering before podding	NSW, NT, Qld and WA only	350 mL	DO NOT spray plants bearing pods. Add Chemwet 1000 at 100 mL/100L of water for best results.
<i>Prosopis velutina</i>		Qld only	670 mL	
Mistflower	Spring to autumn	ACT, NSW and Qld only	350 mL	Apply as a thorough foliar spray.
Mother-of-millions	Flowering	NSW and Qld only	500 mL	Add Chemwet 1000 at 100 mL/100L of water
Paddy's lucerne	Active growth	NSW only		Plants that have been continually slashed or grazed over many seasons may be difficult to control and regrowth may occur.
Parkinsonia	Under 2 m tall	NT, Qld and WA only	350 mL	Add Uptake* Oil at 500 mL/100L water. Avoid spraying under dry conditions when plants are stressed or bearing pods. Thoroughly wet foliage.
Prickly pear (common) Smooth tree pear	Active phyllode growth	All States	500 mL	Apply as a thorough foliage spray. Regrowth may occur, so a follow-up application may be necessary.
Rubber vine (Not infected with rust)	Up to 1.5 m tall at flowering	NT, Qld and WA only	350 mL	Spray all leaves and stems just to the point of run off and thoroughly spray the base of the plant. With larger, more dense stands regrowth may appear. Subsequent control of any regrowth should be done by basal bark spraying.
	Dense stands greater than 1.5 m tall at flowering		500 mL	
Siam weed	Active growth	Qld and WA only	350 mL	Add Chemwet 1000 at 100 mL/100 L of water for best results.
Sicklepod	Up to flowering	NT, Qld and WA only	200 mL	DO NOT apply to podding plants. Add Chemwet 1000 at 100 mL/100L of water for best results.
St. John's wort	From flowering to early seed set	All States	500 mL	Late spring to early summer.
Sweet briar	Up to 1.5 m tall	All States	350 mL	Add Brushkiller™ 600/Associate® at 10g/100L water to obtain more reliable results with the lower rate of Gallop. Full leaf to ripe fruit prior to leaf fall. Thorough wetting including the crown is recommended.
			500 mL	
Tobacco weed	Actively growing plants	NT, Qld and WA only	300 mL	Add Chemwet 1000 at 100 mL/100L of water for best results.

**DIRECTIONS FOR USE:** continued**Table A: High Volume Spraying**See **GENERAL INSTRUCTIONS - APPLICATION** section for application method details.

<b>AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHT-OF-WAYS</b>				
<b>WEEDS CONTROLLED</b>	<b>WEED GROWTH STAGE</b>	<b>STATE</b>	<b>RATE /100L WATER</b>	<b>CRITICAL COMMENTS</b>
<b>Wattle (<i>Acacia</i> spp.) (except corkwood wattle)</b>	1 to 3 m tall	All States	350 mL	Add Chemwet 1000 at 100mL/100L of water for best results.
<b>Wild rosemary (<i>Cassinia laevis</i>)</b>	Active growth, 0.5 to 1.0 m tall	Qld only	350 to 500 mL	Use lower rate on seedlings 0.5m tall. Apply as a thorough foliar spray.
<b>Wild tobacco tree</b>	Spring to autumn up to 2 m tall	ACT, NSW and Qld only	350 mL	

**Table B: Aerial Application**See **GENERAL INSTRUCTIONS - APPLICATION** section for application method details.**AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHT-OF-WAYS**

<b>WEEDS CONTROLLED</b>	<b>WEED GROWTH STAGE</b>	<b>STATE</b>	<b>RATE /100L WATER</b>	<b>CRITICAL COMMENTS</b>
<b>Blackberry</b>	Summer to autumn	NSW, Qld, SA, Vic and WA only	10L	Where herbicides other than Group I herbicides have been used, allow two seasons regrowth to occur before respraying with Gallop. WARNING: <i>Eucalyptus</i> species up to 4 m may be killed if sprayed during this treatment. Mature trees which are 15 to 20 m tall may be partially defoliated but are likely to recover.
<b>Gorse</b>		Tas only		Helicopter application only
<b>Cockspur thorn Crofton weed Lantana Mistflower</b>	Late autumn	NSW, NT and Qld only (helicopter only)	1.5 L plus 6L Amine 625	Spray with calibrated equipment using the half overlap opposite pass technique applying a minimum spray volume of 150 L/ha. Follow-up respraying will be required.
<b>Lantana</b>			10L	Helicopter application only
<b>Rubber vine (Not infected with rust)</b>	When flowering	NT and Qld only (helicopter only)	3 L to 5 L	Use rates will depend upon the density and height of the rubber vine stand. The higher rate should be used on dense stands. However, complete coverage and penetration may be difficult. Follow-up respraying will be required. Any regrowth should be sprayed with a suitable basal bark herbicide.
<b>St. John's wort</b>	Flowering to early seed set (Nov-Jan)	NSW only	4 L	Helicopter application only. Follow up spraying will be required in the following season.

**AGRICULTURAL NON-CROP AREAS ON FLOOD PLAINS**

<b>WEEDS CONTROLLED</b>	<b>WEED GROWTH STAGE</b>	<b>STATE</b>	<b>RATE / ha</b>	<b>CRITICAL COMMENTS</b>
<b>Parkinsonia</b>	Seedlings 1 - 2 m tall, or 12 - 24 months old	Qld and NT only (helicopter only)	3 L	Add Uptake* Oil at 1 L/ha

Table C: Controlled Droplet Application (C.D.A.)

See GENERAL INSTRUCTIONS - APPLICATION section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHT-OF-WAYS				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE / ha	CRITICAL COMMENTS
Blackberry in association with: Docks Ragwort St. John's wort Thistles	Summer to autumn	All States	Apply undiluted	One application may give satisfactory control but subsequent regrowth and seedlings should be re-sprayed after hardening off. Where herbicides other than Group I herbicides have been used, allow two seasons regrowth to occur before respraying with Gallop.

Table D: Low Volume High Concentrate Application Techniques  
(GAS POWERED GUN, SPRINKLER SPRAYER)

See GENERAL INSTRUCTIONS - APPLICATION section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHT-OF-WAYS				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE / 10L WATER	CRITICAL COMMENTS
Blackberry	Late Spring to autumn	ACT, NSW, Qld, SA, TAS and WA only	335 mL	Apply to actively growing bushes which are able to be sprayed on all sides. For larger bushes, the high volume application technique is recommended.
Camphor laurel Cockspur thorn Crofton weed	Less than 1.5 m high	ACT, NSW and Qld only	500 mL	
<i>Eucalyptus</i> species	Seedlings up to 2 m tall	All States	335 mL	<b>Gas powered Gun only:</b> Apply to actively growing bushes not more than 1.5 m tall that have not more than 5 stems from the crown.
Mistflower	Less than 1.5 m high	ACT, NSW and Qld only	500 mL	
Sweet briar	1.5 m tall, full leaf to ripe fruit	NSW only		
St. John's wort	During flowering to early seed set	NSW, Tas and Vic only		<b>Gas powered Gun only:</b> One application should provide control. Minor regrowth and seedlings may be re-treated the following summer.
Wild tobacco tree	Less than 1.5 m high	ACT, NSW and Qld only		Apply to actively growing bushes which are able to be sprayed on all sides. For larger bushes, the high volume application technique is recommended.

Table E: Boom Application

See GENERAL INSTRUCTIONS - APPLICATION section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHT-OF-WAYS				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE /ha	CRITICAL COMMENTS
Galenia	Fresh growth during spring to summer	NSW only	5L	Rough mine sites will require adequate spray equipment such as boomless nozzles for effective coverage.
Sicklepod	Up to flowering	NT and Qld only	3L	DO NOT apply to podding plants. Add Chemwet 1000 at 100 mL/100L of water.
St. John's wort	Flowering to early seed set (Nov-Jan)	NSW only	2 to 4 L	Use the higher rate on dense infestations and when longer residual control is required. Follow-up respraying will be required in the following season.

## 2. FALLOW SITUATIONS

Table F: Boom Application

See GENERAL INSTRUCTIONS - APPLICATION section for application method details.

FALLOW				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE /ha	CRITICAL COMMENTS
Blackberry nightshade - Suppression only	10 to 25 cm tall prior to flowering	NSW and Qld only	200 to 400 mL + 1.2L Gladiator or Roundup <sup>^</sup> CT + LI 700 or 1L Credit <sup>®</sup> plus 1L Bonus <sup>®</sup> or 1L Roundup <sup>^</sup> PowerMAX <sup>^</sup>	FOR USE BY GROUND EQUIPMENT ONLY. Plants must be actively growing
Camel melon Prickly paddy melon Cucumber melon ( <i>Cucumis melo</i> )	From 2 leaf to 50 cm diameter			Use the lower rate on the smaller weeds, as specified in the weed growth stage (or up to 5 cm diameter for <i>Polymeria pusilla</i> ).
Common sowthistle	From 8 leaf to flowering			Refer to mixture product label for use of LI 700 Surfactant.
Cow vine	From 2 to 5 leaf up to 15 cm diameter, prior to flowering			DO NOT add an adjuvant to mixtures containing Credit and Bonus.
Lucerne (established)	Active growth, 15 to 25 cm high, during spring			DO NOT plant susceptible crops for up to nine months after application, as specified in General Instructions - Minimum Re-cropping Periods - Black Cracking Clay Soils, Northern NSW & Qld.
<i>Polymeria pusilla</i>	2 to 12 leaf up to 20cm diameter, prior to flowering			Dry conditions after application will increase the re-cropping interval.
			300 to 500 mL + 1.2L Gladiator or Roundup <sup>^</sup> CT + LI 700 or 1L Credit <sup>®</sup> plus 1L Bonus <sup>®</sup> or 1L Roundup <sup>^</sup> PowerMAX <sup>^</sup>	
			200 to 400 mL + 1.2L Gladiator or Roundup <sup>^</sup> CT or 1L Credit <sup>®</sup> plus 1L Bonus <sup>®</sup> or 1L Roundup <sup>^</sup> PowerMAX <sup>^</sup>	

Table B: Blanket Wiper Application

See GENERAL INSTRUCTIONS - APPLICATION section for application method details.

FALLOW				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE	CRITICAL COMMENTS
<b>Bitterbark</b> ( <i>Alstonia constricta</i> )	From summer to end of autumn	Qld	1:4 (1 part Gallop to 4 parts water)  2% solution for spot spray (eg. 100 mL Gallop in 5 L water)	For use with blanket wipers only.  For best results apply in autumn to tall (>60 cm) plants using two opposite directional passes (up and back)  Follow up "missed" plants with a spot spray application. These will be obvious after 6 weeks.  Blanket wiper application can be made in summer when plants are smaller but follow up spot spraying may be necessary.  Do not disturb (cultivate) the treated patches for at least 3 months. Best long term control is achieved when patches are left undisturbed for as long as possible after treatment (at least 6 months).  Spot spraying "missed" plants: thoroughly wet all stems and leaves without producing any solution run off. Avoid any spray reaching the soil surface.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**IN TASMANIA FOR BLACKBERRY:** DO NOT treat bushes carrying mature or near mature fruit.

**FOR NATIVE VEGETATION:** Use of Gallop on native vegetation must be done in accordance with STATE and/or LOCAL legislation.

**WITHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED.**

**GENERAL INSTRUCTIONS****FALLOW SITUATIONS****MINIMUM RECROPPING PERIODS- Black  
Cracking Clay Soils, Northern NSW & Qld.****Table F: Boom Application**

Plant-back periods for crops following the application of Gallop for rates up to 600 mL/ha				
RATE mL/ha	200	300	400	600
CROP	MONTHS			
Wheat	2	2	4	4
Barley	2	2	4	4
Canola	2	4	4	4
Faba Beans	4	4	6	6
Chickpeas	4	6	6	6
Lucerne	6	9	9	9

These plantback periods are based on a normal rainfall pattern. During drought conditions (or when the rainfall is less than 100 mm for a period of 4 months or greater) the plantback period may be significantly longer.

**Table G: Blanket Wiper Application**

Plant-back periods for crops following blanket wiper application.	
CROP	MONTHS
Broadleaf crops	18
Lucerne	6 This will allow any potential soil residues to dissipate, if any, and allow effective control.

**NOTE:** Before using Gallop in tank mixes with other herbicides, check the plant-back information on all product labels. The most residual product with the longest plant-back period, will determine the time between spraying and planting.

**COMPATIBILITY**

Follow any regional restrictions, and all directions and restrictions on the label, for any chemical mixed with Gallop (e.g 2,4,-D amine).

Gallop is compatible with the following herbicides: Amine 625, Lantana DP600, Brushkiller™/Associate®, Gladiator, Roundup<sup>CT</sup>, Credit<sup>®</sup> plus Bonus<sup>®</sup> and Roundup<sup>^</sup> PowerMAX<sup>^</sup>.

Gallop is compatible with the following adjuvants, as per Directions for Use:  
Chemwet 1000, Pulse<sup>^^</sup> Penetrant, LI 700\* and crop oils.

**MIXING**

Mix only with water.

Half fill the spray unit with water, and add the required amount of Gallop. Add the remaining water with the agitator running. If required, then add spray oils or wetters (surfactants). Maintain mechanical or by-pass agitation in the spray tank during spraying. Only mix sufficient solution for immediate daily use and avoid storing.

**APPLICATION****1. WOODY WEED SITUATIONS**

Weeds need to be actively growing for herbicides to have optimum effect. Delay treatment until all regrowth has had time to grow approximately 1 metre in situations which have been bulldozed, slashed, burnt, ploughed or areas having a previous chemical treatment.

**Table A. High Volume Spraying**

Thorough coverage of foliage to the point of run-off is essential, however, avoid excess spraying which is wasteful of chemical.

**Hand Gun**

Apply the recommended mix to give full coverage of leaves and stems through a No. 6 to 8 tip at 700 to 1500 kPa (400 to 500 kPa for St John's wort). A spray volume of 3000 to 4000 L per infested hectare of 1 to 2 metre high blackberry (30 to 40 L/100m<sup>2</sup>) should be used. Use 2000 L of spray mixture/ha of galenia infestation (i.e. 20L/100 m<sup>2</sup> infested area).

**Knapsack**

Apply the recommended spray mix to give full coverage of leaves and stems. The final volume of application should be similar to hand gun. A spray volume of 3 to 4 L/10m<sup>2</sup> infested area should be used. A spray volume of 2L/10m<sup>2</sup> should be used for an area infested with galenia.

**Table B. Aerial Application**

Apply in 200 L of water/ha using an aircraft to apply 100L per pass on a double overlap pattern using nozzle configurations to produce droplets of 250 to 350 micron diameter. The potential for damage from drift can be greatly reduced by avoiding unsuitable spraying conditions and using spray pressure and nozzles to minimise the production of small droplets. DO NOT spray when wind exceeds 15 km/hr and /or air temperature reaches 35°C.

**Table C. Controlled Droplet Application (C.D.A)**

Results similar to high volume spraying can be obtained using Micron Herbi\* or similar equipment. Select a nozzle to give a flow rate of 2 mL/sec and sweeping action of approximately 1m/sec to ensure a droplet density of 20/cm<sup>2</sup>. Use a marking agent, as recommended by the equipment manufacturers, to check spray coverage. Also, consult directions provided with C.D.A unit.

**Table D. Low Volume High Concentrate Application Techniques**

Good control will be achieved, similar to high volume application, where bush size enables good coverage of entire bush. Use a marking agent, as recommended by the equipment manufacturers, to check spray coverage.

**Gas powered gun**

Apply 50mL shots to obtain uniform coverage of 4 to 5 m<sup>2</sup> of surface area of bush. This relates to 20 droplets/cm<sup>2</sup> of leaf surface.

**Sprinkler sprayer**

This technique involves using a micro sprinkler that is connected to a hollow fibre glass rod attached to a pneumatic knapsack sprayer. Use at low pressures (50 to 200 kPa) and apply with a slow sweeping action over the top of the plants ensuring even coverage on the leaves.

**Table E. Boom Application**

Application in a minimum spray volume of 200 L/ha for galenia and St. John's wort and 600 L of water/ha for sicklepod. Flat fan nozzles are recommended, using pressure in the range of 200 - 300 kPa. Boom height must be set to ensure double overlap of nozzle patterns.

**2. FALLOW SITUATIONS****Table F. Boom Application**

Application of Gallop in a minimum spray volume of 50 L/ha is recommended. Flat fan nozzles are recommended, using pressure in the range of 200 - 300 kPa. Boom height must be set to ensure double overlap of nozzle patterns.

**Table G. Blanket Wiper Application**

Blanket needs to be made from durable and wettable material with a rigid backing.

Blanket should be rigidly mounted behind motorised vehicle (eg. tractor, 4-wheel drive vehicle) and set low but never touching the ground. The chemical solution should be fed to the blanket at a flow rate sufficient to keep the blanket wet but not dripping. In thick patches the blanket may require more frequent solution recharge (re-wetting).

Ideally, a scraper bar should be mounted at the front of the blanket in order to scrape or damage the bark (but not sever the stems) prior to the blanket wiping stems. This scraper may be mounted at the front of the vehicle.

Two passes (in opposite direction) with the blanket increases the contact with the plant. Ground speeds of 10 - 15 kph are ideal for blanket wiping application.

**CLEANING SPRAY EQUIPMENT****Rinsing**

After using Gallop empty the spray unit completely and drain the whole system. Thoroughly wash inside the unit using a pressure hose. Drain the spray unit, and clean any filters in the tank, pump, lines, hoses and nozzles.

After cleaning the spray unit as above, quarter fill with clean water and circulate through the pump, lines,

hoses and nozzles. Drain and repeat the rinsings procedure twice.

**Decontamination**

Before spraying cotton and other sensitive crops with equipment that has been used to apply Gallop, see PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS section.

Wash the tank and rinse the system as above. Then quarter fill the tank and add an alkali detergent such as Spraymate Tank and Equipment Cleaner at 500 mL/100 L of water and circulate throughout the system for at least 15 minutes.

Drain the whole system. Remove filters and nozzles and clean them separately. Finally flush the system with clean water and allow to drain.

**Rinse water should be discharged onto a designated disposal area or, if this is unavailable, onto unused land away from desirable plants and watercourses.**

**RESISTANCE WEEDS WARNING****GROUP I HERBICIDE**

'Gallop' Herbicide is a member of the pyridines group of herbicides. Gallop has the disruptors of plant cell growth mode of action. For weed resistance management, Gallop is a Group I herbicide.

Some naturally-occurring weed biotypes resistant to Gallop and other Group I herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Gallop or other Group I herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Crop Care Australasia Pty Ltd accepts no liability for any losses that may result from the failure of Gallop to control resistant weeds.

Strategies to minimise the risk of herbicide resistance are available. Contact your farm chemical supplier, consultant, local Department of Agriculture, or local Crop Care Australasia Pty Ltd representative.

**PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS**

Crops susceptible to Gallop include, but are not limited to: peas, lupins, lucerne, navy beans, peanuts, soybeans and other legumes, cotton, flowers, fruit, hops, ornamentals, shade trees and *Pinus* spp., potatoes, safflower, sugar beet, sunflowers, tobacco, tomatoes, vegetables and vines. Gallop is damaging to susceptible crops during both growing and dormant periods. Grasses are normally unaffected and establish quickly after treatment.

Picloram, one of the active constituents in this product can remain active in the soil for extended periods depending on soil type and application rate, rainfall, temperature, humidity, soil moisture and soil organic matter.

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands, pastures, waterways or native vegetation.

DO NOT allow physical spray drift onto waterways, native vegetation and susceptible crops.  
DO NOT apply close to, or in areas, containing roots of desirable vegetation, where treated soil may be washed onto areas growing (or areas to be planted with) desirable plants.  
DO NOT apply on sites where surface water from heavy rain can be expected to run off to areas containing, or to be planted with susceptible crops or plants. DO NOT move soil, which may have been treated to areas where desirable plants are to be grown.

#### PROTECTION OF LIVESTOCK

Poisonous plants may become more palatable after spraying and stock should be kept away from these plants until they have died down. Many plants remain poisonous after death, and stock should not be allowed access, as there is a likelihood that they may graze the dead material. Such material should be burnt if possible.

#### PROTECTION OF WILDFISH, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

#### STORAGE AND DISPOSAL

Store in closed, original container in a dry, cool well-ventilated area. DO NOT store for prolonged periods in direct sunlight.  
Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should NOT be burnt.

#### SMALL SPILL MANAGEMENT

Wear protective equipment (See Safety Directions). Apply absorbent material such as earth, sand, clay granules or cat litter to the spill. Sweep up material for disposal when absorption is completed and contain in a refuse vessel for disposal. (See Storage and Disposal section). If necessary, wash the spill area with an alkali detergent and water and absorb as above the wash liquid for disposal.

#### SAFETY DIRECTIONS

Harmful if swallowed. May irritate the eyes and skin. Avoid contact with eyes and skin. When preparing spray, wear cotton overalls buttoned to the neck and wrists, a washable hat, elbow-length PVC gloves and a face shield or goggles. If the product is in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

#### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

#### MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet which can be obtained from your supplier.

#### CONDITIONS OF SALE

Crop Care Australasia Pty Ltd will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply or use of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Crop Care Australasia Pty Ltd is limited to the replacement of the goods or (at the option of Crop Care Australasia Pty Ltd) the refund of the price paid and is conditional upon a claim being made in writing and where possible sufficient part of the goods to enable proper examination being returned to Crop Care Australasia Pty Ltd within thirty days of delivery.

In a Transport Emergency Dial <b>000</b> Police or Fire Brigade	SPECIALIST ADVICE IN AN EMERGENCY ONLY <b>1800 033 498</b> ALL HOURS- AUSTRALIA WIDE
--	---



Crop Care Australasia Pty Ltd  
ACN 061 362 347  
Unit 15/16 Metroplex Avenue  
Murarrie Qld 4172  
Ph : (07) 3909 2000

APVMA Approval No: 61664/0407

#### In case of emergency: Phone 1800 03 3498

Ask for shift supervisor. Toll free 24 hours.

- ® Associate, Credit and Bonus are registered trademarks of Nufarm Australia Limited.
- \* Other registered Trademarks
- ^^ Pulse is a registered trademark of Nufarm Technologies USA Pty Ltd
- ™ Trademark pending
- ^ Roundup and Roundup PowerMAX are registered trademarks of Monsanto Technology LLC used under licence by Nufarm Australia Limited.