

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Farmoz

Prometrex® 500 SC

HERBICIDE

ACTIVE CONSTITUENT: 500 g/L PROMETRYN

GROUP C HERBICIDE

Controls early competing weeds in cotton, chickpeas, carrots, potatoes and other crops as specified in the Directions for Use table.



www.farmoz.com.au

DIRECTIONS FOR USE:

RESTRAINTS: DO NOT spray when dry conditions prevail.

DO NOT apply by aircraft.

CROP & SITUATION	WEEDS CONTROLLED	STATE	RATE/HA	CRITICAL COMMENTS
Carrots pre-emergent	Amaranth (Powell's, Slim and Redroot), Amsinckia, Blackberry Nightshade, Capeweed, Charlock, Chickweed, Common Spurry, Corn Gromwell, Fumitory (Common, White and Dense Flowered), Fat Hen, Hedge Mustard, Hexham Scent, Indian Hedge Mustard, Lesser Nettle, Mountain Sorrel, Rough Poppy, Three-Cornered Jack, Wild Radish Pre-emergent: suppression of Annual Ryegrass, Barnyard Grass, Prairie Grass and Summer Grass	NSW, Vic, Tas only	2.2 to 3.3 L	Apply onto bare moist soil within 2 to 3 days of planting. Use the lower rate on sandy soils.
	Common Sowthistle, Common Thornapple, Deadnettle, Fat Hen, Green Amaranth, Less Swinecress, Pigweed, Potato Weed, Stinking Roger Pre-emergent: suppression of Barnyard Grass and Crowsfoot Grass	Qld only SA, WA only	2.2 to 3.3 L 1.1 L	
Carrots at least 1 true leaf	Less than 4 true leaves: Amaranth (Powell's, Slim and Redroot), Amsinckia, Blackberry Nightshade, Capeweed, Chickweed, Common Spurry, Corn Gromwell, Fumitory (Common, White and Dense Flowered), Fat Hen, Lesser Nettle, Mountain Sorrel, Rough Poppy, Three-cornered Jack	NSW, Vic, Tas, SA, WA only	1.7 to 2.2 L	Spray when weeds are young and actively growing. Use the higher rate on heavy soils and for Three-cornered Jack.
Celery up to 3 weeks after transplanting	Amaranth, Bladder Ketmia, Chickweed, Dwarf Nettle, Fat Hen, Nettles, Pigweed, Potato Weed, Radish, Shepherd's Purse, Sowthistle, Wild Turnip	Qld only	1 to 3.3 L	Apply to moist soil. Use the lower rate on sandy soils. Some grasses are also controlled.
Celery up to 4 weeks after transplanting	Amaranth (Powell's, Slim and Redroot), Amsinckia, Blackberry Nightshade, Capeweed, Charlock, Chickweed, Common Spurry, Corn Gromwell, Fumitory (Common, White and Dense Flowered), Fat Hen, Hedge Mustard, Hexham Scent, Indian Hedge Mustard, Lesser Nettle, Mountain Sorrel, Rough Poppy, Three-cornered Jack, Wild Radish Pre-emergent: suppression of Annual Ryegrass, Barnyard Grass, Prairie Grass and Summer Grass	NSW, Vic, SA, WA only	2.2 to 3.3 L	Apply onto moist soil. Use the lower rate in SA.

CROP & SITUATION	WEEDS CONTROLLED	STATE	RATE/HA	CRITICAL COMMENTS
Chickpeas	Deadnettle, Indian Hedge Mustard, Lesser Swinecress, Milk Thistle, Prickly Lettuce, Purple Goosefoot, Shepherd's Purse, Turnip Weed, Wireweed, and suppression of Black Bindweed and Paradoxa Grass	Qld, NSW, SA only	1.5 L + 1.5 L Flowable Simazine (500 g/L)	Apply immediately post-planting. Application should not be made to ridged or excessively cloddy soil. For reliable results significant rain, 20-30 mm, is necessary within 2-3 weeks of sowing.
Cotton (pre-emergent)	Barnyard Grass, Bathurst Burr, Bellvine, Blackberry Nightshade, Black Pigweed, Bladder Ketmia, Boggabri Weed, Caltrop (Yellow Vine), Caustic Weed, Common Sowthistle, Common Thornapple, Cowvine (Peachvine), Fat-hen, Green Amaranth, Mintweed, Mossman River Grass, Noogoora Burr, Pigweed, Prickly Paddy Melon, Redroot Amaranth, Slim Amaranth, Summer Grass, Wild Gooseberry	Qld, NSW only	3.3 to 4.5 L	Apply onto bare moist soil or irrigate within 3 days of spraying. Apply the lower rate if overhead irrigating. Some mechanical cultivating may be required.
Cotton 15 cm high			1.1 to 2.2 L	Early spraying: Apply as a directed spray when the weeds are not more than 5 cm high. Add surfactant such as WETSPRAY 1000® Surfactant. Use lower rate on light soils.
Cotton after last cultivation			2.2 to 4.5 L	Late spraying: Apply as a directed spray. The weed should not be more than 8 cm high and actively growing. Add a surfactant such as WETSPRAY 1000 Surfactant. Use lower rate on light soils.
Cotton Maintenance of fallow areas and after furrowing-up prior to planting	The above plus winter annuals such as: Amsinckia, Capeweed, Chickweed, Common Sowthistle, Common Spurry, Corn Gromwell, Deadnettle, Fumitory (Common, White and Dense Flowered), Lesser Nettle, Mountain Sorrel, Rough Poppy, Three-cornered Jack Less than 4 true leaves: Hedge Mustard, Hexham Scent, Indian Hedge Mustard, Powell's Charlock, Wild Radish	Qld, NSW only	2.2 to 4.5 L	Apply to bare moist soil or when rain is imminent as a pre-emergent treatment to weeds or as an early post-emergent treatment to weeds after cultivation or listing (furrowing-up). Use the lower rate when short term weed control is required. If conditions remain dry after application, as a pre-emergent treatment, lightly incorporate to a depth of approximately 5 cm.
Pastures from 5 leaf stage onwards (not Legume seed crops)	Less than 4 true leaves: Amaranth (Powell's, Slim and Redroot), Amsinckia, Blackberry Nightshade, Capeweed, Chickweed, Common Spurry, Corn Gromwell, Deadnettle, Fumitory (Common, White and Dense Flowered), Fat-hen, Lesser Nettle, Mountain Sorrel, Rough Poppy, Three-cornered Jack Less than 4 true leaves: Charlock, Hexham Scent, Hedge Mustard, Indian Hedge Mustard, Wild Radish	Sthn NSW, Vic, SA, WA only	550 to 850 mL 550 to 850 mL plus 480 mL of a 242 g/L MCPA	Use the higher rate where weeds are in the 4 true leaf stage. DO NOT add 2,4-D or MCPA if pastures are undersown with Lucerne, Clover or Medics, unless some damage is acceptable.
Perennial Grass Seed Crops Sirocco, Phalaris, Demeter Fescue, Currie Cocksfoot, Medea Ryegrass 3 to 6 leaf stage	Less than 4 true leaves: Amaranth (Powell's, Slim and Redroot), Amsinckia, Blackberry Nightshade, Capeweed, Chickweed, Common Spurry, Corn Gromwell, Fumitory (Common, White and Dense Flowered), Fat Hen, Lesser Nettle, Mountain Sorrel, Rough Poppy, Three-cornered Jack	NSW, Vic, SA, WA only	1.1 to 2.2 L	Spray when weeds are young and actively growing. Use the higher rate on heavy soils and for Three-cornered Jack. DO NOT use more than 1.1 litres per ha on Medea Ryegrass.
Perennial Ryegrass Seed Crops 3 to 6 leaf stage		Tas only		
Potatoes prior to emergence of the crop	Less than 4 true leaves and pre-emergent: Amsinckia, Blackberry Nightshade, Capeweed, Charlock, Chickweed, Corn Gromwell, Common Spurry, Deadnettle, Fat-hen, Fumitory (Common White and Dense Flowered), Hedge Mustard, Hexham Scent, Indian Hedge Mustard, Lesser Nettle, Mountain Sorrel, Powell's Slim and Redroot Amaranth, Rough Poppy, Three-cornered Jack, Wild Radish Pre-emergent: Annual Ryegrass, Prairie Grass, Barnyard Grass and Summer Grass	Qld, NSW, Vic, Tas, WA, only	2.3 to 3.3 L	Apply in the late Autumn and Winter. Spray when weeds are actively growing. Use the lower rate on light soils and the higher rate on heavy soils.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THE LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

CHICKPEAS: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 9 WEEKS AFTER APPLICATION.

OTHER CROPS: NOT REQUIRED WHEN USED AS DIRECTED.

GENERAL INSTRUCTIONS

FARMOZ PROMETREX 500 SC is a selective systemic herbicide, absorbed by the leaves and roots of weeds. This product controls many early competing weeds in cotton, when applied to the actively growing weeds. When applied to bare soil in celery, chickpeas, potatoes, carrots and cotton, many common broadleaf weeds are controlled and the density of grasses is reduced.

RESISTANCE WEEDS WARNING

FARMOZ PROMETREX 500 SC Herbicide is a member of the Triazine group of herbicides. This product has the photosynthesis at photosystem II inhibitor mode of action. For weed resistance management PROMETREX 500 SC is a Group C herbicide.

Some naturally-occurring weed biotypes resistant to PROMETREX 500 SC and other Group C herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by PROMETREX 500 SC or other Group C herbicides.

However resistance should not be assumed without first viewing the method of application, the coverage and the timing of application.

Resistance to PROMETREX 500 SC cannot be predicted and may occur at any time. It is not due to a fault in the product and FARMOZ Pty. Ltd. accepts no liability for any losses that may result from the failure of PROMETREX 500 SC to control resistant weeds.

MIXING INSTRUCTIONS

Settling may occur after storage for some weeks. Stir product or invert container several times before opening. Pour the flowable into the spray vat through a strainer to remove any dry particles or flakes, which can occasionally occur under hot storage conditions. Add the full quantity of chemical to the partly filled spray tank while agitating. Fill tank and agitate to ensure thorough mixing. Continue agitation while spraying. Agitate vigorously from the bottom if allowed to stand. Reseal part-used containers immediately.

PRECAUTION

DO NOT apply more than 4.5 L per ha. of product post-emergent per season in cotton. Severe injury to cotton and chickpeas may result if heavy rain occurs between pre-emergence applications and crop emergence, or shortly after post-emergence spraying and this risk must be assumed by the user.

PROTECTIONS OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands, or pastures.

DO NOT use on Carrots on light sandy soils.

DO NOT plant crops other than Carrots in PROMETREX 500 SC treated Cotton land for at least six months after the Cotton crop has been removed.

DO NOT plant crops other than transplanted Celery, Chickpeas or Carrots in areas within 6 months following an application of the product at rates above 2.2 litres per hectare.

DO NOT add 2,4-D or MCPA if pastures are undersown with Lucerne, Clover or Medics, unless some damage is acceptable.

DO NOT add 2,4-D or MCPA if spraying Cotton, vegetables or seedling pastures.

DO NOT apply more than 4.5 litres per hectare of product post-emergent per season in Cotton. Heavy rain following application prior to emergence may cause damage to Chickpeas.

PROTECTIONS OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to fish. DO NOT contaminate streams, rivers or waterways with the chemical or used container. Avoid use in channels and drains where roots of desirable plants may extend and in other situations where desirable trees grow close to treated areas.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

May irritate the eyes and skin. Avoid contact with eyes and skin. DO NOT inhale spray mist. After use and before eating, drinking or smoking wash hands, arms and face thoroughly with soap and water.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

MSDS

Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for PROMETREX 500 SC is available from FARMOZ on request. Call Customer Service on (02) 9431 7800.

CONDITIONS OF SALE: The use of FARMOZ PROMETREX 500 SC Herbicide being beyond the control of the manufacturer, no warranty expressed or implied is given by FARMOZ Pty. Ltd., regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and FARMOZ Pty. Ltd. accepts no responsibility for any consequence whatsoever from the use of this product.

® Registered trademarks of a Makhteshim-Agan Company

NOT A DANGEROUS GOOD ACCORDING TO THE AUSTRALIAN DANGEROUS GOODS (ADG) CODE.

PACK SIZE: 20 L