

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



Aurora Turf®

Systemic Fungicide

ACTIVE CONSTITUENT: 250 g/L PROPICONAZOLE

For Control of Dollar Spot in Bent and Queensland Blue Couch,
and Spring Dead Spot in Couch
as specified in the Directions for Use table.

DIRECTIONS FOR USE

RESTRAINTS:

DO NOT apply to turf under heat or moisture stress.

DO NOT apply more than one application per year on the following couch varieties: G29, Greenless Park, National Park, Tifway, Sportsway and Wagga City.

SITUATION	DISEASE	STATE	RATE	CRITICAL COMMENTS
Bent Grass in bowling greens, parks and sporting areas.	Dollar Spot (<i>Sclerotinia homeocarpa</i>)	Vic only	30 to 60 mL/ 100 m ² in 10 to 20 L water.	Spray when conditions are warm and humid, from September to March. Make a second application 14-28 days later if conditions continue to favour disease development. Use rates towards the lower end of the range as a preventive program and against light to moderate infection. Use rates towards the higher end of the range and shorter intervals as a preventive or curative treatment when conditions are highly favourable for the disease.
Bent, Queensland Blue Couch grasses in bowling greens, golf greens, parks and sporting areas.		NSW, Qld, SA, WA only		
Couch Turf in bowling greens, parks and sporting areas.	Spring Dead Spot (<i>Leptosphaeria</i> spp.)	NSW, Qld, Vic, SA, WA only	60 mL/100 m ² in 150 L of water.	Apply as a soil drench and water in immediately, ensuring thorough mixing with the soil. Spray in January to March, after renovation and recovery of active growth. Make a second application one month later where infection is severe. DO NOT renovate treated greens until active growth has recommenced in Spring. DO NOT spray in the Spring/Summer period prior to renovation. AURORA TURF may cause bleaching of the grass after application in late Summer/Autumn and also produce a greening effect in the following Spring. Couch may be slow to recommence active growth in the Spring, particularly in cooler regions. These effects should be allowed to grow out before treating again. DO NOT apply more than twice per year.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.



GENERAL INSTRUCTIONS

MIXING

Add the required amount of product directly to the spray tank and mix well.

COMPATIBILITY

AURORA TURF is compatible with chlorothalonil.

FUNGICIDE RESISTANCE WARNING

AURORA TURF is a member of the DMI **GROUP C FUNGICIDE** group of fungicides. For fungicide resistance management AURORA TURF is a Group C fungicide. Some naturally occurring individual fungi resistant to AURORA TURF or any other Group C fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. The resistant fungi will not be controlled by AURORA TURF or other Group C fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, FARMOZ Pty. Ltd. accepts no liability for any losses that may result from the failure of AURORA TURF to control resistant fungi.

PROTECTION OF LIVESTOCK

DO NOT graze treated turf or feed grass clipping from treated area to poultry or livestock.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to fish. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area.

DO NOT store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Avoid contact with eyes and skin. DO NOT inhale spray mist.

FIRST AID

If poisoning occurs, contact a Doctor or Poisons Information Centre on 131126.

MSDS

Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for AURORA TURF is available from FARMOZ on request. Call Customer Service on (02) 9363 3611.

CONDITIONS OF SALE: The use of FARMOZ AURORA TURF Systemic Fungicide being beyond the control of the manufacturer, no warranty expressed or implied is given by FARMOZ Pty. Ltd., regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and FARMOZ Pty. Ltd. accepts no responsibility for any consequence whatsoever resulting from the use of this product.

® Registered trademark of FARMOZ Pty. Ltd.

PACK SIZE: 1 L, 5 L

