

**POISON**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

***Omite***<sup>®</sup>

**300W WETTABLE POWDER MITICIDE**

**ACTIVE CONSTITUENT: 300 g/kg PROPARGITE**

**GROUP 12C INSECTICIDE**

**For the control of mites in crops as per the DIRECTIONS OF USE table.**



**Net Contents: 10 kg**

Chemtura Australia Pty Ltd  
Level 7, 435 King William Street Adelaide SA 5000  
Tel: (08) 8112 0900 Fax: (08) 8112 0999

 **Chemtura**  
AGROSOLUTIONS™

**DIRECTIONS FOR USE:****RESTRAINTS:**

DO NOT add wetting agents or crop oils to spray solution.

| Crop                              | Pest Controlled                                   | States                            | Rate     |         | Withholding Period   | Critical Comments   |
|-----------------------------------|---|-----------------------------------|----------|---------|--|---|
|                                   |   |                                   | per 100L | per ha  |  |   |
| Stone fruit                       | Two spotted mite                                  | All States                        | 200 g    | 5.5 kg  | 7 days   | Apply as a high volume spray as soon as mites appear. Further applications may be necessary later in the season. Whatever volume is applied a minimum of 5.5 kg/ha is essential.  |
|                                   | European red mite                                 | Qld & WA only                     |          |         |  |   |
| Apples                            | Two spotted mite                                  | All States                        | 200 g    | 5.5 kg  |  | Apply as necessary as a high volume spray from pink onwards. Whatever spray volume is applied a minimum of 8.5 kg/ha is essential (except in Tasmania).   |
|                                   | European red mite                                 | Vic, NSW, ACT, SA, WA & Qld only  |          | 8.5 kg  |  |   |
|                                   |   | Tas only                          |          | 5.5 kg  |  |   |
| Apples and Stonefruit             | Biological control of mites                       | All States                        | 100 g    | –       |  | Apply where biological control is being practised. Complete spray coverage is essential for adequate control.   |
| Pears                             | Two spotted mite                                  | All States                        | 200 g    | –       | Post-harvest spray only: Apply as high volume spray to achieve a minimum of 5.5kg/ha. This product may cause leaf spoiling and possible defoliation, this effect is more likely in hot conditions. |   |
| Strawberries                      | Two spotted mite ( <i>Tetranychus urticae</i> )   | Vic, Tas, NSW, ACT, WA & Qld only | 100 g    | –       | 3 days   | May be phytotoxic on some varieties when used at higher concentrations. Spray when pests first appear. A maximum of 3 sprays applied over a period of 3 to 4 weeks may be necessary depending on the level of infestation. Complete spray coverage is essential for adequate control. Do NOT use higher concentration or low volume high concentrate spray application. Do NOT mix with other pesticides or foliar nutrients. DO NOT spray when temperature exceeds 27°C. |
|                                   | Biological control of mites                       |                                   |          |         |  | Apply where biological control is being practised. Complete spray coverage is essential for adequate control. DO NOT mix with other pesticides or foliar nutrients. DO NOT spray when temperature exceeds 27°C.   |
| Bananas                           | Two spotted mite (red spider)                     | Qld, NSW, ACT & WA only           | 100 g    | 1.25 kg | 7 days   | Spray when bunches covered and when pests appear and repeat as necessary. Avoid spraying under hot-humid-slow drying conditions when young bananas may be susceptible to insecticide damage.  |
| Passionfruit                      | Passionvine mite ( <i>Brevipalpus phoenicis</i> ) | Qld & WA only                     | 100 g    | –       |  | Apply as pest populations indicate. Do NOT apply more than twice per season.  |
| Beans, Tomatoes, other vegetables | Two spotted mite                                  | All States                        | 100 g    | –       |  | Apply when pests appear and repeat as necessary. Good spray cover is necessary.   |
|                                   | Spider mites ( <i>Tetranychus</i> spp.)           | Qld & WA only                     |          |         |  |   |
| Hops                              | Two spotted mite                                  | Vic, WA & Tas only                | 200 g    | –       |  | Apply as a high volume spray on first appearance of mites. e.g. when growth is 2–3 metres high. Thoroughly spray all foliage with special attention to base of plants, repeat as necessary.   |
| Ornamental plants                 | Two spotted mite                                  | All States                        | 100 g    | –       | –  | Spray when mites appear and repeat when necessary.  |
|                                   | European red mite                                 | Qld & WA only                     |          |         |  | <b>Note:</b> (1) This product may be quite phytotoxic to some ornamentals. (2) May be used in integrated mite control with the Chilean predatory mite <i>Phytoseiulus persimilis</i> for control of two spotted mite.   |

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.****WITHHOLDING PERIOD:****STRAWBERRIES: DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION.****APPLES, BANANAS, HOPS, PASSIONFRUIT, PEARS, STONEFRUIT AND VEGETABLES: DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION.****RE-ENTRY:**

DO NOT allow entry into treated areas for low exposure activities such as irrigation, scouting or weeding for 21 days unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

DO NOT allow entry into treated areas for high exposure activities such as harvesting, pruning, training or tying for 32 days unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

**GENERAL INSTRUCTIONS:**

DO NOT apply using spraying equipment carried on the back of the user.

**Mixing**

Slowly add the required amount of product to the water under agitation in the spray vat.

**Application**

This product is not an ovicide. For fruit trees, two sprays 10–14 days apart are desirable for long term results. If lower volume sprays are applied, the 2 spray programme is essential. Early applications are recommended. These should be made before the mite population is sufficient to cause serious damage. This product is non-systemic in action and high volume sprays and complete coverage are recommended.

**Compatibility**

A mixture of this product with any other product or more than one product may be ineffective or may cause serious damage. The use of such a mixture is, therefore, entirely at the user's risk. DO NOT add wetting agents or crop oils to spray solution as phytotoxicity may occur.

**INSECTICIDE RESISTANCE WARNING****GROUP 12C INSECTICIDE**

For insecticide resistance management Omite® 300W is a Group 12C insecticide.

Some naturally occurring insect biotypes resistant to Omite® 300W and other Group 12C insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Omite® 300W or other Group 12C insecticides are used repeatedly. The effectiveness of Omite® 300W on resistant individuals can be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Chemtura Australia Pty Ltd accepts no liability for any losses that may result from the failure of Omite® 300W to control resistant insects.

Omite® 300W may be subject to specific resistance management strategies. For further information contact your local supplier, Chemtura Australia Pty Ltd representative or local agricultural department agronomist.

**PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

Dangerous to fish. DO NOT contaminate streams, rivers or waterways with the product or used containers.

**STORAGE AND DISPOSAL**

Store in the closed, original container in a dry, well-ventilated area, as cool as possible out of direct sunlight.

Triple, or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and bury containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

**SAFETY DIRECTIONS**

Product is harmful if absorbed by skin contact, inhaled or swallowed. Will irritate the eyes, nose and throat and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. DO NOT inhale spray mist.

When preparing spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves, face shield and goggles. When using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water.

|   |   |
|---|---|
| <b>UN NO. 3077</b>  | <b>ENVIRONMENTALLY HAZARDOUS<br/>SUBSTANCES, SOLID, N.O.S.<br/>(CONTAINS PROPARGITE)<br/>MARINE POLLUTANT</b> |
| <b>In a Transport<br/>Emergency<br/>Dial<br/>000<br/>Police or Fire Brigade</b> | <b>SPECIALIST ADVICE<br/>IN EMERGENCY ONLY<br/>1800 033 111<br/>ALL HOURS – AUSTRALIA WIDE</b>                |
| <b>P.G. III</b>   | <b>HAZCHEM 2 Z</b>  |

After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield, goggles and contaminated clothing.


**FIRST AID**If poisoning occurs, contact a doctor or Poisons Information Centre. Phone *Australia* 131126. If swallowed, do not induce vomiting. Give a glass of water.**MATERIAL SAFETY DATA SHEET:**

Additional information is listed in the Material Safety Data Sheet.

**CONDITIONS OF SALE:**

Chemtura Australia Pty Ltd will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Chemtura Australia Pty Ltd is limited to the replacement of the goods or (at the option of Chemtura Australia Pty Ltd) the refund of the price paid and is conditional upon a claim being made in writing and where possible sufficient part of the goods to enable proper examination being returned to Chemtura Australia Pty Ltd within thirty days of delivery.

® Omite is a registered trademark of a subsidiary of Chemtura Corporation.

APVMA Approval No.  
48086/10/0710  
Printer Code  
BN  
DOM