

Front of Box

CAUTION

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

BURSTTM Herbicide

ACTIVE CONSTITUENT: 500 g/kg PROPYZAMIDE

GROUP K HERBICIDE

For selective control of certain grasses and broad leaf weeds in Lettuce, Sports turf, Home lawns and Legume seed crops and pastures, as per the DIRECTIONS FOR USE TABLE

CONTAINS 10 (20 OR 40) MEASURE PACKS WHICH IT IS ILLEGAL TO SELL SEPARATELY

CONTAINS WATER SOLUBLE PACKAGING

IMPORTANT: READ THE INSTRUCTIONS THOROUGHLY BEFORE USING THIS PRODUCT.

CONTENTS: 5,10,20 kg

Sipcam Pacific Australia Pty Limited
Suite 11, 23-31 Gheringhap Street
GEELONG VIC 3220
ACN 072 176 888
TEL:03 5223 3746

TM Trademark of Sipcam Pacific Australia Pty Ltd

NRA Approval No.: 55783/0602

Batch Number:

Date of Manufacture

Side of the box

General Instructions

Best results are obtained by applying the product to cool, wet soil

Weed Resistance Warning –

GROUP K HERBICIDE

Burst Herbicide is a member of the amide group of herbicides with diverse sites of action. For weed resistance management this is a Group K herbicide.

Some naturally occurring weed biotypes resistant to Burst Herbicide and other amide herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Burst Herbicide or other amide herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Sipcam Pacific Australia Pty Ltd accepts no liability for any losses that may result from the failure of Burst Herbicide to control the resistant weeds.

Advice as to strategies and alternative treatments that can be used should be obtained from your local supplier, consultant, local Department of Agriculture, Primary Industries Department or a Sipcam representative.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT

- **DO NOT** contaminate streams, rivers or waterways by cleaning of equipment, disposal of wastes or used containers.

PROTECTION OF CROPS

- **DO NOT** apply to, or allow spray to drift onto crops, pastures or non target – desirable plants, nor onto agricultural land scheduled for cropping or pasture, as injury to the crop or pasture may occur.

PROTECTION OF LIVESTOCK

- **DO NOT** graze sheep, cut for stock feed (including poultry and pigs) any treated areas (for 25 days after application).
- **DO NOT FEED TURF / LAWN CLIPPINGS TO POULTRY OR LIVESTOCK**

STORAGE AND DISPOSAL

- Keep out of reach of children.
- Store in the closed, original container in a dry, cool, well-ventilated area, out of direct sunlight. Keep from contact with fertilisers, insecticides, fungicides and seeds.
- **DO NOT** re-use container or dispose of undiluted chemicals on site. Shake empty into spray tank. **DO NOT** dispose of undiluted chemicals on-site. Puncture or shred and bury empty containers in a local authority landfill. If not available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

Other side of the box

SAFETY DIRECTIONS

Will irritate the eyes and skin.

Avoid contact with eyes and skin. Avoid inhaling the dust. When preparing the spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist, washable hat and elbow length PVC gloves. After use, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and contaminated clothing.

FIRST AID

- If poisoning occurs, contact a doctor or Poisons Information Centre (131126).

MATERIAL SAFETY DATA SHEET

- For further information refer to the Material Safety Data Sheet which is available from the supplier.

Spray Preparation

Burst Herbicide is a wettable powder formulation to be blended with water and applied as a spray. Partly (30%) fill the boomspray-tank with water. Put the required amount of Burst Herbicide for the area to be sprayed into water containers. Add this correct amount of pre-mixed Burst Herbicide into the boomspray-tank with the agitation system operating. Fill up to the correct volume with water. The mixture must be kept in suspension at all times by continuous agitation.

Tank-mixtures should be tested for physical compatibility and efficacy before use. **DO NOT** mix with MCPA. If tank mixing, Burst Herbicide must be in suspension before adding other herbicides, insecticides or surfactant.

- Wash sprayer thoroughly after use

DO NOT handle the water soluble packaging with wet hands or wet gloves. Keep dry.

EQUIPMENT

Always ensure the sprayer is clean, using a suitable spray and boom cleaner as directed.

It is recommended that for certain turf situations, up to 100L / ha spray volume is required.

NOTICE TO BUYER

This product is not to be used for any purpose or in any way contrary to label instructions. The buyer by use or application of this product acknowledges so doing upon the terms and conditions agreed between Sipcam Pacific Australia Pty Limited and the buyer but not limited to the following.

To the extent permitted by law all conditions and other warranties or other rights of action which buyer or any other user may have against Sipcam Pacific Australia Pty Limited (Sipcam) or Seller are hereby excluded. Sipcam hereby gives notice to buyer and other users that it will not accept responsibility for any indirect or consequential loss arising from reliance on product information or advice provided by Sipcam or on its behalf unless it can be established that such information or advice was provided negligently and that the product has been used strictly as directed. Notwithstanding the generality of this notice Sipcam will not be held liable for any loss, injury or damage indirect or consequential arising from the sale, supply, use or application of this product. Sipcam's liability shall in all circumstances be limited to replacement of the product or a refund of the purchase price paid therefor.

<p>IN A TRANSPORT EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE FOR SPECIALIST ADVICE, CONTACT UNITED TRANSPORT SERVICES EMERGENCY RESPONSE COORDINATOR: 1800 033 111 (24 HOURS)</p>
--

Rear of box

Directions for Use

Crops	Weeds Controlled	State	Rate kg / ha	Critical Comments	
LETTUCE (direct sown)	Barnyard Grass, Blackberry Nightshade, Chickweed, Portulaca, Winter grass, Rye grass, Summer Grass, Wireweed, Prince of Wales feather, Shepherd's Purse, Nettles.	All States	4.5 in 330 – 550 L	Apply pre-emergent, immediately after sowing. For best results, prepare the seed bed free from clods. Sufficient irrigation to wet the top 3 to 4 cm of soil should be applied within 24 hours	
Transplanted				Apply immediately after transplanting. For best results, prepare the seed bed free from clods. Sufficient irrigation to wet the top 3 to 4 cm of soil should be applied within 24 hours	
			Vic only.	2.3 in 300 – 550 L	Apply on light sandy soils, low in organic matter
Turf & Lawns Common couch	Winter grass	All States	1.2 (12g / 100 m ²)	Rate/Bowling Green 37 x 37 m 170 g	Apply during the Autumn and Winter, if established turf is infested with seedling winter grass. Ensure a uniform coverage. Irrigate immediately after application with 12 – 25 mm irrigation or rain and again 24 hour later. Repeat treatment if necessary on Qld blue couch after 4 weeks. DO NOT RESEED FOR 60 DAYS AFTER TREATMENT Use ONLY on established turf. DO NOT use on bluegrass, ryegrass, fescue or bent grass lawns.
Queensland blue couch				Qld only	
Buffalo grass		W A only	1.2 (12g / 100 m ²)	170 g	
Lucerne, Clover, Medics etc. Grown for forage, Hay or Seed production	Annual Ryegrass, Winter Grass, Barley Grass, Silver grass. Prairie Grass, Canary Grass, Great Brome, Squirrel tail Fescue and Rats tail Fescue,	NSW, ACT, S A, W A, Tas only	1-1.5	SEEDLING CROPS: Apply after the third trifoliate leaf is fully developed. ESTABLISHED CROPS: Apply any time before the end of July WEEDS: Apply when weeds are at post or Pre-emergent stages	
	Wild oats, Perennial Ryegrass, Bent grass, English couch, Yorkshire fog grass, Chickweed, Seedling Sorrel, Black- berry nightshade, Patterson's curse, seedling Wireweed		1.5 - 2	BEST RESULTS: When applied during cool and moist conditions. Rain or irrigation is necessary to thoroughly wet the top 3 to 4 cm of soil. This is similar to 25 mm rain. Use rates towards the higher end of the range on heavy soils, if conditions are not optimal or where a heavy grass population is expected. Grazing or cutting PRIOR to treatment is recommended.	

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORIZED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD

DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD OR FOR SEED FOR 25 DAYS AFTER APPLICATION

NOT CLASSIFIED AS DANGEROUS GOODS FOR TRANSPORT BY ROAD AND RAIL

Labelling for **every** measure pack

CAUTION

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

BURST™ Herbicide

ACTIVE CONSTITUENT: 500 g/kg PROPYZAMIDE

MEASURE ONLY - ILLEGAL TO BE SOLD
EXCEPT IN UNOPENED PRIMARY PACK

WATER SOLUBLE PACKAGING. KEEP DRY

BEFORE USE: READ ALL DIRECTIONS ON THE LABEL
OF THE PRIMARY PACK

Net Contents: 500 g

Sipcam Pacific Australia Pty Limited
Suite 11, 23-31 Gheringhap Street
GEELONG VIC 3220
ACN 072 176 888
TEL:03 5223 3746

NRA Approval No.: 55783/0602 