

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

TERRACLOR®

SOIL FUNGICIDE

ACTIVE CONSTITUENT: 750g/kg QUINTOZENE

GROUP **Y** FUNGICIDE

For control of soil borne diseases of vegetables, cotton,
turf, peanuts and ornamentals as specified in the
DIRECTIONS FOR USE table.

CONTENTS: 2.5 kg



BARMAC INDUSTRIES

ABN 21 009 674 953

3 Mary Street
Blackstone Qld 4304
Ph: (07) 3280 3000

Batch
Number:

Date of
Manufacture:

Terraclor2.5L0803V1

Name of file: Terraclor.pdf Label Dimensions: 160mm (w) x 160 mm (h) Scale of print: A4 - 100% A3 - 100%
Colours: Pantone 814 (left side of label) Pantone 811 (right side of label) Reflex Blue (logo) Black

DIRECTIONS FOR USE

Crop	Disease	States	Rate	Critical Comments
TOMATO	Southern Blight (<i>Asterium ralskyi</i>)	QLD, NSW, ACT, WA, SA & TAS	300-500 g in 100 L water	Apply as a drench at transplanting using 250 mL of solution per plant. Use higher rate for heavier soils and higher disease levels.
LETTUCE	Bottom Rot (Rhizoctonia)	All States	100-150 g in 100 L water	Apply 100 L of solution to 20 m ² after thinning when plants are 5-7 cm high. Use higher rate for heavier soils and higher disease levels.
BEANS	Stem & Root Rot (Rhizoctonia)		16.5-22 kg/ha	Mix in a convenient volume of water and apply in a 20-30 cm band over the row. Incorporate into the top 5-10 cm of soil prior to planting. Use higher rate for heavier soils and higher disease levels.
PEANUTS	Stem Rot (<i>Asterium ralskyi</i>)	QLD, NSW, ACT & WA		For row application at pegging time, direct spray to centre of row (30 cm band) so as to cover crown of plant and pegging zone. Use higher rate for heavier soils and higher disease levels.
COTTON	Damping Off (Rhizoctonia)		5 kg/ha	Mix in a convenient volume of water and spray over seed and surrounding soil at planting.
POTATO	Black Scurf (<i>Rhizoctonia solani</i>)	All States	33-44 kg/ha	Row spray application. Spray onto soil in a 30 cm band over row prior to planting time. Cultivate to 10-15 cm and cover seed with treated soil only. Use higher rate for heavier soils and higher disease levels.
CABBAGE, CAULIFLOWER, BROCCOLI	Club Foot (Plasmopthora) Wire Stem & Black Rot		16.5-22 kg/ha	Spray in a 20-30 cm band over rows and incorporate to depth of 10 cm prior to planting out. Use higher rate for heavier soils and higher disease levels.
IN SEED BED			500-750 g per 100 m ²	Mix in a convenient volume of water, spray or drench and incorporate to depth of 10 cm. Use higher rate for heavier soils and higher disease levels.
TURF	Brown Patch (Rhizoctonia)		750 g in 200 L water per 100 m ²	Mix in 200-500 L water and apply evenly when disease appears (usually in hot weather). Repeat application in 14 days.
BULBOUS IRIS	Crown Rot Black Rot		150 g per 10 m ²	Mix in a convenient volume of water. Spray or drench evenly over area and incorporate to 10-15 cm prior to planting.
LARKSPUR, CALENDULA, SWEET PEA, SNAPDRAGONS	Stem Rot (Rhizoctonia)		120 g per 10 m ²	Mix in a convenient volume of water. Spray or drench evenly over area and incorporate to 10-15 cm prior to planting.
CARNATION, POINSETTA, CHRYSANTHEMUM, BEGONIA			500-700 g per 100 m ²	Either apply to seed bed as a drench in 200-500 L of water or as a dust, then incorporate into the top 7-10 cm soil. Use higher rate for heavier soil and higher disease levels.
APPLES	Collar Rot (<i>Asterium ralskyi</i>)	QLD, SA, WA & TAS	1 kg in 100 L water. Apply 500 mL solution per tree.	Drench the seedling butt and planting site with 500 mL of the mixture.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHOLDING PERIOD: VEGETABLES AND PEANUTS: DO NOT HARVEST FOR 4 WEEKS AFTER APPLICATION. OTHER CROPS: NOT REQUIRED WHEN USED AS DIRECTED.

GENERAL INSTRUCTIONS

This product may be applied to the soil and mixed into the top 10-15 cm or mixed with water and applied to the soil surface.

Fungicide Resistance Warning:

GROUP Y FUNGICIDE
For fungicide resistance management, Terracolor Soil Fungicide is a Group Y fungicide.

Some naturally occurring individual fungi resistant to Terracolor Soil Fungicide and other Group Y fungicides may exist through normal genetic variability in any fungal population. The resistant

individuals can eventually dominate the fungal populations if these fungicides are used repeatedly. These resistant fungi will not be controlled by Terracolor Soil Fungicide or other Group Y fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence or resistant fungi is difficult to detect prior to use, Barnac Industries Pty Ltd accepts no liability for any losses that may result from the failure of Terracolor Soil Fungicide to control resistant fungi.

PROTECTION OF LIVESTOCK:

DO NOT graze any treated area or cut for stock food.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT:

Dangerous to fish. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL:

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If no landfill is available, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS:



Avoid contact with eyes and skin. Wash hands after use.

FIRST AID:

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131 126.

MATERIAL SAFETY DATA SHEET:


For additional information, refer to the Material Safety Data Sheet.



NOTICE: This product is designed only for the particular purposes indicated on the label. Failure to use the product strictly as directed may be illegal, prove dangerous and render the product ineffective. The buyer or user is responsible for any misuse or negligence in the handling, storage, use or disposal of this material. Barnac Industries Pty Ltd will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply and use of this product other than those responsibilities mandatorily imposed by Statutes applicable to the sale and supply of these goods.

UN NO. 3077	ENVIRONMENTALLY HAZARDOUS SUBSTANCES, SOLID, N.O.S. (CONTAINS QUINTOZENE) MARINE POLLUTANT
In a Transport Emergency Dial 000 Police or Fire Brigade	
P.G.-III	HAZCHEM 2 X

©Terracolor is a trademark of Amvac Chemical Corporation.



APVMA Approval No. 55633/10/1206
Terracolor 25L0803V1

Terracolor 2.5kg Back Size: Height 160mm deep X 255mm Wide Prints at 100% A4 100% A3