

E-LABEL SUBMISSION

UNITED FARMERS SANCTION 99.5EC HERBICIDE - 5/10Litres

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT plant cereal crops into the treated area for a period of 18 weeks after application of United Farmers Sanction 99.5EC Herbicide.
DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping land or pastures.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

DO NOT store tank-mixes of United Farmers Sanction 99.5EC Herbicide with other materials. Store in the closed original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, relace cap and return clean container to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the container below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Will irritate the eyes and skin. When preparing spray wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and face shields or goggles. If product in eyes, wash out immediately with water. Wash hands after use. After each day's use wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (PHONE AUSTRALIA: 13 11 26). If swallowed, do NOT induce vomiting. Give a glass of water.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet, which is available from the supplier.

CONDITIONS OF SALE

The use of United Farmers Sanction 99.5EC Herbicide being beyond the control of the manufacturer no warranty expressed or implied is given by United Farmers Cooperative regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and United Farmers Cooperative accepts no responsibility for any consequence whatsoever resulting from the use of this product.



UN No: 3082	BIPYRIDILIUM PESTICIDES LIQUID, TOXIC, N.O.S. (contains Quizalofop-p-ethyl)
In a transport Emergency Dial 000 Police or Fire Brigade	SPECIALIST ADVICE IN AN EMERGENCY ONLY 1800 033 111
PG II	HAZCHEM 2X



POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



United Farmers
SANCTION 99.5EC
HERBICIDE

ACTIVE CONSTITUENT: 99.5 g/L QUIZALOFOP-P-ETHYL
SOLVENT: 801 g/L HYDROCARBON LIQUID

GROUP **A** HERBICIDE

For the control of certain grass weeds in broadleaf crops as shown in the Directions for Use table.

IMPORTANT
READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT

Batch Number:

NET CONTENTS APVMA Approval Number

5 Litres

58610/5L/0208

Date of Manufacture:

10 Litres

58610/10L/0208



United Farmers Co-operative Company Limited
2 Birksgate Road, Rous Head,
PMB 10 North Fremantle, Western Australia 6159
Telephone 08 9430 2222 Facsimile 08 9430 2233



100% ACTUAL SIZE
(388mm x 160mm)

E-LABEL SUBMISSION

UNITED FARMERS SANCTION 99.5EC HERBICIDE - 20Litres

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT plant cereal crops into the treated area for a period of 18 weeks after application of United Farmers Sanction 99.5EC Herbicide.
DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping land or pastures.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

DO NOT store tank-mixes of United Farmers Sanction 99.5EC Herbicide with other materials. Store in the closed original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, relace cap and return clean container to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the container below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Will irritate the eyes and skin. When preparing spray wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and face shields or goggles. If product in eyes, wash out immediately with water. Wash hands after use. After each day's use wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (PHONE AUSTRALIA: 13 11 26). If swallowed, do NOT induce vomiting. Give a glass of water.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet, which is available from the supplier.

CONDITIONS OF SALE

The use of United Farmers Sanction 99.5EC Herbicide being beyond the control of the manufacturer no warranty expressed or implied is given by United Farmers Cooperative regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and United Farmers Cooperative accepts no responsibility for any consequence whatsoever resulting from the use of this product.



UN No: 3082	BIPYRIDILIUM PESTICIDES LIQUID, TOXIC, N.O.S. (contains Quizalofop-p-ethyl)
In a transport Emergency Dial 000	SPECIALIST ADVICE IN AN EMERGENCY ONLY 1800 033 111
Police or Fire Brigade	
PG II	HAZCHEM 2X

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

United Farmers
SANCTION 99.5EC
HERBICIDE

ACTIVE CONSTITUENT: 99.5 g/L QUIZALOFOP-P-ETHYL
SOLVENT: 801 g/L HYDROCARBON LIQUID

GROUP **A** HERBICIDE

For the control of certain grass weeds in broadleaf crops as shown in the Directions for Use table.

IMPORTANT
READ THE ATTACHED LEAFLET
BEFORE USING THIS PRODUCT

CONTENTS 20 Litres


Batch Number:

APVMA Approval Number
58610/20L/0208



United Farmers Co-operative Company Limited
2 Birksgate Road, Rous Head,
PMB 10 North Fremantle, Western Australia 6159
Telephone 08 9430 2222 Facsimile 08 9430 2233

Date of Manufacture:



100% ACTUAL SIZE
(388mm x 160mm)

E-LABEL SUBMISSION

UNITED FARMERS SANCTION 99.5EC HERBICIDE - 110/200/1000Litres



PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT plant cereal crops into the treated area for a period of 18 weeks after application of United Farmers Sanction 99.5EC Herbicide.
DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping land or pastures.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

DO NOT store tank-mixes of United Farmers Sanction 99.5EC Herbicide with other materials. Store in the closed original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, relace cap and return clean container to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the container below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Will irritate the eyes and skin. When preparing spray wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and face shields or goggles. If product in eyes, wash out immediately with water. Wash hands after use. After each day's use wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (PHONE AUSTRALIA: 13 11 26). If swallowed, do NOT induce vomiting. Give a glass of water.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet, which is available from the supplier.

CONDITIONS OF SALE

The use of United Farmers Sanction 99.5EC Herbicide being beyond the control of the manufacturer no warranty expressed or implied is given by United Farmers Cooperative regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and United Farmers Cooperative accepts no responsibility for any consequence whatsoever resulting from the use of this product.



UN No: 3082	BIPYRIDILIUM PESTICIDES LIQUID, TOXIC, N.O.S. (contains Quizalofop-p-ethyl)
In a transport Emergency Dial 000 Police or Fire Brigade	SPECIALIST ADVICE IN AN EMERGENCY ONLY 1800 033 111
PG II	HAZCHEM 2X

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

United Farmers
SANCTION 99.5EC
HERBICIDE

ACTIVE CONSTITUENT: 99.5 g/L QUIZALOFOP-P-ETHYL
SOLVENT: 801 g/L HYDROCARBON LIQUID

GROUP **A** HERBICIDE

For the control of certain grass weeds in broadleaf crops as shown in the Directions for Use table.

IMPORTANT
READ THE ATTACHED LEAFLET
BEFORE USING THIS PRODUCT

Batch Number:

NET CONTENTS APVMA Approval Number


- 110 Litres 58610/110L/0208
- 200 Litres 58610/200L/0208
- 1000 Litres 58610/1000L/0208

Date of Manufacture:



United Farmers Co-operative Company Limited
2 Birksgate Road, Rous Head,
PMB 10 North Fremantle, Western Australia 6159
Telephone 08 9430 2222 Facsimile 08 9430 2233

100% ACTUAL SIZE
(388mm x 160mm)



ACTIVE CONSTITUENT: 99.5 g/L QUIZALOFOP-P-ETHYL
SOLVENT: 801 g/L HYDROCARBON LIQUID

United Farmers
SANCTION 99.5EC
HERBICIDE

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

GROUP **A** HERBICIDE

DIRECTIONS FOR USE

RESTRAINT:

DO NOT apply to crops that are stressed by any cause (such as severe cold weather, disease, drought or waterlogging).
DO NOT apply to weeds that are stressed by any cause (such as weather conditions, drought, waterlogging) or not actively growing. Grass weeds under stress frequently become less susceptible to herbicide activity. This may result in an incomplete kill or suppression of weeds only.
DO NOT apply under conditions of high temperature and low humidity.
DO NOT apply if rainfall is expected within 3 hours.
DO NOT apply less than 50L prepared spray/ha.

BOOM SPRAYING: WINTER GROWING WEEDS: ALL STATES

CROP	WEEDS CONTROLLED	WEEDS STAGE	RATE/ha	WITHHOLDING PERIOD		CRITICAL COMMENTS
				Harvest	Grazing	
Beetroot Cabbage Canola Carrots Cauliflower Chickpeas Clover & Sub-clover pastures Cucumbers Faba Beans Field Peas Honey Dew Melons Lentils Lucerne Lupins Medic pasture Mung beans Navy Beans Onions Potatoes Pumpkins Radish Tomatoes Vetch	Annual Ryegrass (<i>Lolium rigidum</i>) Brome Grasses (<i>Bromus</i> spp.)	3 leaf to early tillering	300mL or 375mL	14 days	-	Apply when weeds are actively growing. Use higher rate under heavy weed pressure, and/or when weeds have commenced tillering*.
				9 weeks	4 weeks	
				11 weeks	-	
				10 weeks	-	
Barley grass (<i>Hordeum</i> spp) Volunteer Barley Volunteer Wheat	3 leaf to early tillering	250mL	14 days	-	4 weeks	Apply when weeds are actively growing*.
				12 weeks	4 weeks	
				9 weeks	4 weeks	
				9 weeks	4 weeks	
Wild Oats (<i>Avena</i> spp)	3 leaf to mid tillering	125mL or 250mL	12 weeks	4 weeks	Apply when weeds are actively growing. Use the lower rate only under light weed pressure and/or when weeds are at the 1-3 leaf stage. Use higher rate under heavy weed pressure and/or when weeds are tillering. The higher rate will give better or more reliable control*.	
			6 weeks	4 weeks		
			12 weeks	4 weeks		
			12 weeks	4 weeks		
			18 weeks	4 weeks		
			10 weeks	-		
			9 weeks	-		
			9 weeks	-		
			21 days	-		
			4 weeks	-		
			12 weeks	28 days		

* Always add a surfactant/wetting agent at the recommended Rate see section on "use of surfactant/wetting agent" for details.

• Important: DO NOT apply United Farmers Sanction 99.5EC Herbicide before the five true leaf stage of the crop.

BOOM SPRAYING: SUMMER GROWING WEEDS: ALL STATES

CROP	WEEDS CONTROLLED	WEEDS STAGE	RATE/ha	WITHHOLDING PERIOD		CRITICAL COMMENTS
				Harvest	Grazing	
Beetroot Cabbage Cauliflower Carrots Cucumbers Green Beans Honey Dew Melons Lucerne Mung beans Navy Beans Onions Peanuts Potatoes Pineapples Pumpkins Radish Soybeans Sunflower Tomatoes	Awnless Barmyard Grass (<i>Echinochloa colona</i>) Barmyard Grass (<i>Echinochloa crusgalli</i>)	3 leaf to early tillering	500mL or 750mL	14 days	-	Apply when weeds are actively growing. Use higher rate under heavy weed pressure, and/or when weeds have commenced tillering*.
				9 weeks	-	
				14 days	-	
				10 weeks	-	
Couch grass (<i>Cynodon dactylon</i>)	2-4 leaf seedling	500mL	5 weeks	-	4 weeks	Apply when weeds are actively growing. This rate will only control seedling Couch Grass which is less than 2 months old and established from seed*.
				9 weeks	4 weeks	
				12 weeks	4 weeks	
				12 weeks	4 weeks	
Dinebra (<i>Dinebra retroflexa</i>)	3 leaf to early tillering	250mL or 375mL	12 weeks	-	4 weeks	Apply when weeds are actively growing. Use higher rate under heavy weed pressure, and/or when weeds have commenced tillering*.
				18 weeks	4 weeks	
				11 weeks	11 weeks	
				10 weeks	-	
Crowfoot grass (<i>Eleusine indica</i>)	2 leaf to early tillering	500mL	7 days	-	-	Apply when weeds are actively growing*.
				12 weeks	-	
				21 days	-	
				12 weeks	4 weeks	
Foxtail Millet (<i>Setaria italica</i>)	2 leaf to early tillering	250mL or 500mL	12 weeks	-	4 weeks	Apply when weeds are actively growing. Use higher rate under heavy weed pressure, and/or when weeds have commenced tillering*.
				9 weeks	11 weeks	
				9 weeks	-	
				4 weeks	-	
Columbus grass (<i>Sorghum x alatum</i>) Johnson grass (<i>Sorghum halepense</i>) Liverseed grass (<i>Orochloa panicoides</i>)	2 leaf to 3 tiller	500mL or 750mL	5 weeks	-	-	Apply when weeds are actively growing. Use higher rate under heavy weed pressure, and/or when weeds have commenced tillering*.
				9 weeks	-	
				12 weeks	-	
				12 weeks	-	

* Always add a surfactant/wetting agent at the recommended Rate see section on "use of surfactant/wetting agent" for details.

• Important: DO NOT apply United Farmers Sanction 99.5EC Herbicide before the five true leaf stage of the crop.

HAND GUN SPRAYING: ALL STATES

CROP	WEEDS CONTROLLED	WEEDS STAGE	RATE/100 L	WITHHOLDING PERIOD		CRITICAL COMMENTS
				Harvest	Grazing	
Beetroot Cabbage Cauliflower Carrots Canola Cucumbers Chickpeas Clover & Sub-clover pastures Cucumbers Faba Beans Field Peas Grapes Green beans Honey Dew Melons Lentils Lucerne Lupins Medic pasture Mung beans Navy Beans Non crop Onions Peanuts Pineapples Potatoes Pumpkins Radish Soy beans Sunflowers Tomatoes Vetch	Awnless Barmyard Grass (<i>Echinochloa colona</i>) Crowfoot grass (<i>Eleusine indica</i>)	3 leaf to early tillering	125mL	14 days	-	Apply when weeds are actively growing. Use a minimum of 800L of prepared spray/ha. Thoroughly wet target weeds (especially well established clumps) without causing runoff. Repeat spraying may be necessary for perennial grass species. Always add a surfactant/wetting agent at the recommended rate – see section on "Use of Surfactant/Wetting agent" for details.
				9 weeks	4 weeks	
				14 days	4 weeks	
				10 weeks	14 days	
Paspalum (<i>Paspalum dilatatum</i>)	Seedling to Established Clumps	Seedling to early tillering	250mL	14 days	-	Always add a surfactant/wetting agent at the recommended rate – see section on "Use of Surfactant/Wetting agent" for details.
				12 weeks	4 weeks	
				9 weeks	4 weeks	
				9 weeks	4 weeks	
Johnson grass (<i>Sorghum halepense</i>)	Seedling 2 leaf to early tillering	Rhizome 2 leaf to early tillering	250mL	5 weeks	-	Always add a surfactant/wetting agent at the recommended rate – see section on "Use of Surfactant/Wetting agent" for details.
				9 weeks	4 weeks	
				12 weeks	4 weeks	
				12 weeks	4 weeks	
Kikuyu (<i>Pennisetum clandestinum</i>)	Seedling to Established plants	Seedling to Established plants	250mL	6 weeks	-	Always add a surfactant/wetting agent at the recommended rate – see section on "Use of Surfactant/Wetting agent" for details.
				12 weeks	4 weeks	
				12 weeks	4 weeks	
				18 weeks	11 weeks	
Pineapples Potatoes Pumpkins Radish Soy beans Sunflowers Tomatoes Vetch	Kikuyu (<i>Pennisetum clandestinum</i>)	Seedling to Established plants	250mL	7 days	-	Always add a surfactant/wetting agent at the recommended rate – see section on "Use of Surfactant/Wetting agent" for details.
				9 weeks	-	
				9 weeks	-	
				21 days	4 weeks	
Summer grass (<i>Digitaria ciliaris</i>)	3 leaf to early tillering	Rhizome 2 leaf to early tillering	500mL or 1000mL	12 weeks	-	Always add a surfactant/wetting agent at the recommended rate – see section on "Use of Surfactant/Wetting agent" for details.
				12 weeks	4 weeks	
				18 weeks	4 weeks	
				11 weeks	11 weeks	

* Always add a surfactant/wetting agent at the recommended Rate see section on "use of surfactant/wetting agent" for details.

• Important: DO NOT apply United Farmers Sanction 99.5EC Herbicide before the five true leaf stage of the crop.

BOOM SPRAYING: SUMMER GROWING WEEDS: ALL STATES

CROP	WEEDS CONTROLLED	WEEDS STAGE	RATE/ha	WITHHOLDING PERIOD		CRITICAL COMMENTS
				Harvest	Grazing	
Beetroot Cabbage Cauliflower Carrots Cucumbers Green Beans Honey Dew Melons Lucerne Mung beans Navy Beans Onions Peanuts Potatoes Pineapples Pumpkins Radish Soy beans Sunflowers Tomatoes	Rhodes grass (<i>Chloris gayana</i>) Stink grass (<i>Eragrostis cilianensis</i>)	3 leaf to early tillering	250mL or 500mL	14 days	-	Apply when weeds are actively growing. Use higher rate under heavy weed pressure, and/or when weeds have commenced tillering*.
				9 weeks	-	
				14 days	-	
				10 weeks	-	
Queensland Blue grass (<i>Dichanthium sericeum</i>)	3 leaf to early tillering	500mL	5 weeks	-	4 weeks	Apply when weeds are actively growing*.
				9 weeks	4 weeks	
				9 weeks	4 weeks	
				12 weeks	4 weeks	
Summer grass (<i>Digitaria ciliaris</i>)	3 leaf to early tillering	500mL or 1000mL	12 weeks	-	Apply when weeds are actively growing. Use higher rate under heavy weed pressure, and/or when weeds have commenced tillering*.	
			12 weeks	4 weeks		
			18 weeks	4 weeks		
			11 weeks	11 weeks		
			10 weeks	-		
			7 days	-		
			9 weeks	-		
			9 weeks	-		
			21 days	4 weeks		
			12 weeks	11 weeks		
			9 weeks	-		
			4 weeks	-		

* Always add a surfactant/wetting agent at the recommended Rate see section on "use of surfactant/wetting agent" for details.

• Important: DO NOT apply United Farmers Sanction 99.5EC Herbicide before the five true leaf stage of the crop.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD

HARVEST

Beetroot, Cauliflower, Cucumbers: **DO NOT APPLY LATER THAN 14 DAYS AFTER APPLICATION.**

Cabbage, Field Peas, Honey Dew Melon, Pumpkins, Sunflowers: **DO NOT APPLY LATER THAN 9 WEEKS AFTER APPLICATION.**

Carrots, Potatoes: **DO NOT APPLY LATER THAN 10 WEEKS AFTER APPLICATION.**

Chickpeas, Faba Beans, Lentils, Mung Beans, Navy Beans, Soybeans, Vetch: **DO NOT APPLY LATER THAN 12 WEEKS AFTER APPLICATION.**

Canola, Peanuts: **DO NOT APPLY LATER THAN 11 WEEKS AFTER APPLICATION.**

Green Beans: **DO NOT APPLY LATER THAN 5 WEEKS AFTER APPLICATION.**

Lupins: **DO NOT APPLY LATER THAN 6 WEEKS AFTER APPLICATION.**

Onions: **DO NOT APPLY LATER THAN 18 WEEKS AFTER APPLICATION.**

Pineapples: **DO NOT APPLY LATER THAN 7 DAYS AFTER APPLICATION.**

Radish: **DO NOT APPLY LATER THAN 21 DAYS AFTER APPLICATION.**

Tomatoes: **DO NOT APPLY LATER THAN 4 WEEKS AFTER APPLICATION.**

GRAZING

Canola, Chickpeas, Faba Beans, Field Peas, Green Beans, Lentils, Lucerne, Lupins, Mung Beans, Navy Beans, Soybeans, Vetch: **DO NOT GRAZE OR CUT FOR STOCK FEED FOR 4 WEEKS AFTER APPLICATION.**

Clover, Medic Pastures (except Lucerne): **DO NOT GRAZE OR CUT FOR STOCK FEED FOR 14 DAYS AFTER APPLICATION.**

Peanuts, Sunflowers: **DO NOT GRAZE OR CUT FOR STOCK FEED FOR 11 WEEKS AFTER APPLICATION.**

GENERAL INSTRUCTIONS

United Farmers Sanction 99.5EC Herbicide is an emulsifiable concentrate herbicide used post emergence for the control of certain grass weeds in specified broadleaf crops. United Farmers Sanction 99.5EC Herbicide is a systemic herbicide, which is rapidly absorbed and readily translocates to the treated foliage to the root system and growing points of the plant. Applications made to grass weeds stressed due to insufficient moisture or cold temperatures may result in unsatisfactory control.

RESISTANT WEEDS WARNING

GROUP **A** HERBICIDE

United Farmers Sanction 99.5EC Herbicide is a member of the "Fop" (aryloxyphenoxy propanoate) group of herbicides. United Farmers Sanction 99.5EC Herbicide has the inhibitors of acetyl coA carboxylase mode of action. For weed resistance management United Farmers Sanction 99.5EC Herbicide is a Group A herbicide. Some naturally occurring weed biotypes resistant to United Farmers Sanction 99.5EC Herbicide and other Group A herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by United Farmers Sanction 99.5EC Herbicide or other Group A herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, United Farmers Cooperative accepts no liability for any losses that may result from failure of United Farmers Sanction 99.5EC Herbicide to control resistant weeds. Strategies to minimise the risk of herbicide resistance are available. Consult your farm chemical supplier, consultant, local Department of Agriculture or Primary Industries.

TIME OF APPLICATION

For maximum benefit, weeds should be sprayed when young and actively growing. Application may be made at any stage of crop growth (except to cucurbit crops), but sufficient time must be allowed for the prescribed withholding period to expire before harvest and/or grazing or cutting for stockfeed.

IMPORTANT

DO NOT apply United Farmers Sanction 99.5EC Herbicide to cucurbit crops before the five true leaf stage.

SPRAY PREPARATION

USE OF A SURFACTANT/WETTING AGENT

United Farmers Sanction 99.5EC Herbicide is an emulsifiable concentrate to be mixed with water and applied as a spray. Add the amount of United Farmers Sanction 99.5EC Herbicide directly to the spray tank and mix well. If tank mixing with benomyl or Mancozeb, add these to the spray tank and mix thoroughly PRIOR to adding United Farmers Sanction 99.5EC Herbicide.

United Farmers Sanction 99.5EC Herbicide may be used with either of the following adjuvant system:

1) A non-ionic surfactant (1000 gai/L strength) at 100mL/100L (i.e. 0.1% v/v) and a mineral spray oil at 1L/100L of final spray volume (i.e. 1% v/v).

2) A non-ionic surfactant (1000 gai/L strength) at 200mL/100L of final spray volume (i.e. 0.2% v/v).

DO NOT use the non-ionic surfactant + mineral oil adjuvant system on vegetable crops, use a non-ionic surfactant only.

DO NOT use a spray oil alone, a surfactant must always be added. DO NOT use non-methylated spray oils of crop origin as spray adjuvants.

COMPATIBILITIES

United Farmers Sanction 99.5EC Herbicide is compatible with the insecticides deltamethrin, omethoate, dimethoate, methomyl, endosulfan, the fungicides carbendazim and mancozeb and the herbicides diflufenican and clopyralid **when using the surfactant only adjuvant system.**

United Farmers Sanction 99.5EC Herbicide is compatible with the herbicide sethoxydim and clethodim **when using the surfactant plus spray oil adjuvant system.**

When tank mixing United Farmers Sanction 99.5EC Herbicide and broadleaf herbicides, DO NOT use a spray oil, use a surfactant only.

Application of United Farmers Sanction 99.5EC Herbicide /diflufenican tank mix may result in crop effects. The effect is temporary and usually appears as yellow or white bleaching on the leaves. New growth and subsequent yield, however, will not be affected.

DO NOT add crop oil to tank mixtures of United Farmers Sanction 99.5EC Herbicide /diflufenican.

To minimise crop effects apply United Farmers Sanction 99.5EC Herbicide a minimum of 5 days prior to diflufenican application or a minimum 10 days after diflufenican application (14-21 days if weeds have been stressed).

GROUND SPRAYING

Use sprayers with flat fan or hollow cone nozzles at 210-350kPa to give an even distribution of droplets with no less than 50% of the spray volume in the 150-300 micron range. 110° flat fan nozzles are preferred.

Use between 50 and 100 litres of prepared spray/ha, the higher spray volume gives more reliable results. If weeds are large and the crop canopy is dense, increase the volume to 150 litres prepared spray/ha.

For proper spray coverage adjust the boom and nozzle height according to the specification listed by the nozzle manufacturers. Ensure the spray is evenly distributed and a double overlap spray pattern is obtained.

Avoid overlapping boom spray widths and shut off spray booms while starting, slowing or stopping.

SPRAYER CLEANUP

Thoroughly clean all traces of United Farmers Sanction 99.5EC Herbicide from application equipment immediately after use. Flush tank, pump, hoses and boom with several changes of water after removing nozzle tips and screens (clean those parts separately). Failure to thoroughly clean the equipment may result in injury to subsequently sprayed cereal and grass crops.

If spray equipment has been previously used for the application of sulfonylurea herbicides eg metsulfuron methyl, thifensulfuron methyl, chlorsulfuron, triasulfuron or terbuthryn, ensure that the recommended sprayer cleanup procedure of the product previously sprayed has been followed before using the equipment for the application of United Farmers Sanction 99.5EC Herbicide.

PRECAUTION: RE-ENTRY

DO NOT allow entry into treated areas until the spray has dried. When prior entry is necessary, wear chemical resistant gloves. Clothing must be laundered after each day's use.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT plant cereal crops into the treated area for a period of 18 weeks after application of United Farmers Sanction 99.5EC Herbicide.

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping land or pastures.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

DO NOT store tank-mixes of United Farmers Sanction 99.5EC Herbicide with other materials.

Store in the closed original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, relace cap and return clean container to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the container below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots.

Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Will irritate the eyes and skin. When preparing spray wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and face shields or goggles. If product in eyes, wash out immediately with water. Wash hands after use. After each day's use wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (PHONE AUSTRALIA: 13 11 26). If swallowed, do NOT induce vomiting. Give a glass of water.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet, which is available from the supplier.

CONDITIONS OF SALE

The use of United Farmers Sanction 99.5EC Herbicide being beyond the control of the manufacturer no warranty expressed or implied is given by United Farmers Cooperative regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and United Farmers Cooperative accepts no responsibility for any consequence whatsoever resulting from the use of this product.



UN No: 3082	BIPYRIDILUM PESTICIDES LIQUID, TOXIC, N.O.S. (contains Quizalofop-p-ethyl)
In a transport Emergency Dial 000 Police or Fire Brigade	SPECIALIST ADVICE IN AN EMERGENCY ONLY 1800 033 111
PG II	HAZCHEM 2X

