

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Crop Care
Simazine
Flowable Herbicide

ACTIVE CONSTITUENT: 500 g/L SIMAZINE



Controls weeds in Orchards, Chickpeas, Vineyards, Asparagus, Field Lupins, TT Canola, Berry Fruits, Gladioli, Hops, Faba Beans, Almonds (SA only), non-agricultural situations and other crops, in certain states as specified in the Directions for Use table.

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE

CONTENTS: 20, 110, 200 & 1000 Litres

Crop Care Australasia Pty Ltd 77 Tingira Street Pinkenba Qld 4008

STORAGE AND DISPOSAL (20L, 200L)

KEEP OUT OF REACH OF CHILDREN. Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse the containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury the empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should NOT be burnt.

(110 & BULK ONLY)

KEEP OUT OF REACH OF CHILDREN. Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage. Do NOT dispose of undiluted chemicals on-site.

SAFETY DIRECTIONS

Avoid contact with eyes and skin. DO NOT inhale spray mist. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. (Tel Australia 13 11 26)

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet which can be obtained from the supplier.

CONDITIONS OF SALE

Crop Care Australasia Pty Ltd will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than the responsibility for the merchantable quality of these goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by these Statutes the liability of Crop Care Australasia Pty Ltd is limited to the replacement of the goods or (at the option of Crop Care Australasia Pty Ltd) the refund of the price paid and is conditional upon a claim being made in writing and where possible sufficient part of the goods to enable proper examination being returned to Crop Care Australasia Pty Ltd within Thirty days of delivery.

Bar codes to be inserted here

APVMA Approval No: 52582/200L/0205
NRA Approval No: 52582/1101 (for 20L, 110L, 1000L)

BN

DOM

In a Transport Emergency Dial 000 Police or Fire Brigade	SPECIALIST ADVICE IN EMERGENCY ONLY 1800 033 498 ALL HOURS AUSTRALIA WIDE
--	---



Crop Care Australasia Pty Ltd
ACN 061 362 347
77 Tingira Street
PINKENBA QLD 4008

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Crop Care
Simazine

Flowable Herbicide



ACTIVE CONSTITUENT: 500 g/L SIMAZINE

Controls weeds in Orchards, Chickpeas, Vineyards, Asparagus, Field Lupins, TT Canola, Berry Fruits, Gladioli, Hops, Faba Beans, Almonds (SA only), non-agricultural situations and other crops, in certain states as specified in the Directions for Use table.

IMPORTANT: READ THIS LEAFLET BEFORE USE

Crop Care Australasia Pty Ltd 77 Tingira Street Pinkenba Qld 4008

CROP/ SITUATION	WEEDS CONTROLLED	STATE	RATE			CRITICAL COMMENTS
			Per Hectare	Per 100 Litres	Per 15 Litre Knapsack	
Canola – (Triazine Tolerant varieties only) Pre emergence or post sowing pre emergence only	Capeweed, Charlock, Clover, Corn Cromwell, Doublegee, Fumitories, Geraniums, Ivy-leaf Speedwell, London Rocket, Mustards, , Turnips, Paterson's Curse, Shepherd's Purse, Silver Grass (Vulpia). Suppression of Annual Ryegrass, Barley Grass, Brome Grass, Wild Oats and Wild Radish	All States	2 to 4 Litres			This use is subject to an <i>Integrated Weed Management Strategy</i> for the use of triazine herbicides in triazine tolerant (TT) canola. See General Instructions: <i>Integrated Weed Management Strategy for TT-Canola.</i> Can be applied up to a week before sowing or post-sowing pre- emergence (ideally Incorporated by harrows). For best results, apply to bare moist soil, either immediately before seeding or as a pre-emergence treatment at or within 7 days of planting. Sufficient rainfall (20-30 mm) to wet the soil through the weed root zone is necessary within 2-3 weeks of application. Application should not be made to ridged or excessively cloddy soil. When applied before seeding, incorporate to a depth of 5 cm.
Faba Beans	Annual Ryegrass, Barley Grass, Brome Grass, Capeweed, Corn Gromwell (Sheepweed), Fumitories, Geranium, Ivyleaf Speedwell, Mustards, Soursob, Wireweed and suppression of Wild Oats	SA only	2 to 2.5 litres			Apply either pre-seeding or immediately post-sowing which is preferred on light soils. Sow the crop at least 5cm deep. Use the lowest rate on light soils. Application should not be made to ridged or excessively cloddy soil. For fully reliable results significant rainfall, 20-30mm is necessary within 2-3 weeks of sowing.
			1 to 1.5 litres plus 1 litre of a Trifluralin 400g/L Selective Herbicide			Apply to bare moist soil and incorporate to a depth of 5cm just prior to sowing. Incorporation should be made within 4 hours of application. Use this mixture where Annual Ryegrass and Wild Oats are the major problem. Application should not be made to ridged or excessively cloddy soil. For fully reliable results significant rainfall, 20-30mm is necessary within 2-3 weeks of application. Do NOT use rates higher than 2L/ha on soils with pH 8.0 and above as unacceptable crop damage may occur.
Dams, Tanks, Troughs	Filamentous Blue- green Algae	WA only	4mL per 1000 litres of water			Mix in a convenient amount of water and apply when Algae development is first noticed.

CROP/ SITUATION	WEEDS CONTROLLED	STATE	RATE			CRITICAL COMMENTS
			Per Hectare	Per 100 Litres	Per 15 Litre Knapsack	
Non-agricultural areas such as Footpaths, Factory Sites, Oil Installations, Fence Lines, Road Shoulders, Power Stations, Gutters, Drains, Headlands, Driveways, Railway Tracks, Aerodromes, Public Utilities.	Annual Ryegrass, Annual Thistles, Barley Grass, Barnyard Grass, Bindy-eye, Brome Grass, Capeweed, Chickweed, Common Sowthistle, Creeping Oxalis, Fathen, Geranium, Ivyleaf Speedwell, Nettles, Potato Weed, Powell's Amaranth, Redroot Amaranth, Redshank, Shepherd's Purse, Slim Amaranth, Summer Grass, Turnips (not NSW), Wild Mustard, Wild Oats, Winter Grass, Wireweed (not Tas.) and suppression of Soursob	All States	8.8 litres	880mL	140mL	This rate gives seasonal maintenance control only. Apply to bare moist soil prior to rainy season to ensure herbicide is moved into the soil.
			18 to 44 litres	1.8 to 4.4 litres	260 to 600mL	Long residual control in summer rainfall areas. Apply to bare moist soil prior to rainy season to ensure herbicide is moved into the soil.
			44 to 88 litres	4.4 to 8.8 litres	600mL to 1.2 litres	Long residual control in summer rainfall areas. Apply to bare moist soil prior to rainy season to ensure herbicide is moved into the soil.
Asparagus	Annual Ryegrass, Annual Thistles, Barley Grass, Bindy-eye, Brome Grass, Capeweed, Chickweed, Common Sowthistle, Creeping Oxalis, Fathen, Geranium, Ivyleaf Speedwell, Nettles, Potato Weed, Powell's Amaranth, Redroot Amaranth, Redshank, Shepherd's Purse, Slim Amaranth, Turnips (not NSW), Wild Mustard,	All States	2.3 to 4.5 litres			Apply to bare moist soil after last cultivation and before spear emergence. Use the highest rate on heavy soils, also for Wild Oats. Apply to bare moist soil on established plants only. DO NOT apply to foliage or when fruit is present. Use the highest rate for Wild Oats.
Berry Fruits, (Boysenberries, Currants, Loganberries, Raspberries) Established for 12 months	Wild Oats, Winter Grass, Wireweed (not Tas.) and suppression of Soursob.					
Citrus Established for 12 months				110 to 220mL	25 to 45 mL	Apply to bare moist soil immediately after cultivation and before weed emergence.
Almonds Established for 3 years		SA only	1.7 to 3.5 litres			In almonds, lower rates (990mL to 1.7 litres per hectare) can be used in combination with other pre-emergence herbicides to enhance their broadleaf weed control.
Gladioli		All States	2.2 litres	110mL per approx. 500 square metres	25mL per approx. 100 square metres	Apply to bare moist soil after planting. High rates may cause crop damage on sandy soils low in organic matter.
Hops			2.3 to 4.5 litres			Apply to bare moist soil in late winter before hop emergence. Hops should be covered by 50mm of soil.

CROP/ SITUATION	WEEDS CONTROLLED	STATE	RATE			CRITICAL COMMENTS
			Per Hectare	Per 100 Litres	Per 15 Litre Knapsack	
Apples and Pears		Qld only	7 litres			Apply to bare moist soil. Use the highest rate in heavy soil.
		NSW, VIC, SA, WA, TAS only	3.2 to 4.5 litres	160 to 220 mL	30 to 45 mL	
Roses Established for 12 months		NSW, VIC, SA, WA, TAS only	2.3 to 4.5 litres	110 to 220mL	25 to 45mL -	
Vineyards Vines Established for 3 years in NSW, VIC, SA, WA, QLD and Tas.		Qld only	3.5 – 7 litres			Use lowest rates on sandy alkaline soils.
		NSW, Vic, SA, WA, Tas only	2.3 – 4.5 litres	110 to 220mL	25 to 45mL	Use lowest rates on sandy alkaline soils. In first year of use split applications are preferred.
Chickpeas	Deadnettle, Indian Hedge Mustard, Lesser Swinecress, Milk Thistle, Prickly Lettuce, Purple Goosefoot, Shepherd's Purse, Turnip Weed, Wireweed and suppression of Black Bindweed and Paradoxa grass.	NSW, QLD, SA only	1.5 litres plus 1.5 litres Prometryn Flowable liquid herbicide (500g/L)			Apply immediately post-planting. Application should not be made to ridged or excessively cloddy soil. For reliable results significant rain, 20-30mm, is necessary within 2-3 weeks of sowing.
Strawberries	Amaranthus (Redshank), Annual Ryegrass, Annual Thistles, Barley Grass, Bindy-eye, Brome Grass, Common Sowthistle, Capeweed, Chickweed, Creeping Oxalis, Fathen, Geranium, Ivyleaf Speedwell, Nettles, Potato Weed, Shepherd's Purse, Wild Mustard, Winter Grass, Wireweed (not Tas.), suppression of Wild Oats and Soursob.	All States	2.2 litres			Apply to bare moist soil between polythene covered beds.
Sub Clover Established lucerne and perennial grass pastures	Vulpia (Silvergrass), Rat's tail Fescue, Squirrel-tail Fescue, Sand Fescue.	NSW, SA & VIC only	1 to 1.6 litres			Apply 6-10 weeks after emergence of the pasture, but not before the 3rd trifoliate leaf stage of sub clover. Best results are obtained from application to young, well grazed, and actively growing plants. Some damage to sub clover may occur especially at the higher rates. DO NOT apply to sandy soils and use the lower rates on light textured soils. Apply under moist soil conditions; rain following application enhances results. DO NOT apply to stressed plants. Insect infestations should be controlled and sub clover allowed to recover before the Crop Care Simazine Flowable Herbicide is applied. DO NOT use herbicides for broad-leaved weed control within 3 weeks of application.

CROP/ SITUATION	WEEDS CONTROLLED	STATE	RATE			CRITICAL COMMENTS
			Per Hectare	Per 100 Litres	Per 15 Litre Knapsack	
Sub Clover Established lucerne and perennial grass pastures cont.		NSW, VIC only	1 to 1.5 litres plus 125 to 200mL of a 200g/L paraquat			Apply 6-10 weeks after emergence of the pasture, but not before the 3rd trifoliate leaf stage of sub clover. Best results are obtained from application to young, well grazed, and actively growing plants. However, the addition of paraquat improves the control of well established plants. Add a non-ionic surfactant at 0.2% v/v (200mL per 100L). Under conditions of good soil moisture, control of other grasses and some broad-leaved weed seedlings may occur. DO NOT apply to sandy soils and use the lower rates on light textured soils. Apply under moist conditions; rain following application enhances results. DO NOT apply to stressed plants. Insect infestations should be controlled and sub clover allowed to recover before the Crop Care Simazine plus paraquat is applied. DO NOT use herbicides for broad- leaved weed control within 3 weeks of application.
			Per Hectare	Per 100 Litres	Per 15 Litre Knapsack	
		SA only	750mL to 1 litre plus 125 to 200mL of 200g/L paraquat			
	Vulpia (Silvergrass), Rat's tail Fescue	WA only	750mL to 1 litre			Apply within 8 weeks of emergence of sub clover and grasses. Under conditions of good soil moisture control of other grasses and some broadleaf weeds may occur. Ensure that there is a good stand of subterranean clover present before spraying. Use the lower rate on light textured soils. DO NOT use broadleaf weed herbicides within 3 weeks of using Crop Care Simazine. DO NOT tank mix with other herbicides or add crop oils or wetting agents. DO NOT use on medics or red or white clover. DO NOT overlap when spraying, otherwise damage may be observed.

CROP/ SITUATION	WEEDS CONTROLLED	STATE	RATE			CRITICAL COMMENTS
			Per Hectare	Per 100 Litres	Per 15 Litre Knapsack	
Lupins	Annual Ryegrass, Barley Grass, Capeweed, Corn Gromwell, Fumitories, Geraniums, Ivyleaf Speedwell, Mustards, Paterson's Curse, Shepherd's Purse, Turnip Weed, Wild Turnip and Winter Grass. Suppression of Brome Grass, Soursob and Wild Oats.	NSW, VIC, SA, TAS only	1.5 to 2 litres on light soils 2.5 to 4 litres on loam soils 1 to 2 litres on light soils			Can be applied up to a week before sowing or post-sowing pre-emergent (ideally incorporated by harrows). Best results are achieved when application is made to bare moist soil and when significant rain to wet the soil through the weed root zone (20-30 mm) occurs within 2-3 weeks of application. Results can be variable if seasonal conditions are dry prior to sowing and lupins are sown into a dry or low moisture seedbed. Heavy, intense rainfall after application can cause crop damage. DO NOT apply to ridged or excessively cloddy soils. Apply 50 to 100 litres of spray per ha. Use a tank mix of Simazine and 1 litre of a 400g/L trifluralin where annual ryegrass and wild oats are the major problem. Incorporate the tank mixture to a depth of 5cm just prior to sowing. Incorporation of the tank mixture should be made within 4 hours of application.
Lupins When no weeds are present at time of sowing	Capeweed, Clover, Dock, Doublegee, Mustard, Radish, Self Sown Cereals, Silver Grass, Turnip and suppression of Barley Grass, Brome Grass, Ryegrass and Wild Oats.	WA only	2 to 3 litres on gravelly loam soils			For best results apply to bare moist soil, either immediately before seeding or as a pre-emergent treatment at or within 7 days of planting. Sufficient rainfall to wet the soil through the weed root zone (20-30mm) is necessary within 2-3 weeks of application. Results can be variable if seasonal conditions are dry prior to sowing and lupins are sown into a dry or low moisture seedbed. Application should not be made to ridged or excessively cloddy soil. When applied before seeding incorporation by the sowing operation should not be greater than 5cm. Use 1.0 to 1.5 L/ha on yellow sands and 2L/ha on all other types. DO NOT use on whitish or grey sands.

CROP/ SITUATION	WEEDS CONTROLLED	STATE	RATE			CRITICAL COMMENTS
			Per Hectare	Per 100 Litres	Per 15 Litre Knapsack	
Lupins When no weeds are present at time of sowing cont.			1 to 2 litres plus 1.5 litres of a Trifluralin 400 g/L Selective Herbicide			Can be applied up to a week before sowing or post-sowing pre-emergent (ideally incorporated by harrows). Best results are achieved when application is made to bare moist soil and when significant rain to wet the soil through the weed root zone (20-30 mm) occurs within 2-3 weeks of application. Results can be variable if seasonal conditions are dry prior to sowing and lupins are sown into a dry or low moisture seedbed. Heavy, intense rainfall after application can cause crop damage. DO NOT apply to ridged or excessively cloddy soils. Apply 50 to 100 litres of spray per ha. Use a tank mix of Simazine and 1 litre of a 400g/L trifluralin where annual ryegrass and wild oats are the major problem. Incorporate the tank mixture to a depth of 5cm just prior to sowing. Incorporation of the tank mixture should be made within 4 hours of application.
Lupins Where weeds are present at time of sowing			1 to 1.5 litres with the recommended rate of a knock- down herbicide			For best results apply to bare moist soil, either immediately before seeding or as a pre- emergent treatment at or within 7 days of planting. Sufficient rainfall to wet the soil through the weed root zone (20-30mm) is necessary within 2-3 weeks of application. Results can be variable if seasonal conditions are dry prior to sowing and lupins are sown into a dry or low moisture seedbed. Application should not be made to ridged or excessively cloddy soil. When applied before seeding incorporation by the sowing operation should not be greater than 5cm. Use 1.0 to 1.5 L/ha on yellow sands and 2L/ha on all other types. DO NOT use on whitish or grey sands.
Lupins post-emergent Top up application following a pre-emergent application of Crop Care Simazine	Capeweed, Clover, Dock, Doublegee, Mustard, Radish, Self Sown Cereals, Silvergrass, Turnip and suppression of Barley Grass, Brome Grass, Ryegrass and Wild Oats.	WA only	750mL to 2 litres			To be used in addition to a pre- emergent application of Crop Care Simazine Flowable Herbicide, but not as an alternative to a pre- emergent application. Apply to moist soil. Use rates of 750mL to 1L/ha within 4 weeks after seeding. DO NOT apply to Lupins which are showing symptoms of Simazine damage from the pre-emergent herbicide application.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD:

- CHICKPEAS:** DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 9 WEEKS AFTER APPLICATION.
FABA BEANS: DO NOT USE TREATED PRODUCE FOR HUMAN CONSUMPTION WITHIN 24 WEEKS OF TREATMENT. DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 8 WEEKS AFTER APPLICATION.
SUB CLOVER: DO NOT GRAZE FOR 14 DAYS AFTER APPLICATION. DO NOT CUT FOR STOCK FOOD FOR 21 DAYS AFTER APPLICATION.
CANOLA: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 15 WEEKS AFTER APPLICATION.

FOR OTHER CROPS: NOT REQUIRED WHEN USED AS DIRECTED

GENERAL INSTRUCTIONS

Crop Care Simazine Flowable Herbicide is a pre-emergent herbicide which selectively controls certain broadleaf weeds and grasses in asparagus, chickpeas, berry fruits, citrus, faba beans, almonds (SA only) gladioli, hops, pome fruits, roses, vineyards and lupins. In non-agricultural and other crop areas applied at higher rates, it will provide long control of a wide range of weeds and grasses. Established perennial species are not satisfactorily controlled. Since the product enters weeds mainly through their roots its effectiveness depends on rainfall or irrigation after application to move it down into the weed root zone.

Duration and effectiveness of control depends on the amount of chemical applied, soil type, rainfall and particular weed species.

When susceptible weeds start to appear uniformly the Crop Care Simazine Flowable Herbicide residue has probably dissipated.

Wash sprayer thoroughly with clean water after use.

INTEGRATED WEED MANAGEMENT STRATEGY FOR TT CANOLA

The use of this product in TT canola is subject to an *'Integrated Weed Management Strategy for the use of triazine herbicides in TT canola'* The Strategy encompasses:

- an *Integrated Weed Management (IWM)* strategy;
- the management of crop plant volunteers and outcrossing; and
 - triazine herbicide residue management (agronomic and environment).

Compliance with registered label directions and adoption of the principles outlined in the Code will assist with implementation of crop management practices that **minimise** the development of herbicide resistance in treated weeds; **reduce** the levels of triazine residues in the environment; and **manage** volunteer plants and outcrossing.

Copies of the *Strategy* are available from a Crop Care Australasia Pty Ltd representative or by contacting 1800 650 361 or (07) 3867 9100.

It is advised that consultation be undertaken with an appropriate agronomist, consultant or Departmental adviser prior to the use of Crop Care Atrazine Flowable Herbicide on TT Canola.

To minimise herbicide resistance:

- Avoid dry sowing in heavily weed infested paddocks. Wait for a weed germination after the opening rains in weedy paddocks. Use a pre-plant knockdown or cultivation. No weeds should be allowed to survive at this stage.
- Adapt the weed control program to the anticipated weed spectrum and pressure:

Broadleaf Weeds and Ryegrass: Use (simazine 500g/L) or (atrazine 500g/L) plus trifluralin pre-emergence. A follow-up with a Group A herbicide (if ryegrass is susceptible) or atrazine 500g/L may be necessary.

Broadleaf Weeds only: Use atrazine 500g/L post-emergence.

- **DO NOT** use atrazine 500g/L or simazine 500g/L if the area to be treated had a triazine herbicide applied to it last season.
- Watch for escapes, especially in paddocks with a long history of Group C herbicide use.
- **DO NOT** use Group C herbicides in consecutive years.

Resistant Weeds Reporting

Growers should collect plant or seed samples where weeds that are normally susceptible to atrazine and simazine may be resistant, get them tested and seek professional advice.

To avoid Triazine carry-over:

On acid soils (pH less than 6.5) - The maximum rate of atrazine 500g/L or simazine 500g/L or a combination of the two products to be applied to the crop during the growing season is 4 L/ha.

On alkaline soils (pH greater than 6.5) - The maximum rate of atrazine 500g/L or simazine 500g/L or a combination of the two products to be applied to the crop during the growing season is 2 L/ha.

Post-emergence use - It is recommended that atrazine 500g/L only be used, and at rates of 2 L/ha or less, on both acid or alkaline soils.

RESISTANT WEEDS WARNING

GROUP C HERBICIDE

Crop Care Simazine Flowable Herbicide is a member of the Triazine group of herbicides and has the inhibitors of Photosynthesis at Photosystem II mode of action. For weed resistance management this product is a group C Herbicide. Some naturally-

occurring weed biotypes resistant to Crop Care Simazine Flowable Herbicide and other inhibitors of Photosynthesis at Photosystem II herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other Inhibitors of Photosynthesis at Photosystem II herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use. Crop Care Australasia Pty Ltd accepts no liability for any losses that may result from the failure of Crop Care Simazine Flowable Herbicide to control resistant weeds.

MIXING

Settling may occur after storage for some weeks. Stir product or invert container several times before opening. Pour the flowable into the spray vat through a strainer to remove any dry particles or flakes, which can occasionally occur under hot storage conditions. Add the full quantity of Crop Care Simazine Flowable Herbicide to the partly filled spray tank while agitating. Fill tank and agitate to ensure thorough mixing. Continue agitation while spraying. Agitate vigorously from the bottom if allowed to stand. Reseal part used container immediately.

TT-Canola: Application

DO NOT apply to TT-canola by aircraft. Apply only with a low boom sprayer with a 60 m buffer zone downwind of treated fields to natural or impounded lakes or dams, and a 20 m buffer zone for any well, sink hole, intermittent or perennial stream. Apply only to areas where run-off is unlikely to occur or where run-off may be captured by farm earthworks.

Incorporation (Pre-plant and At Sowing application)

This product acts mainly by root absorption. Its effectiveness depends on the occurrence of rainfall or irrigation to move it down into the weed root zone. Sufficient rain or irrigation to thoroughly wet the soil through the weed root zone should occur or be made immediately after application to provide appropriate weed control. Delay in activation of the product may result in some weed growth.

In flood or furrow irrigation situations complete and continued activation of the product may not occur due to a thin band of dry soil on the surface during or after irrigation. Mechanical incorporation after application, using light harrows to incorporate the product into the soil not more than 4 cm deep is required to ensure the irrigation water activates the product.

Always apply the product to an even unridged seedbed.

COMPATIBILITY

Crop Care Simazine Flowable Herbicide is compatible with Flowable Gesagard, Roundup (glyphosate), Spinnaker, paraquat, diquat or paraquat-diquat mixtures, provided the mixture is agitated. If allowed to stand, agitate vigorously.

PROTECTION OF CROP, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT spray foliage of desirable plants, except lupins (WA only).

DO NOT make more than two applications during any one year in asparagus, berry fruit, citrus, almonds, gladioli, hops, apples, pears, roses, vineyards and lupins.

DO NOT use near newly planted shrubs, young ornamentals, and species with shallow surface roots. DO NOT plant crops other than those recommended on this label for at least nine months following treatments of this product at rates up to 4.5 litres per hectare.

DO NOT apply under meteorological conditions or from spraying equipment which could be expected to cause spray drift onto nearby susceptible plants, adjacent crops, crop lands or pastures.

When rates exceed 4.5 litres per hectare, planting's may not be possible for very long periods afterwards. Avoid deep cultivation of orchards, vineyards, asparagus, roses, berry fruit and hops which may throw untreated soil over sprayed areas as this may seriously reduce weed control.

Heavy rain following application prior to emergence may cause damage to chickpeas.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT

DO NOT apply this product within 60 m of natural or impounded lakes or dams.

DO NOT use in channels or drains where roots of desirable plants may extend. Wash sprayer thoroughly with clean water after use.

DO NOT contaminate dams, waterways or drains with chemical or used containers. This product is very highly toxic to algae and aquatic macrophytes.

RE-ENTRY PERIOD

DO NOT enter treated area without protective clothing until spray has dried.

Storage and Disposal (20L)

KEEP OUT OF REACH OF CHILDREN. Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse the containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury the empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should NOT be burnt.

(110 & BULK ONLY)

KEEP OUT OF REACH OF CHILDREN. Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of

supply for refill or storage.
DO NOT dispose of undiluted chemicals on-site.

SAFETY DIRECTIONS

Avoid contact with eyes and skin. DO NOT inhale spray mist. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. (Tel: Australia 13 11 26)

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet which can be obtained from the supplier.

CONDITIONS OF SALE

Crop Care Australasia Pty Ltd will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than the responsibility for the merchantable quality of these goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by these Statutes the liability of Crop Care Australasia Pty Ltd is limited to the replacement of the goods or (at the option of Crop Care Australasia Pty Ltd) the refund of the price paid and is conditional upon a claim being made in writing and where possible sufficient part of

In a Transport Emergency Dial 000 Police or Fire Brigade	SPECIALIST ADVICE IN EMERGENCY ONLY 1800 033 498 ALL HOURS AUSTRALIA WIDE
--	--



the goods to enable proper

Crop Care Australasia Pty Ltd
ACN 061 362 347
77 Tingira Street
PINKENBA QLD 4008

examination being
returned to Crop Care Australasia Pty Ltd within thirty
days of delivery.

Gesagard is a Registered Trade Mark of a Syngenta Group Company, Basel, Switzerland.

Roundup is a Registered Trade Mark of Monsanto.

Spinnaker is a Registered Trade Mark of American Cyanamid Corporation.

APVMA Approval No: 52582/0205