



RLP
Approved

RELEVANT LABEL PARTICULARS

READ SAFETY DIRECTIONS BEFORE OPENING OR USING



Farmoz


Simanex® 900 WG

HERBICIDE

ACTIVE CONSTITUENT: 900 g/kg SIMAZINE

GROUP C HERBICIDE

For the control of weeds in chickpeas, lupins, TT-Canola, orchards, vineyards and certain other horticultural and non-crop situations as per the Directions for Use table.



www.farmoz.com.au

CONTENTS: 10 kg, 15 kg,

DIRECTIONS FOR USE:

CROP	WEEDS CONTROLLED	STATE	RATE/ha		CRITICAL COMMENTS
			LIGHT SOIL	HEAVY SOIL	
Almonds	Amaranthus, Annual Ryegrass, Annual Thistles, Barley Grass, Bindy-eye, Capeweed,	SA only	1 kg or 60 g/100 L water	2 kg or 120 g/100 L water	Use only if trees at least 3 years old. Apply to bare moist soil immediately after cultivation and before weeds emerge.
Asparagus	Chickweed, Common Sowthistle, Corn Gromwell (Sheepweed), Creeping Oxalis, Doublegee, Fat Hen, Fumitory, Ivy-Leaf Speedwell, Mustard,	All States	1.3 kg	2.5 kg	Apply to bare moist soil after last cultivation and before spear emergence. DO NOT apply to foliage or when fruit is present. Use on established plants only.
Berry Fruits	Native Geranium, Nettles, Potato Weed, Red Shank, Shepherd's Purse, Turnip, Winter Grass, Wireweed (not Qld, Tas)	All States	1.2 kg or 60 g/100 L water	1.2 kg or 60 g/100 L water	Use only if trees are at least one year old. Apply to bare moist soil after planting. High rates may cause crop damage on sandy soils low in organic matter.
Citrus	Suppression of: Brome Grass, Soursob, Wild Oats	Qld only	2 kg	4 kg	Use only if vines are at least two years old.
Gladioli		NSW, Vic, SA, Tas, WA only	1.25 kg OR 60 g/100 L water	2.5 kg OR 120 g/100 L water	Use only if vines are at least 12 months old. In first year of use split applications are preferred.
Grapevines		All States	1.3 kg	2.5 kg	Apply to moist weed-free soil in late Winter, early Spring before hop emergence.
Hops		All States	1.1 kg	2.2 kg	Important: This use is subject to adherence to the INTEGRATED WEED MANAGEMENT STRATEGY for TT-Canola. See General Instructions: Integrated Weed Management Strategy for TT-Canola. Can be applied up to a week before sowing or post-sowing pre-emergence (ideally incorporated by harrows). For best results apply to bare moist soil, either immediately before seeding or as a pre-emergence treatment at or within 7 days of planting. Sufficient rainfall (20-30 mm) to wet the soil through the weed root zone is necessary within 2-3 weeks of application. Application should not be made to ridged or excessively cloddy soil. When applied before seeding, incorporate to a depth of 5 cm.
Canola – (Triazine Tolerant varieties only) Pre-emergence or post-sowing pre-emergence only	Capeweed, Charlock, Clover, Corn Cromwell, Doublegee, Fumitories, Geraniums, Ivy-Leaf Speedwell, London Rocket, Mustards, Turnips, Paterson's Curse, Shepherd's Purse, Silver Grass (Vulpia) Suppression of Annual Ryegrass, Barley Grass, Brome Grass, Wild Oats and Wild Radish	All States	1.1 kg	2.2 kg	

RELEVANT LABEL PARTICULARS

CROP	WEEDS CONTROLLED	STATE	RATE/ha		CRITICAL COMMENTS
			LIGHT SOIL	HEAVY SOIL	
Chickpeas	Deadnettle, Indian Hedge Mustard, Lesser Swinecress, Milk Thistle, Prickly Lettuce, Purple Goosefoot, Shepherd's Purse, Turnip Weed, Wireweed and suppression of Black Bindweed and Paradoxa Grass	NSW, Qld, SA only	800 g + 1.5 L Prometrex 500		Apply immediately post-planting. Application should not be made to ridged or excessively cloddy soil. For reliable results significant rain, 20-30 mm is necessary within 2-3 weeks of sowing.
	Capeweed, Clover, Dock, Doublegee, Mustard, Radish, Self Sown Cereals, Silver Grass, Turnip Weed and suppression of Barley Grass, Ryegrass and Wild Oats	WA only	550 g	1.1 kg	For best results apply to bare moist soil, either immediately before seeding, or as a pre-emergence treatment at or within 7 days of planting. Sufficient rainfall (20-30 mm) to wet the soil through the weed root zone is necessary within 2-3 weeks of application. Results can be variable if seasonal conditions are dry prior to sowing, and chickpeas are sown into a dry or low moisture seedbed. Application should not be made to ridged or excessively cloddy soil. When applied before seeding, incorporation by the sowing operation should not be greater than 5 cm. Use 0.55-0.775 kg on lighter soils and in the Northern Agricultural areas, and up to 1.1 kg on heavier soil types. DO NOT use on whitish or grey sands.
Faba Beans	Annual Ryegrass, Barley Grass, Brome Grass, Capeweed, Corn Gromwell (Sheepweed), Deadnettle, Fumitories, Geranium, Ivy-Leaf Speedwell, Medics, Mustards	NSW, Vic, SA, WA only	1.1 kg	1.4 kg	Apply either pre-seeding or immediately post-sowing which is preferred on light soils. Sow the crop at least 5 cm deep. Application should not be made to ridged or excessively cloddy soil. For fully reliable results significant rainfall, 20-30 mm, is necessary within 2 to 3 weeks of sowing. DO NOT use rates higher than 2 litres per hectare on soils with pH 8.0 and above as unacceptable crop damage may occur.
	Paradoxa Grass, Saffron Thistle, Soursob, Volunteer Canola, Wireweed and suppression of Wild Oats		560 g	830 g plus 800 mL of Trilogy	Apply to bare moist soil and incorporate to a depth of 5 cm just prior to sowing. Incorporation should be made within 4 hours of application. Use this mixture where Annual Ryegrass and Wild Oats are the major problems. Application should not be made to ridged or excessively cloddy soil. For fully reliable results significant rainfall, 20-30 mm, is necessary within 2 to 3 weeks of application.
Lupins	Amaranthus, Annual Ryegrass, Annual Thistles, Barley Grass, Bindy-eye, Capeweed, Chickweed, Common Sowthistle, Corn Gromwell (Sheepweed), Creeping Oxalis, Doublegee, Fat Hen, Fumitory, Ivy-Leaf Speedwell, Mustard, Native Geranium, Nettles, Potato Weed, Red Shank, Shepherd's Purse, Turnip, Winter Grass, Wireweed (not Qld, Tas) Suppression of: Brome Grass, Soursob, Wild Oats	Qld, NSW, Vic, SA, Tas only	1.3 kg	2.2 kg	Apply within 2 days of sowing. Seedbed should be moist and free of weeds, large clods and ridges. Apply higher rate for suppression of Wild Oats and Soursob. Use the lower rates (NSW, SA, Vic only) in combination with 800 mL of Trilogy.
		NSW, Vic, SA only	830 g	1.3 kg plus 1.25 L of Trilogy	Apply any time from 4 weeks before sowing until just before sowing. Apply as a tank mixture to a moist bare level soil and incorporate to a depth of 50 mm prior to sowing. Incorporation must be within 4 hours of spraying especially under warm conditions. More effective than SIMANEX 900 WG on its own in dry conditions and where grass pressures are high. Use 830 g/ha SIMANEX 900 WG on poorer light sands, 1.1 kg/ha on sandy loams and on all other soil types. Increase the rate to 1.3 kg/ha where Brome grass is the major problem.
		WA only	550 g	1.1 kg	DO NOT exceed 500 g/ha on Lupin Variety Ultra. Apply immediately before seeding or within 6 days after sowing. Seed bed should be moist and free of weeds, large clods and ridges. Use 550 g to 800 g/ha on yellow sands, 1.1 kg/ha on other soil types. Note: Not for use on white or grey sands.
			550 g	800 g plus 800 mL of Trilogy	Apply to bare moist soil just before sowing. Incorporate to a depth of 4-5 cm within 4 hours of spraying. Soil should be free of large clods or ridges. Choose SIMANEX 900 WG rate according to comments in panel above. Note: Not for use on white or grey sands.
			550 g	800 g plus recommended rate of other knockdown herbicide	Apply 1-6 days before direct drilling the crop. Use the higher rate of SIMANEX 900 WG on weeds up to two weeks old and the lower rate of SIMANEX 900 WG with a high rate of knockdown herbicide on weeds more than 2 weeks old. DO NOT exceed 500 g/ha on Lupin Variety Ultra. Note: Not for use on white or grey sands.
Lupins where weeds are present at time of sowing					
Pome Fruit, Apples, Pears		Qld only	3 kg OR 90 g/100 L water	4 kg OR 120 g/100 L water	Use only if trees are at least 2 years old. Apply to bare moist soil.
		NSW, Vic, SA, Tas, WA only	1.8 kg/ha OR 90 g/100 L water	2.5 kg/ha OR 120 g/100 L water	Use only if trees are at least 2 years old. Apply to bare moist soil.

RELEVANT LABEL PARTICULARS

CROP	WEEDS CONTROLLED	STATE	RATE/ha		CRITICAL COMMENTS
			LIGHT SOIL	HEAVY SOIL	
Roses	Amaranthus, Annual Ryegrass, Annual Thistles, Barley Grass, Bindy-eye, Capeweed, Chickweed, Common Sowthistle,	NSW, Vic, SA, Tas, WA only	1.3 kg OR 60 g/100 L water	2.5 kg OR 120 g/100L water	Apply to bare moist soil. DO NOT use on plants less than 1 year old.
Strawberries	Corn Gromwell (Sheepweed), Creeping Oxalis, Doublegee, Fat Hen, Fumitory, Ivy-Leaf Speedwell, Mustard, Native Geranium, Nettles, Potato Weed, Red shank, Shepherd's Purse, Turnip, Winter Grass, Wireweed (not Qld, Tas) Suppression of: Brome Grass, Soursob, Wild Oats	All States	1.2 kg	1.2 kg	Apply between polythene covered beds. DO NOT use on white or grey sands in WA.

CROP	WEEDS CONTROLLED	STATE	RATE/ha		CRITICAL COMMENTS
			LIGHT SOIL	HEAVY SOIL	
Sub Clover established lucerne and perennial grass pastures	Vulpia (Silvergrass), Rat's Tail Fescue, Squirrel-tail Fescue, Sand Fescue.	NSW, Vic, SA only	550 to 900 g		Apply 6 to 10 weeks after emergence of the pasture, but not before the 3rd trifoliolate leaf stage of Sub Clover. Best results are obtained from application to young, well grazed, and actively growing plants. Some damage to Sub Clover may occur especially at the higher rates. DO NOT apply to sandy soils and use the lower rates on light textured soils. Apply under moist soil conditions; rain following application enhances results. DO NOT apply to stressed plants. Insect infestations should be controlled and Sub Clover allowed to recover before the SIMANEX 900 WG is applied. DO NOT use herbicides for broad-leaved weed control within 3 weeks of application.
		NSW, Vic only	550 to 800 g plus 100 to 160 mL of Spraytop 250 SL		Apply to 6 to 10 weeks after emergence of the pasture, but not before the 3rd trifoliolate leaf stage of Sub Clover. Best results are obtained from application to young, well grazed, and actively growing plants. However, the addition of Spraytop 250 SL improves the control of well established plants. Add a non-ionic surfactant at 0.2 % v/v (200 mL per 100 litre). Under conditions of good soil moisture, control of grasses and some broad-leaved weed seedlings may occur. DO NOT apply to sandy soils and use the lower rates on light textured soils. Apply under moist conditions; rain following application enhances results. DO NOT apply to stressed plants. Insect infestations should be controlled and Sub Clover allowed to recover before the SIMANEX 900 WG plus Spraytop 250 SL is applied. DO NOT use herbicides for broad-leaved weed control within 3 weeks of application.
		SA only	400 to 550 g plus 100 to 160 mL of Spraytop 250 SL		
	Vulpia (Silvergrass), Rat's Tail Fescue	WA only	400 to 550 g		Apply within 8 weeks of emergence of Sub Clover and grasses. Under conditions of good soil moisture, control of other grasses and some broadleaved weeds may occur. Ensure that there is a good stand of Subterranean Clover present before applying. Use the lower rate on light textured soils. DO NOT use with broadleaf weed herbicides within 3 weeks of using SIMANEX 900 WG. DO NOT tank mix with other herbicides or add crop oils or wetting agents. DO NOT use on Medics, or Red or White Clover. DO NOT overlap when spraying, otherwise damage may be observed.

RELEVANT LABEL PARTICULARS

CROP	WEEDS CONTROLLED	STATE	RATE/ha		CRITICAL COMMENTS
			LIGHT SOIL	HEAVY SOIL	
Forestry including farm tree plantations <i>Pinus</i> , <i>Araucaria</i> , <i>Corymbia</i> and <i>Eucalyptus</i> tree plantings pre- and post-planting (see General Instructions for further information on tree species)	Pre-emergence control of many annual and some perennial grasses and broadleaved weeds including <i>Amaranthus</i> , Annual (Wimmera) Ryegrass, Barley Grass, Barnyard Grass, Bindii, Brome Grass, Capeweed, Chickweed, Common Sowthistle, Corn Gromwell (Iron/Sheepweed), Creeping Oxalis, Fat Hen, Fumitories, Geranium, Ivy-Leaf Speedwell, Mustards, Native Geranium, Nettles, Potato Weed, Powell's Amaranth, Red Root Amaranth, Red Shank, Shepherd's Purse, Summer Grass, Thistles - annual, soldier, variegated, Turnips, Turnip Weed, Wild Mustard, Wild Radish, Wild Turnip, Yellow Wood Sorrel Weed suppression at higher rates: Soursob, Wild oats. Weeds suppressed WA: Annual Ryegrass, Brome Grass, Dock, Double Gee, Radish, Sorrel, Soursob, Wild Turnip, Wireweed	All States	1.6-6.7 kg/ha		SIMANEX 900 WG may be applied prior to or after planting. For most situations use 5.0 to 5.5 kg/ha. On gravelly coarse textured soils use 2.8 to 4.0 kg/ha. In WA and SA in medium rainfall zones (400-600 mm) use 3.0 to 4.0 kg/ha and for sandy low organic soils use 1.6 to 3.0 kg/ha. On sandy soils allow two weeks and at least 50 mm of rain between spraying and planting. Use a maximum rate of 1.6 kg/ha for overspraying eucalyptus plantations on sandy, low organic matter soils in WA and SA. For short term weed control or when applied with other pre-emergent herbicides use half rates according to the situation. Apply to bare ground. Best results will be achieved if applied to moist, finely tilled soil. If weeds are present add a knockdown or other partner herbicide as a tank mix. If spraying after planting check tree tolerance to knockdown or other partner herbicide before using. Application in a band 1.5 to 2.0 m wide, over the planting row is recommended although broadcast application can also be used. Rates above refer to sprayed hectares and not field hectare. Some tree damage may occur, especially with post-plant application. Recovery is normally rapid. Directed application is preferred for post-planting application especially when the higher rates are used.
Non-crop uses, Commercial Industrial, Rights of Way, Public Utility areas, Road shoulders, Drains, Headlands, Driveways, Railway tracks, Aerodromes, Gutters, Footpaths	As above and including Barnyard Grass, Summer Grass	All States	5 kg OR 500 g/100 L water	5 kg OR 500 g/100L water	Provides season control of annual weed germinations. Apply to bare moist ground.
			10 kg OR 1 kg/100 L water	24 kg OR 2.4 kg/100 L water	Provides long residual control in Winter rainfall areas. Apply to bare moist ground.
			24 kg OR 2.4 kg/100 L water	48 kg OR 4.8 kg/100 L water	Provides long residual control in Summer rainfall areas. Apply to bare moist ground. Read the remarks under Crop Safety.
Dams, Tanks, Troughs	Filamentous Blue-Green Algae	WA only	2 g per 100 litres of water		Mix in a convenient amount of water and apply when algae development first noticed.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

HARVEST:

ALL CROPS – NOT REQUIRED WHEN USED AS DIRECTED.

GRAZING:

CANOLA: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 15 WEEKS AFTER APPLICATION.

SUB CLOVER: DO NOT GRAZE FOR 14 DAYS AFTER APPLICATION.

DO NOT CUT FOR STOCK FOOD FOR 21 DAYS AFTER APPLICATION.

CHICKPEAS: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 9 WEEKS AFTER APPLICATION.

FABA BEANS: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 8 WEEKS AFTER APPLICATION.

GENERAL INSTRUCTIONS

FARMOZ SIMANEX® 900 WG is a pre-emergent herbicide which is most effective when applied to bare, moist soil. Because it acts mainly through the roots of weeds, best results are obtained when rain or irrigation water move the herbicide down to the root zone within 2 weeks of application. The duration and effectiveness of control depends on the amount applied, the type of soil, intensity of leaching and the species of weeds present.

RESISTANT WEEDS WARNING

FARMOZ SIMANEX® 900 WG Herbicide is **GROUP C HERBICIDE** a member of the Triazine group of herbicides. SIMANEX 900 WG has the inhibitor photosynthesis mode of action. For weed resistance management SIMANEX 900 WG is a Group C herbicide. Some naturally-occurring weed biotypes resistant to SIMANEX 900 WG and other Group C herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by SIMANEX 900 WG or other Group C herbicides. However resistance should not be assumed without first viewing the method of application, the coverage and the timing of application. Resistance to SIMANEX 900 WG cannot be predicted and may occur at any time. It is not due to a fault in the product and FARMOZ Pty. Ltd. accepts no liability for any losses that may result from the failure of SIMANEX 900 WG to control resistant weeds.

INTEGRATED WEED MANAGEMENT STRATEGY FOR TT-CANOLA

An Integrated Weed Management Strategy for TT-Canola (the Strategy) has been developed and funded by the Canola Association of Australia in cooperation with certain companies including FARMOZ Pty. Ltd. The Strategy outlines recommendations, measures and options for weed management, including management of herbicide resistance in weed populations. The Strategy is available from a FARMOZ representative and the Canola Association of Australia. A program has been developed that outlines sound agronomic practices and integrated weed management programs designed to optimise the performance of TT-Canola. It is advised that consultation on IWM be undertaken with an accredited agronomist prior to use of SIMANEX 900 WG on TT-Canola.

TO MINIMISE HERBICIDE RESISTANCE

- Avoid dry sowing in heavily weed infested paddocks. Wait for a weed germination after the opening rains in weedy paddocks. Use a pre-plant knockdown or cultivation. No weeds should be allowed to survive at this stage.
- Adapt the weed control program to the anticipated weed spectrum and pressure:
 - Broadleaf Weeds and Ryegrass:** Use SIMANEX 900 WG or Farnozine 900 WDG plus trifluralin pre-emergence. A follow-up with a Group A herbicide (if ryegrass is susceptible) or Farnozine 900 WDG may be necessary.
 - Broadleaf Weeds only:** Use Farnozine 900 WDG post-emergence.
- DO NOT use Farnozine 900 or SIMANEX 900 WG if the area to be treated had a triazine herbicide applied to it last season.
- Watch for escapes, especially in paddocks with a long history of Group C herbicide use.
- DO NOT use Group C herbicides in consecutive years.

TO AVOID TRIAZINE CARRY-OVER

On acid soils (pH less than 6.5) – The maximum rate of Farnozine 900 WDG or SIMANEX 900 WG or a combination of the two products to be applied to the crop during the growing season is 2.2 kg/ha.

On alkaline soils (pH greater than 6.5) – The maximum rate of Farnozine 900 or SIMANEX 900 WG or a combination of the two products to be applied to the crop during the growing season is 1.1 kg/ha.

Post-emergence use – It is recommended that Farnozine 900 only be used, and at rates of 1.1 kg/ha or less, on both acid or alkaline soils.

RESISTANT WEEDS REPORTING

Growers should collect plant or seed samples where weeds that are normally susceptible to atrazine and simazine may be resistant, get them tested and seek professional advice.

MIXING

Half fill tank with clean water. Commence vigorous agitation. Add granules steadily. DO NOT dump. Ensure complete dispersion before filling to volume with water. Continue agitation during spraying. DO NOT leave spray solution in tank overnight.

APPLICATION

Use a boom sprayer or knapsack sprayer for small areas. 50-200 L of mixture per hectare should be applied with the boom, and a minimum equivalent to 500 L per hectare through the knapsack sprayer. Clean spray equipment immediately after use using several flushings of water with detergent.

TT-CANOLA: APPLICATION

DO NOT apply to TT-Canola by aircraft. Apply only with a low boom sprayer with a 60 m buffer zone downwind of treated fields to natural or impounded lakes or dams, and a 20 m buffer zone for any well, sink hole, intermittent or perennial stream. Apply only to areas where run-off is unlikely to occur or where run-off may be captured by farm earthworks.

COMPATIBILITY

This product may be mixed with Amitrole, Atrazine (Farnozine), Diuron, Glyphosate (Wipe-Out®, Wipe-Out® Plus), Diquat, Paraquat (Spraytop® 250 SL) and Paraquat-diquat mixtures, Prometryn (Prometrex® 500) and Trifluralin (Trilogy®). Always add the granules to the tank first as per the mixing instructions and ensure the product is completely dispersed before adding any other product.

Because products may vary from time to time, a small scale compatibility test should be carried out before mixing in the spray tank. For residual control of annual weeds in non-crop areas, pome and citrus orchards and vineyards only, mixtures of this product and Wipe-Out 360 herbicide may be used. Half fill the spray tank with clean water before adding this product at the recommended rate, and mix thoroughly. Then add Wipe-Out 360 at 3 L/ha or 700 mL/100 L (or Wipe-Out Plus at 2.1 L/ha or 500 mL/ha) and the remaining water. Maintain adequate agitation and use the mixture promptly. DO NOT use lower rates of Wipe-Out 360 (or Wipe-Out Plus) where control of perennial weeds is desired.

PRECAUTION

Re-Entry Period: DO NOT enter treated areas without protective clothing until spray has dried.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT spray foliage of desirable plants, except lupins (WA only). DO NOT use under peach, apricot, plum trees, ornamental prunus or near newly planted shrubs. DO NOT plant crops other than those recommended for at least nine (9) months following treatments at rates up to 2.5 kg/ha. At higher rates planting may not be possible for much longer periods afterwards. DO NOT use on sandy or porous soils near desirable plants or trees or where roots of such may extend or are near the surface. Crops such as red beet, lettuce, cabbage, tomato, cucumber and carrots are particularly sensitive to Simazine residues. DO NOT exceed highest rate recommended for any particular crop. DO NOT use on white or grey sands (WA only). DO NOT use in channels or drains where the roots of desirable plants may extend.

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

Araucaria, Corymbia, Eucalyptus and Pinus Plantations: Field trials and use experience has indicated that the following species are tolerant. Test tolerance before treating other species.

Araucaria: All commercial species.

Corymbia: Maculata.

Eucalyptus: Agglomerate, calophylla, camaldulensis, cladocalx, cloeziana, dunnii, globulus, grandis, laevopinea, nitens, pilularis, pseudoglobulus, regnans, rudis, saligna, wandoo.

Pinus: Caribaea and hybrids, elliottii, pinaster, radiata.

SOIL MOISTURE

If sufficient rain to thoroughly wet the soil through the weed root zone has not fallen within 14 days of application, water in with the equivalent of 13 mm of rain. If applied under dry conditions and irrigation is not available, mechanical incorporation to a depth of not more than 4 cm within 7 days of sowing may assist results.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate dams, rivers or streams with pesticide or used containers. DO NOT use the container for any other purpose. This product is very highly toxic to algae and aquatic macrophytes.

RELEVANT LABEL PARTICULARS

STORAGE AND DISPOSAL

KEEP OUT OF REACH OF CHILDREN. Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Shake empty bag into spray tank. DO NOT dispose of undiluted chemicals on site. Break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

SAFETY DIRECTIONS

Avoid contact with eyes and skin. DO NOT inhale dust or spray mist. Wash hands after use.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 131126.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for FARMOZ SIMANEX 900 WG is available from FARMOZ on request. Call Customer Service on (02) 9431 7800.

CONDITIONS OF SALE: The use of FARMOZ SIMANEX 900 WG Herbicide being beyond the control of the manufacturer, no warranty expressed or implied is given by FARMOZ Pty. Ltd., regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and FARMOZ Pty. Ltd. accepts no responsibility for any consequence whatsoever from the use of this product.

® Registered trademarks of a Makhteshim-Agan Company

NOT A DANGEROUS GOOD ACCORDING TO THE AUSTRALIAN DANGEROUS GOODS (ADG) CODE.

Farmoz Pty. Ltd. ABN 55 050 328 973

Suite 1, Level 4, Building B,
207 Pacific Highway
St Leonards NSW 2065 Australia
Tel: (02) 9431 7800 Fax: (02) 9431 7700

APVMA Approval No.: 62917/57494

Batch No.

Date of Manufacture