

CAUTION

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

EJECT HERBICIDE

ACTIVE CONSTITUENT: 750g/kg SULFOMETURON METHYL

GROUP	B	HERBICIDE
--------------	----------	------------------

For the control of certain annual and perennial grasses and broadleaf weeds in commercial and industrial areas: around agricultural buildings and rights of way as per directions for use.

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE

Net Contents : 200g, 500g, 1kg, 1.5kg

Sipcam Pacific Australia Pty Limited
A C N 073 176 888
Suite 11
23 – 31 Gheringhap Street
Geelong Vic 3220

STORAGE AND DISPOSAL

During storage keep from contact with fertilisers, other pesticides and seeds.

Store in the closed original container in a well-ventilated area, as cool as possible but out of direct sunlight.

Triple or (preferably) pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Avoid contact with eyes and skin. DO NOT inhale dust. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (TEL: 13 11 26).

Note

This product has the potential to be carried in water from the treated area to damage non-target plants. Refer to "Protection of Crops, Native and other Non-Target Plants" section of this label/leaflet.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet which can be obtained from the supplier.

NOTICE TO BUYER

This product is not to be used for any purpose or in any way contrary to label instructions. The buyer by use or application of this product acknowledges so doing upon the terms and conditions agreed between Sipcam Pacific Australia Pty Limited and the buyer but not limited to the following.

To the extent permitted by law all conditions and other warranties or other rights of action which buyer or any other user may have against Sipcam Pacific Australia Pty Limited (Sipcam) or Seller are hereby excluded. Sipcam hereby gives notice to buyer and other users that it will not accept responsibility for any indirect or consequential loss arising from reliance on product information or advice provided by Sipcam or on its behalf unless it can be established that such information or advice was provided negligently and that the product has been used strictly as directed. Notwithstanding the generality of this notice Sipcam will not be held liable for any loss, injury or damage indirect or consequential arising from the sale, supply, use or application of this product. Sipcam's liability shall in all circumstances be limited to replacement of the product or a refund of the purchase price paid therefor.

**For specialist advice in an emergency dial
1 800 033 111
24 hours Australia wide.**

Batch No:

Date of Manufacture:

NRA Approval No:

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

EJECT HERBICIDE

ACTIVE CONSTITUENT: 750g/kg SULFOMETURON METHYL

GROUP	B	HERBICIDE
-------	---	-----------

For the control of certain annual and perennial grasses and broadleaf weeds in commercial and industrial areas: around agricultural buildings and rights of way as per directions for use.

IMPORTANT: READ THIS LEAFLET BEFORE USE

Sipcam Pacific Australia Pty Limited
A C N 073 176 888
Suite 11, 23 – 31 Gheringhap Street
Geelong VIC 3220

DIRECTIONS FOR USE**RESTRAINTS**

DO NOT Apply EJECT to drainage ditches and channels

DO NOT apply EJECT during periods of intense rainfall, or to soils saturated with water as off-target movement may occur

DO NOT store tank mixes of EJECT with other materials for more than 24 hours as product breakdown may occur

DO NOT store a suspension of EJECT for more than 2 days as it may degrade, particularly in acid solutions.

FOR ALL STATES

SITUATION	WEEDS	Rate			CRITICAL COMMENTS
		Boom g/ha	Hand gun g/100L	Knap sack g/15L	
Commercial & Industrial areas, around Agricultural buildings, Rights-of-Way including road sides, around guide posts, railways, power lines and telephone lines	<u>Easy-to-Kill Annual Weeds</u> Grasses - Annual Ryegrass, Barnyard grass, Summer grass/Crab grass, Wild Oats, and Winter grass. Broadleaves – Blue Billygoat Weed, Capeweed, Corkscrew/Erodium, Paterson's Curse/Salvation Jane, Subterranean Clover and Turnip Weed <u>Easy-to-Kill Perennial Weeds</u> Grasses – Prairie grass Broadleaves - Dandelion, Dock, White Clover	200 or 400	20 or 40	3 or 6	<p>For roadside applications confine treatment to the area between the edge of the bitumen and the roadside shoulder.</p> <p>For weeds advanced beyond the 4 leaf stage or when more rapid knockdown is required, tank mix with Glyphosate 360 Herbicide in accordance with its label recommendations. Always add a surfactant - except when tank mixing with Glyphosate 360 Herbicide. Use the higher rate plus Glyphosate 360 Herbicide (in accordance with its label recommendations), for dense weed infestations and where weeds are greater than 15cm in height or diameter.</p> <p>Use the higher rate for longer term residual control.</p> <p>For handgun or knapsack application, apply the spray evenly to the required area. Spray to wet the foliage of the weeds present, but not excessively so as to cause run-off. Over application may reduce knockdown results and significantly increase period of residual control.</p>

SITUATION	WEEDS	Rate			CRITICAL COMMENTS
		Boom g/ha	Hand gun g/100L	Knap sack g/15L	
Commercial & Industrial areas, around Agricultural buildings, Rights-of-Way including road sides, around guide posts, railways, power lines and telephone lines	<u>Hard-to-Kill Annual Weeds</u> Grasses - Rats Tail Fescue, Squirrel Tail Fescue. Broadleaf – Black Thistle/Spear Thistle	200 or 400	20 or 40	3 or 6	For roadside applications confine treatment to the area between the edge of the bitumen and the roadside shoulder. Always tank mix with Glyphosate 360 Herbicide in accordance with its label directions - a surfactant is not required. Use the higher rate plus Glyphosate 360 Herbicide (in accordance with its label recommendations), for dense weed infestation and where weeds are greater than 15cm in height or diameter. Use the higher rate for longer term residual control. For handgun or knapsack application apply the spray evenly to the required area. Spray to wet the foliage of the weeds present, but not excessively so as to cause run-off. Over application may reduce knockdown results and significantly increase the period of residual control.
	<u>Hard-to-Kill Perennial Weeds</u> Grasses - African Lovegrass, Columbus grass, Couch, Elastic grass, Guinea grass, Johnson grass, Kikuyu, Paspalum and Phalaris	400 or 800	40 or 80	6 or 12	

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED

GENERAL INSTRUCTIONS

Eject is a soil residual herbicide with some knockdown action. It controls susceptible weeds through both post-emergence and pre-emergence (residual) action.

Eject may be applied pre-emergence or post-emergence to the weeds; best results are obtained if application is made before or during the early stages of weed growth.

Adequate soil moisture is required for Eject to be activated so it can provide optimum weed control. Under limited rainfall conditions, Eject may not provide satisfactory control of hard-to-kill annuals and perennial weeds.

The higher use rates give longer term residual control.

The degree of control and duration of effect will vary with the rate of application, soil texture, organic matter content, soil pH, rainfall and other environmental conditions.

As soil pH increases the rate of Eject breakdown decreases so that longest term residual control is obtained on alkaline soils. As soil moisture increases the rate of Eject breakdown decreases, so that longest term residual control is obtained when rain-fall is low.

As soil organic matter content increases the rate of Eject breakdown increases, so that longest term residual control is obtained in low organic matter soils.

RESISTANT WEEDS WARNING

Group B Herbicide

Eject is a member of the sulfonylurea group of herbicides. Eject has the acetolactate synthase (ALS) inhibitor mode of action. For weed resistance management Eject is a Group B herbicide.

Some naturally-occurring weed biotypes resistant to Eject and other ALS inhibitor herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Eject or other ALS inhibitor herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Sipcam Pacific Australia Pty Ltd accepts no liability for any losses that may result from the failure of Eject to control resistant weeds.

Strategies to minimise the risk of herbicide resistance are available. Consult your farm chemical supplier, consultants local Department of Agriculture of Primary Industries, or Sipcam representative.

SPRAY PREPARATION

Eject is a dry flowable formulation to be mixed with water and applied as a spray. Partially fill the spray tank with water. Using the graduated measuring cylinder provided, measure the amount of Eject required for the area to be sprayed. DO NOT shake the Eject down as the measuring cylinder is graduated for Eject in an uncompacted state. Add the correct amount of Eject to the spray tank with agitation system engaged. Top up the correct volume with water. THE MATERIAL MUST BE KEPT IN SUSPENSION AT ALL TIMES BY CONTINUOUS AGITATION.

(NB: The measuring cylinder provided is graduated in grams of Eject ONLY. DO NOT use for measuring other materials.)

In tank mixes, Eject must be in suspension before adding the companion herbicide or surfactant.

Where prepared spray solutions have been allowed to stand, thoroughly reagitate before using.

USE OF SURFACTANT

Always add a non-ionic surfactant at the rate of 0.25% v/v (eg Surfactant Wetting Agent Maxiwet 1000 at 250 mL/100 L) when Eject is used alone. When Eject is tank mixed with Glyphosate 360 Herbicide, no surfactant addition is necessary

SPRAY EQUIPMENT

DC NOT apply by aerial application equipment or mist spray application equipment

BOOM SPRAY

Eject is recommended for application by ground spraying equipment only. Use a boom spray (or off-centre nozzle if necessary), properly calibrated to a constant speed and rate of delivery, to ensure thorough coverage and a uniform spray pattern and rate of application over the area to be treated. Apply a minimum of 50 L prepared spray/ha.

HANDGUN/KNAPSACK

Wet the foliage of the weeds present but not excessively so as to cause run-off. Handgun - Use 1000 - 2000 Litres of spray mix per hectare. Eject residues in the spray tank can damage sensitive plants. Following a Eject application the spray tank should not be used for other than non-crop applications.

COMPATIBILITY

Eject is compatible with Bromacil 800, KROVAR¹, Glyphosate 360 formulations and VELPAR¹ L.

SPRAYER CLEANUP

Eject residues in the spray tank can damage sensitive plants. It is recommended that a sprayer be dedicated to the use of Eject. Where equipment is also used for crop spraying, it is particularly important that all traces of Eject be removed from mixing and spraying equipment immediately after spraying. The recommended sprayer cleanup is as follows:

1. Drain tank, then flush tank, boom, filters and hoses with clean water for a minimum of 10 minutes.
2. Fill the tank with clean water and add 300 mL chlorine bleach (containing 4% chlorine) per 100 L water. Flush through boom, filters and hoses, then allow to sit for 15 minutes with agitation engaged, then drain.
3. Repeat Step 2.
4. Nozzles and screens should be removed and cleaned separately. To remove traces of chlorine bleach, rinse the tank thoroughly with clean water and flush through boom, filters and hoses.

CAUTION:

DO NOT use chlorine bleach with ammonia.

If mixing with glyphosate, DO NOT mix or store or apply this product in galvanised steel, unlined steel containers or spray tanks, since a highly flammable gas mixture may be formed. So use only spray tanks and equipment made of plastic or plastic lined, fibreglass, rubber, aluminium, brass, copper, or stainless steel.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply to drainage ditches (including roadside ditches and table drains) and channels, as some of the chemicals may wash away in drainage water and be deposited in areas where it may cause damage to desirable trees and other plants.

DO NOT apply under weather conditions or from spraying equipment that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures, or onto agricultural land scheduled to be planted to crops or pastures, as injury to the crop or pasture may occur.

PROTECTION OF LIVESTOCK

DO NOT graze treated areas or cut for stock food.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT apply or drain or flush equipment on or near desirable trees or other plants, or on areas where their roots extend, or in locations where the chemical may be washed or moved into contact with their roots.

DO NOT contaminate any body of water by spraying, cleaning of equipment, disposal of waste, including chemical or used containers, or run-off from treated areas. If accidental contamination does occur, the water must not be used for irrigation.

STORAGE AND DISPOSAL

During storage keep from contact with fertilisers, other pesticides and seeds.

Store in the closed original container in a well-ventilated area, as cool as possible but out of direct sunlight.

Triple or (preferably) pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Avoid contact with eyes and skin. DO NOT inhale dust. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Tel: 13 11 26).

Note

This product has the potential to be carried in water from the treated area to damage non-target plants. Refer to "Protection of Crops, Native and Other Non-Target Plants" Section of this label/leaflet.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet which can be obtained from the supplier.

NOTICE TO BUYER

This product is not to be used for any purpose or in any way contrary to label instructions. The buyer by use or application of this product acknowledges so doing upon the terms and conditions agreed between Sipcam Pacific Australia Pty Limited and the buyer but not limited to the following.

To the extent permitted by law all conditions and other warranties or other rights of action which buyer or any other user may have against Sipcam Pacific Australia Pty Limited (Sipcam) or Seller are hereby excluded. Sipcam hereby gives notice to buyer and other users that it will not accept responsibility for any indirect or consequential loss arising from reliance on product information or advice provided by Sipcam or on its behalf unless it can be established that such information or advice was provided negligently and that the product has been used strictly as directed. Notwithstanding the generality of this notice Sipcam will not be held liable for any loss, injury or damage indirect or consequential arising from the sale, supply, use or application of this product. Sipcam's liability shall in all circumstances be limited to replacement of the product or a refund of the purchase price paid therefor.

**For specialist advice in an emergency
dial
1 800 033 111
24 hours Australia wide.**

Batch No:

Date of Manufacture:

NRA Approval No:

¹ Trademark E I du Pont de Nemours & Co Inc