

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

United Farmers

Tebuconazole 430 SC Fungicide

ACTIVE CONSTITUENT: 430g/L Tebuconazole



For the control of various diseases of bananas, peanuts, cereal crops, vegetables and other crops as specified in the 'Directions for Use' Table.

***NOTE - This information is adapted from the United Farmers Tebuconazole 430 EC label.
Please consult the label on the pack before use***



United Farmers Co-operative Company Limited

2 BIRKSGATE ROAD ROUS HEAD
NORTH FREMANTLE, W.A. 6160
Phone 08 9430 2222 Fax 08 9430 2233

CONTENTS: *L (1 Litre, 5 Litres, 20 Litres)
Batch NO.:

APVMA Approval No.: 58028

DIRECTIONS FOR USE

Crop	Disease	Application	Rate	WHP (days)	Critical Comments
Bananas (Qld, NSW, WA, NT only)	Leaf spot (yellow sigatoka) Leaf speckle Black sigatoka	Ground Application Apply by misting machine or airblast sprayer in a convenient volume of water Aerial Application Apply at least 20 L of spray mixture per hectare	230 mL/ha Add 3-6 L water miscible oil/ha	H 1 day	Maintain good deleafing practices to reduce disease inoculum. Very old leaves and leaves with advanced lesions should be removed, or infected parts of the leaf removed, prior to the application of United Farmers Tebuconazole 430 SC Fungicide. Tropical areas (e.g. North Qld, NT, Ord River): Apply a regular schedule of protectant sprays. When conditions favour disease, apply a minimum of 2 and a maximum of 3 consecutive United Farmers Tebuconazole 430 SC Fungicide sprays at 14-day intervals. Do not apply more than 6 United Farmers Tebuconazole 430 SC Fungicide sprays in any 12-month period. Do not apply any United Farmers Tebuconazole 430 SC Fungicide sprays in the months of July, August and September. Sub-tropical areas (e.g. South Qld, NSW): Commence spraying with United Farmers Tebuconazole 430 SC Fungicide at the onset of warm and humid/wet weather, normally December. Repeat at 21 to 28 day intervals using a minimum of 2 consecutive United Farmers Tebuconazole 430 SC Fungicide sprays. Do not apply more than 5 United Farmers Tebuconazole 430 SC Fungicide sprays in any 12-month period. United Farmers Tebuconazole 430 SC Fungicide is approved for use in banana plantations inter-planted with avocados.
Peanuts (South QLD and NSW only)	Early leaf spot Late leaf spot Rust Net blotch	Ground Application Apply in at least 100 L of water/ha Aerial Application Apply at least 30 L of spray mixture per hectare Low disease High disease OR Wet weather High disease AND Wet weather	175 mL/ha +Agridex 1 L/ha 290 mL/ha +Agridex 1 L/ha 440 mL/ha +Agridex 1 L/ha	H 3 weeks G 3 weeks	Regularly check high-risk areas in the crop for disease (eg lower leaves, shaded plants). When leaf spot or net blotch or rust can easily be found then either; - spray <u>immediately</u> after the last disease conducive weather (e.g. rain or heavy dews) OR -spray before the next disease conducive weather. Repeat after 14 days if conditions remain favourable to disease development. If not, repeat before or just after the next disease conducive weather. For resistance strategy, see General Instructions.

DIRECTIONS FOR USE

Crop	Disease	Application	Rate	WHP (days)	Critical Comments
Peanuts (North QLD, WA, NT only)	Early leaf spot Late leaf spot Rust	Ground Application Apply in at least 100 L of water/ha	230 mL/ha +Agridex 1 L/ha	H 3 weeks G 3 weeks	Begin spraying at 3-4 weeks after planting. If band spraying, apply half the appropriate rate as a 45 cm band directly over the row. Sprays should not be banded after 6 weeks from planting. Repeat applications at 14-day intervals. If prolonged wet weather or heavy rains occur, shorten spray interval to 10-12 days.
		Aerial Application Apply at least 30 L of spray mixture per hectare			
		Low disease			
	Moderate disease	290 mL/ha +Agridex 1 L/ha			
	Severe disease	440 mL/ha +Agridex 1 L/ha			
Net blotch	Low disease	290 mL/ha +Agridex 1 L/ha	Apply at 14-day intervals. If prolonged cool moist weather occurs, shorten spray interval to 10-12 days. For resistance strategy, see General Instructions.		
	Moderate – severe disease	440 mL/ha +Agridex 1 L/ha			
Green beans	Rust	Can be applied by aircraft or ground rig	350 mL/ha +Agridex 1 L/ha	H 3 days G 3 days	Spray when rust infection begins or at budding, whichever is the earlier. Repeat application 10-14 days later. A third application may be necessary when infection occurs early or disease pressure is high.
Onions (TAS only)	White root rot	Before sowing, apply United Farmers Tebuconazole 430 SC Fungicide onto lime super. Ensure good coverage of all lime super particles.	1.45 mL/100m of row mixed with 145-218g lime super/100m of row	-	Apply United Farmers Tebuconazole 430 SC Fungicide treated lime super when sowing onion seed. Seed and lime super can either be mixed in the same box on the drill or placed in different boxes and sown down the same tube. Apply in a bandwidth of 2 cm. Ensure that the correct rate of United Farmers Tebuconazole 430 SC Fungicide is used otherwise some delay in emergence and reduced stands of seedlings may occur.

DIRECTIONS FOR USE

Crop	Disease	Application	Rate	WHP (days)	Critical Comments
Papaw	Black spot	Ensure thorough coverage of leaves and fruit.	290 mL/ha	H 3 days	Ensure infected plant material is regularly removed and destroyed to reduce inoculum levels. Spray equipment must be properly calibrated to apply the correct amount of United Farmers Tebuconazole 430 SC Fungicide. Apply United Farmers Tebuconazole 430 SC Fungicide at 14-day intervals. Alternate United Farmers Tebuconazole 430 SC Fungicide with sprays of a protectant fungicide (eg Mancozeb). Do not apply more than 6 sprays of United Farmers Tebuconazole 430 SC Fungicide (or any DMI fungicide) on any block in any 12-month period.
Peas	Powdery Mildew	Ground Application Apply in at least 50 L of water/ha Aerial Application Apply in at least 10 L of water/ha	145 mL/ha	H 3 Days G 3 Days	Apply at flowering or at first sign of disease, whichever occurs first. A second spray 14 days later may be necessary under some conditions.
Pyrethrum	Sclerotinia sclerotiorum		350 mL/ha	-	Apply twice, in rotation with other control measures, at 7-10 day intervals. Commence at 1 to 2% flowering. Use under direction of pyrethrum advisers.
Ryegrass and fescue seed crops	Leaf rust Stem rust	Apply in at least 100 L of water/ha.	290 mL/ha	-	Monitor crops closely and spray at the first signs of disease. Continuing disease pressure or re-infection may require a further application 3-4 weeks later. Ensure thorough coverage and use higher water volumes in dense or advanced crops.
Sugarcane (Variety Q124 Only)	Orange Rust	Ground Application Use droppers and directed sprays and sufficient water volume to ensure thorough coverage. Aerial Application Apply in a minimum sprat volume of 20L per hectare.	290 mL/ha + Agridex 1L/ha.	H 4 Weeks G 4 Weeks	Even low levels of orange rust suppress yields so it is important to apply United Farmers Tebuconazole 430 SC Fungicide early in the development of the disease epidemic. Begin monitoring disease levels early. Check crops at least weekly when climatic conditions favour for the development of disease. Apply as a foliar spray when disease begins to escalate rapidly. Repeat application after 14 days if conditions remain favourable to orange rust spore germination. Do not apply more than 2 United Farmers Tebuconazole 430 SC Fungicide spray per season. DO NOT apply if heavy rains or storms that are likely to cause surface runoff are forecast with greater than 50% probability within 24 hours (48 hours if possible) of application.

DIRECTIONS FOR USE

Crop	Disease	Application	Rate	WHP (days)	Critical Comments
Wheat	Leaf rust Stripe rust Septoria nodorum blotch Yellow leaf spot	Ground Application Apply in at least 50 L of water/ha Aerial Application Apply in at least 10 L of water/ha	145 mL/ha or 290 mL/ha	H 5 weeks G 14 Days	Use higher rate when longer disease control is required. Stripe rust: See spray timings under General Instructions. Other diseases: Apply from full flag leaf emergence to early head emergence. The addition of mineral crop oil (e.g. D-C Trate or equivalent) at 1%, may improve performance of United Farmers Tebuconazole 430 SC Fungicide on wheat, oats and barley.
	Septoria tritici blotch		290 mL/ha		
Oats	Crown rust	145 or 290 mL/ha			
Barley	Scald	145 mL/ha		Apply at late tillering to early jointing.	
	Powdery mildew	145 or 290 mL/ha		Apply when 5% of the leaf area is infected. Use higher rate when longer disease control is required.	
Wheat Oats	Stern rust			Apply if more than 5% of stems become infected between full flag leaf emergence to late flowering. Where stern rust is the major disease, yield responses are usually optimized by delaying application until full head emergence, and using the higher rate. In severe cases, if a majority of stems are infected prior to full head emergence, apply at 145 mL/ha as soon as possible and if necessary, repeat after 3 weeks when heads are fully emerged.	

*H = Harvest, G = Grazing

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

- BANANAS, AVOCADOS:** DO NOT HARVEST FOR 1 DAY AFTER APPLICATION.
- BEANS, PEAS:** DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION AND DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 3 DAYS AFTER APPLICATION.
- CEREALS:** DO NOT HARVEST FOR 5 WEEKS AFTER APPLICATION AND DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 14 DAYS AFTER APPLICATION.
- ONIONS:** NOT REQUIRED WHEN USED AS DIRECTED
- PAPAW:** DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION
- PEANUTS:** DO NOT HARVEST FOR 21 DAYS AFTER APPLICATION AND DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 21 DAYS AFTER APPLICATION
- SUGARCANE:** DO NOT HARVEST FOR 4 WEEKS AFTER APPLICATION. DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 4 WEEKS AFTER APPLICATION.

GENERAL INSTRUCTIONS

Special Warning – bananas

There are certain conditions when the surface of recently emerged fruit is particularly prone to marking damage from spray applications. In circumstances where application will be made to very rapidly growing fruit in hot conditions with strong direct light, it is recommended that all emerged bunches be bagged prior to spraying to minimise risk of fruit marking. **Never include adjuvants** other than water miscible oils with United Farmers Tebuconazole 430 SC Fungicide spays. The inclusion of wetting agents is known to cause phototoxicity to young fruit.

Foliar diseases on cereal crops

Do not apply to cereal crops more than once in a season. Treatment will give approximately three weeks disease suppression. Economic responses may not be gained by spraying crops past flowering stage. The effects of fungicide application will not be clearly seen for 7-10 days after application. Yield potential: crops with potential yield under 2t/ha are unlikely to give economic responses to a fungicide spray except under conditions of very severe disease. Economic responses are most likely with crops with potential yield of over 3t/ha.

Spray timings for stripe rust control

Obtain advisory literature from Department of Agriculture for classification of resistant and susceptible varieties. In South Australia consult Plant Protection Note PPN 21. In other states the following spray program is suggested:

Seedling infections: When approximately 20 out of 100 leaves show first signs of infection during tillering to jointing, apply a spray within one week.

Adult infections – susceptible varieties: When approximately 10 out of 100 leaves show first sign of infection, apply a spray within one week.

Adult infections – moderately susceptible varieties: When approximately 15 to 20 leaves out of 100 leaves show first sign of infection, apply a spray within one week. Do not delay.

Adult infections – moderately resistant and resistant varieties: Monitor carefully. If rust appears and spreads, spray within one week.

Fungicide Resistance Warning

GROUP	C	FUNGICIDE
--------------	----------	------------------

United Farmers Tebuconazole 430 SC

Fungicide 430 Fungicide is a member of the DMI group of fungicides. For fungicide resistance management the product is a Group C fungicide. Some naturally occurring individual fungi resistant to United Farmers Tebuconazole 430 SC and other Group C fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by United Farmers Tebuconazole 430 SC Fungicide or other Group C fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, United Farmers Cooperative accepts no liability for any losses that may result from the failure of United Farmers Tebuconazole 430 SC Fungicide to control resistant fungi.

Resistance management recommendation – peanuts

Apply no more than 3 consecutive sprays of DMI fungicide (eg United Farmers Tebuconazole 430 SC Fungicide) before switching to a non-DMI fungicide. Apply no more than 5 DMI sprays per season.

Export of treated produce:

Growers should note that MRLs or import tolerances do not exist in all markets for edible produce treated with United Farmers Tebuconazole 430 SC Fungicide. If you are growing edible produce for export, please check with United Farmers Cooperative for the latest information on MRLs and import tolerances before using this product.

Mixing:

Prior to pouring, shake container vigorously, then add the required quantity of United Farmers Tebuconazole 430 SC Fungicide to water in the spray vat while stirring or with agitators in motion. Add the required amount of Agridex® (peanuts, beans) or water miscible oil (bananas) and mix thoroughly.

Application

Aircraft should fly as low as possible under the prevailing conditions to minimise drift.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate ponds, waterways and drains with this product or containers.

A spray drift minimisation strategy should be employed at all times when aerially applying sprays. The strategy envisaged is exemplified by the cotton industry's Best Management Practices Manual.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

1L Containers Only:

Rinse containers before disposal. Add rinsings to the spray tank. Do not dispose of undiluted chemicals on site. Dispose of at a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots. Empty containers and product should not be burnt.

5, 10, 20L Containers Only:

Triple or preferable pressure rinse containers before disposal. Add rinsings to the spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling break, crush, or puncture and bury containers at a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container, preparing the spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves. After use, and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (Tel: 13 11 26).

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet which is available from the supplier.

CONDITIONS OF SALE

United Farmers Cooperative shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on United Farmers Cooperative's skill of judgment in the purchasing or using the same and every person using this

product does so at his own risk absolutely. No representative of United Farmers Cooperative has the authority to add or alter these.

<p>IN EMERGENCY DIAL</p> <p>000</p> <p>POLICE OR FIRE BRIGADE</p>	<p>FOR SPECIALIST ADVICE IN AN EMERGENCY CALL</p> <p>1800 705 766</p> <p>(24 HOURS)</p>
--	--