

Main panel

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



Scorpio[®]

ORNAMENTAL FUNGICIDE

**ACTIVE CONSTITUENTS: 200 g/L TEBUCONAZOLE
100 g/L TRIFLOXYSTROBIN**

GROUP 3 11 FUNGICIDE

*

For the control of fungal diseases in ornamental plants and Myrtle Rust in ornamentals and non-fruit-bearing plants of the Myrtaceae family as specified in the Directions for Use table.

IMPORTANT: READ THE ATTACHED BOOKLET BEFORE USE.

* 500 mL or 1 L

Rear panel

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product. DO NOT re-use empty containers for any other purpose.

SAFETY DIRECTIONS

When opening the container and preparing spray and using prepared spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length chemical resistant gloves. If applying by spraying equipment carried on the back of the user, wear cotton overalls over normal clothing buttoned to the neck and wrist and elbow-length chemical resistant gloves. Wash hands after use. After each day's use wash gloves and contaminated clothing

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26; New Zealand 0800 764 766.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet which may be found at www.bayeres.com.au.

EXCLUSION OF LIABILITY

This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty Ltd accepts no liability or responsibility for loss or damage from failure to follow such directions and instructions.

APVMA Approval Number: 66328 / 61247

Scorpio® is a Registered Trademark of the Bayer Group.

Bayer CropScience Pty. Ltd.
ABN 87 000 226 022
391-393 Tooronga Rd
Hawthorn East. Vic. 3123
Telephone: 1800 804 479
Website: www.bayeres.com.au

For 24 hour specialist advice in an emergency only phone 1800 033 111

Batch Number:

DOM:

Barcode

Booklet front page

CAUTION

KEEP OUT REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

SCORPIO® ORNAMENTAL FUNGICIDE

ACTIVE CONSTITUENTS: 200 g/L TEBUCONAZOLE

100 g/L TRIFLOXYSTROBIN

For the control of fungal diseases in ornamental plants and Myrtle Rust in ornamentals and non-fruit-bearing plants of the Myrtaceae family as specified in the Directions for Use table.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product. DO NOT re-use empty containers for any other purpose.

SAFETY DIRECTIONS

When opening the container and preparing spray and using prepared spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length chemical resistant gloves. If applying by spraying equipment carried on the back of the user, wear cotton overalls over normal clothing buttoned to the neck and wrist and elbow-length chemical resistant gloves. Wash hands after use. After each day's use wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26; New Zealand 0800 764 766.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet which may be found at www.bayeres.com.au.

EXCLUSION OF LIABILITY

This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty Ltd accepts no liability or responsibility for loss or damage from failure to follow such directions and instructions.

IMPORTANT: READ THIS BOOKLET BEFORE USE.

APVMA Approval Number: 66328 / 61247

Scorpio® is a Registered Trademark of the Bayer Group.

Booklet inside

DIRECTIONS FOR USE

SPRAY DRIFT RESTRAINTS

DO NOT apply with spray droplets smaller than a **MEDIUM** spray droplet size category according to nozzle manufacturers specifications that refer to the ASAE S572 Standard or the BCPC Guideline.

DO NOT apply when wind speed is less than 3 or more than 20 km per hour as measured at the application site.

DO NOT apply during surface temperature inversion conditions at the application site.

Users of this product **MUST make an accurate written record** of the details of each spray application within 24 hours following application and **KEEP** this record for a minimum of 2 years. The spray application details that must be recorded are:

1 date with start and finish times of application;

2 location address and paddock/s sprayed;

3 full name of this product;

4 amount of product used per hectare and number of hectares applied to;

5 crop/situation and weed/pest;

6 wind speed and direction during application;

7 air temperature and relative humidity during application;

8 nozzle brand, type, spray angle, nozzle capacity and spray system pressure measured during application;

9 name and address of person applying this product.

(Additional record details may be required by the state or territory where this product is used.)

MANDATORY NO-SPRAY ZONES

DO NOT apply if there are aquatic or wetland areas including aquacultural ponds within **10 metres** downwind from the application area.

DO NOT apply if there are non-target vegetation or animal habitat within **15 metres** downwind from the application area.

DIRECTIONS FOR USE (continued)

SITUATION	DISEASE	RATE	CRITICAL COMMENTS
Azaleas	Azalea petal blight (<i>Ovulinia azaleae</i>)	75 mL / 100 L water	Apply when disease symptoms first appear and then repeat at 7 – 10 day intervals. When conditions favour disease development the shorter spray interval should be considered. Spray to run-off ensuring thorough coverage of all foliage including the underside of leaves.
Ornamentals	Botrytis blight or Grey mould (<i>Botrytis cinerea</i>); Downy Mildew (<i>Peronospora</i> spp.); Leaf spots (<i>Pseudomonas</i> spp.); Rusts (<i>Uromyces erythronii</i> ; <i>Puccinia</i> spp.)	75 mL / 100 L water	Curative: For existing infections, spray to run-off ensuring thorough coverage of all foliage including the underside of leaves in a series of 3 applications at 5 day intervals. Preventive: Spray to run-off ensuring thorough coverage of all foliage including the underside of leaves every 7 to 14 days according to infection pressure or beginning immediately after onset of the disease.
Roses	Black Spot (<i>Diplocarpon rosae</i>)		
Ornamentals; Roses	Powdery mildew (<i>Sphaerotheca pannosa</i> var. <i>rosae</i> ; <i>Erysiphe cichoracearum</i> ; <i>Oidium</i> spp.)	150 mL / 100 L water	Spray to run-off ensuring thorough coverage of all foliage including the underside of leaves every 7 - 14 days according to infection pressure or beginning immediately after onset of the disease.
Ornamentals and non-fruit-bearing plants of the Myrtaceae family	Myrtle Rust (<i>Uredo rangelii</i>)	75 mL / 100 L water	Apply at first signs of disease or when conditions favour disease development. Curative: For existing infections, apply 2 applications at a 14-day interval. Spray to run-off ensuring thorough coverage of all foliage including the underside of leaves. Preventive: Spray to run-off ensuring thorough coverage of all foliage including the underside of leaves every 14 days according to infection pressure.

**NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS
AUTHORISED UNDER APPROPRIATE LEGISLATION**

GENERAL INSTRUCTIONS

Application: Good control requires good coverage. Application should be made using sufficient water to ensure thorough coverage.

Precaution: Scorpio® Ornamental Fungicide has been used on a range of ornamental plant species (i.e. azaleas, begonias, chrysanthemums, cinerea, geraniums, gerbera, Myrtaceae, native daisy, pelargoniums, roses, snapdragon) without damage. Some species and varieties however are particularly sensitive to chemical sprays and as this sensitivity is often related to local conditions, it is advisable to treat only a small number of plants first, in order to ascertain the reaction before treating the whole target group.

FUNGICIDE RESISTANCE STRATEGY

GROUP	3	11	FUNGICIDE
--------------	----------	-----------	------------------

For fungicide resistance management **Scorpio** is a Group 3 & 11 fungicide. Some naturally occurring individual fungi resistant to **Scorpio** and other Group 3 & 11 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by **Scorpio** and other Group 3 & 11 fungicides, thus resulting in a reduction in efficacy. Since the occurrence of resistant fungi is difficult to detect prior to use, Bayer CropScience accepts no liability for any losses that may result from the failure of **Scorpio** to control resistant fungi.

Compatibility: As formulations of other manufacturers' products are beyond the control of Bayer Environmental Science, all mixtures should be tested prior to mixing commercial quantities.

RE-ENTRY PERIOD:

DO NOT enter treated areas until spray has dried for low and medium exposure activities such as irrigation, scouting, thinning and weeding of immature/low foliage plants and/or mature/high foliage plants, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical-resistant gloves. Clothing must be laundered after each day's use.

DO NOT enter for five (5) days after application for high exposure activities such as hand harvesting, pruning and pinching, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical-resistant gloves. Clothing must be laundered after each day's use.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS: DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT: DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

Scorpio® is a Registered Trademark of the Bayer Group.

Bayer CropScience Pty. Ltd.
A.B.N. 87 000 226 022
391-393 Tooronga Rd
Hawthorn East, Vic. 3123
Phone: (03) 9248 6888
Fax: (03) 9248 6803
Technical enquiries: 1800 804 479
Website: www.bayeres.com.au

For 24 hour specialist advice in an emergency only phone 1800 033 111
