

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



HERBICIDE

ACTIVE CONSTITUENT: 200 g/L TEPRALOXYDIM
SOLVENT: 719 g/L LIQUID HYDROCARBONS

For the control of certain grass weeds in canola, chickpeas, faba beans, field peas, lentils, lupins, subclover and vetch, as per the DIRECTIONS FOR USE table.

GENERAL INSTRUCTIONS

RESISTANT WEEDS WARNING

ARAMO Herbicide is a member of the cyclohexanedione group **GROUP A HERBICIDE** of herbicides. The product has the inhibition of acetyl coA carboxylase mode of action. For weed resistance management, the product is a Group A herbicide.

Some naturally-occurring weed biotypes resistant to the product and other Group A herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other Group A herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, BASF Australia Ltd accepts no liability for any losses that may result from the failure of this product to control resistant weeds.

MIXING

To ensure even mixing, half-fill the spray tank with clean water and add the required amount of product. Add spray additive and agitate thoroughly, then add the remainder of the water. Agitate again before spraying commences.

APPLICATION

Apply ARAMO by ground application only. ARAMO should be applied with calibrated spray equipment producing a median droplet range of 200 to 300 microns VMD. Apply in a minimum of 50 litres of water per hectare. Use 150 L/ha when spraying dense populations.

COMPATIBILITY

ARAMO is compatible with the following grass herbicides: Verdict[†], Shogun[†], Targa[†] and Fusion[†].

ARAMO is also compatible with Lontrel[†].

ARAMO may also be applied in tank mixtures with any one of the following products: Fastac[®] Duo, omethoate, Talstart[†], deltamethrin, chlorpyrifos.

ARAMO may also be mixed with dimethoate but only at the higher rates of ARAMO.

The following adjuvants are compatible with ARAMO: Hasten, D-C Trate[†], Kwickin, Uptake[†] and Enhance[†].

Do NOT mix ARAMO with Basagran[®].

CROP SAFETY

Do NOT apply after the commencement of stem elongation in canola.

RE-ENTRY PERIOD

Do NOT allow entry into treated areas until the spray has dried. If prior entry is necessary, wear cotton overalls buttoned to the neck and wrist and chemical resistant gloves. Clothing must be laundered after each day's use.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

Do NOT apply under weather conditions or from spraying equipment that may cause spray to drift onto nearby susceptible plants/crops, cropping lands, pastures or non-target plants.

Do NOT apply within 5 m upwind of sensitive native grasslands.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Do NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways,

BASF

desirable vegetation and tree roots. Empty containers and product should NOT be burnt.

SAFETY DIRECTIONS

Will irritate the eyes and skin. Avoid contact with eyes and skin. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. Wash hands after use. When opening the container and preparing spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow-length nitrile gloves and face shield or goggles. When using the prepared spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length PVC or nitrile gloves. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Telephone 131126 Australia-wide.

MSDS

Additional information is listed in the Material Safety Data Sheet.

CONDITIONS OF SALE: All conditions and warranties rights and remedies implied by law or arising in contract or tort whether due to the negligence of BASF Australia Ltd. or otherwise are hereby expressly excluded so far as the same may legally be done provided however that any rights of the Buyer pursuant to non excludable conditions or warranties of the Trade Practices Act 1974 or any relevant legislation of any State are expressly preserved but the liability of BASF Australia Ltd. or any intermediate Seller pursuant thereto shall be limited if so permitted by the said legislation to the replacement of the goods sold or the supply of equivalent goods and all liability for indirect or consequential loss or damage of whatsoever nature is expressly excluded. This product must be used or applied strictly in accordance with the instructions appearing hereon. This product is solely sold for use in Australia and must not be exported without the prior written consent of BASF Australia Ltd.

® = Registered trademark of BASF

+ Other Trademarks

© Copyright, 2003

NRA Approval No.: 55220/0603

THIS PRODUCT IS NOT CONSIDERED TO BE A
DANGEROUS GOOD UNDER THE AUSTRALIAN CODE
FOR THE TRANSPORT OF DANGEROUS GOODS BY
ROAD AND RAIL

FOR SPECIALIST ADVICE IN AN
EMERGENCY ONLY
PHONE 1 800 033 111
TOLL FREE – ALL HOURS – AUSTRALIA-WIDE

BASF Australia Ltd

ABN 62 008 437 867

Norwest Business Park, 7 Maitland Place

Baulkham Hills NSW 2153

Customer Service Telephone:

1800 635 550

Customer Service Fax:

1800 630 005

Product Information Line:

1800 501 940

Fax On Demand:

0500 544 044

Website:

www.agro.basf.com.au

BASF

DIRECTIONS FOR USE

RESTRAINTS: Do NOT apply without the addition of an adjuvant.
 Do NOT apply to crops or weeds stressed by factors such as root or foliar diseases, water-logging, nutrient deficiencies, or extremes of temperature and moisture.
 Do NOT apply if rain is expected within one hour of application.
 Do NOT apply ARAMO Herbicide more than once to any one crop.

CROP	WEEDS CONTROLLED	STAGE OF WEED GROWTH	RATE (mL/ha)	CRITICAL COMMENTS
Canola, chickpeas, faba beans, field peas, lentils, lupins, subclover, vetch	Annual ryegrass (<i>Lolium rigidum</i>)	2 leaf to fully tillered	175 to 300	Always apply with Hasten [†] or Kwickin [†] at 1 L/100 L spray volume, or similar adjuvant at the label rate (See COMPATIBILITY section). The lower doses will provide effective control if applied under ideal conditions to susceptible weeds that are smaller, actively growing and free from temperature or water stress. Do NOT apply after the commencement of stem elongation in canola. Use only in clover and vetch crops grown for seed production. Do NOT graze or cut green forage or hay for stockfeed.
	Barley grass (<i>Hordeum leporinum</i>), brome grass (<i>Bromus</i> spp.), paradoxa grass (<i>Phalaris paradoxa</i>), wild oats (<i>Avena</i> spp.), volunteer barley (<i>Hordeum vulgare</i>), volunteer oats (<i>Avena sativa</i>), volunteer triticale (<i>Triticale</i> sp.), volunteer wheat (<i>Triticum aestivum</i>)	2 leaf to fully tillered	175 to 250	

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS**HARVEST**

CANOLA: NOT REQUIRED WHEN USED AS DIRECTED.

CHICKPEAS, FABA BEANS, FIELD PEAS, LENTILS, LUPINS:

DO NOT HARVEST FOR 12 WEEKS AFTER APPLICATION.

GRAZING

CANOLA, CHICKPEAS, FABA BEANS, FIELD PEAS, LENTILS, LUPINS, SUBCLOVER, VETCH:

DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 4 WEEKS AFTER APPLICATION.