

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Farmoz

Storite® Flowable SC

FUNGICIDE

ACTIVE CONSTITUENT: 500 g/L THIABENDAZOLE

GROUP **1** FUNGICIDE

For control of Dry Rot, Gangrene and Silver Scurf of Potato tubers and specific rots and moulds of Fruit and fungal diseases of Bulbs, Corms, Turf and Mushrooms as specified in the Directions for Use table.



www.farmoz.com.au

DIRECTIONS FOR USE:

RESTRAINTS: Material must be kept in suspension at all times by continuous agitation. If agitation stops ensure the mix is resuspended before using.

CROP	PEST	STATE	RATE	CRITICAL COMMENTS
Apples and Pears	Blue Mould (<i>Penicillium expansum</i>), Grey Mould (<i>Botrytis cinerea</i>), Fruit Rot (<i>Gloeosporium album</i>)	NSW, Vic, Qld, SA, WA, Tas only	900 mL STORITE to 450 L water	Immerse fruit for 30 seconds within 24 hours of harvesting. STORITE should only be used for the post-harvest treatment of Apples and Pears where fruit is dipped in either diphenylamine or ethoxyquin prior to storage.
Bananas	Squirter Disease (<i>Nigrospora musae</i>), Black-end Rot (<i>Gloeosporium musarum</i>)	NSW, WA only	27 mL STORITE to 65 L water (200 ppm or 0.2 g/L)	After the Bananas have passed through the dehanding and/or washing operation, the fruit should be completely dipped in the prepared suspension of STORITE for 2-4 minutes.
	Crown Rot and Black-end Rot (<i>Colletotrichum musae</i>) also <i>Fusarium</i> , <i>Deightoniella</i> and <i>Verticillium</i> spp.	Qld, WA only	54 mL STORITE to 65 L water (400 ppm or 0.4 g/L)	
Bulbs and Corms	<i>Fusarium</i> Basal Rot and <i>Penicillium</i> Blue Mould	NSW, Vic, Qld, SA, WA, Tas only	900 mL STORITE to 450 L water	Dip for 15-30 minutes within 24 hours of digging. After treatment, dry bulbs in a shaded and well-ventilated areas.
Citrus	Green and Blue Mould Rot (<i>Penicillium digitatum</i> , <i>P. italicum</i>), Stem End Rot (<i>Phomopsis citri</i>)	NSW, Vic, Qld, SA, WA only	900 mL STORITE to 450 L water (1,000 ppm or 1 g/L)	–
Mushrooms	Wet Bubble Disease (<i>Mycogone perniciosa</i>)	All States	23 mL STORITE per 50 kg bale of peat moss (50 kg peat moss equals approx. 170-200 L)	Incorporate in water used to wet the peat moss at casing.
Potatoes	Dry Rot (<i>Fusarium</i> spp.), Gangrene (<i>Phoma exigua</i>), Silver Scurf (<i>Helminthosporium solani</i>)		1 L of STORITE to 22 L water (provides 22 g/L of thiabendazole)	Apply the suspension as a mist at the rate of 2 litres per ton over the Potatoes on the conveyor line prior to storage.

CROP	PEST	STATE	RATE	CRITICAL COMMENTS
Turf Disease	Dollar Spot (<i>Sclerotinia homeocarpa</i>)	NSW, Vic, Qld, SA, WA only	38 mL STORITE per 100 m ² in 50-100 L of water	Apply as a protective spray when infection first appears and then at 2-3 week interval as required. For extensive damage, initial treatment should be at the high rate and then at 2-3 week intervals at the low rate.
			56 mL STORITE per 100 m ² in 50-100 L of water (900 mL treats 1,600 m ²)	
	Fusarium Patch (<i>Fusarium nivale</i>)	NSW, Vic, Qld, SA, WA, Tas only	76 mL STORITE per 100 m ² in 50-100 L of water (900 mL treats 1,600 m ²)	Water in lightly. Repeat applications after 2-3 weeks may be required until disease is controlled.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS: NOT REQUIRED WHEN USED AS DIRECTED.

GENERAL INSTRUCTIONS

STORITE® FLOWABLE SC is recommended for the post-harvest protection of Apples and Pears, Bananas and Citrus Fruit from specific rots and moulds; and for the treatment of specific fungal diseases in Bulbs, Corms, Mushrooms and Turf.

STORITE FLOWABLE SC also effectively controls Dry Rot (*Fusarium* spp.), Gangrene (*Phome exigua*) and Silver Scurf (*Helminthosporium solani*) in Potatoes.

- Before using STORITE FLOWABLE SC mixed with other chemicals consult the manufacturer.
- Shake the container before use to ensure uniform dispersion of contents before measuring quantity required.
- Clean mixing and spraying equipment before use.
- Fill the mixing tank with water to 3/4 full and begin mechanical agitation.
- For all purposes except Mushrooms and Potatoes, add a wetting agent to the water prior to inclusion of STORITE FLOWABLE SC.
- Fruit and Bulbs should be washed prior to application of STORITE FLOWABLE SC suspension.
- Fruit must be in complete contact with the suspension for at least 30 seconds.
- Fruit should not be rinsed after treatment.
- Ensure that even and complete coverage of Potatoes is obtained by adequate rotation of tubers along the conveyor line.
- Potatoes should be treated within 3 days of harvesting and grading prior to storage. For effective control of Silver Scurf, undertake the treatment within 4 weeks of crop maturity.

FUNGICIDE RESISTANCE WARNING

FARMOZ STORITE FLOWABLE SC Fungicide is a member of the Methyl

GROUP 1 FUNGICIDE

Benzimidazole Carbamate group of fungicides. For fungicide resistance management STORITE FLOWABLE SC is a Group 1 Fungicide.

Some naturally occurring individual fungi resistant to STORITE FLOWABLE SC and other Group 1 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by STORITE FLOWABLE SC or other Group 1 fungicides, thus resulting in a reduction in efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use, FARMOZ Pty. Ltd. accepts no liability for any losses that may result from the failure of STORITE FLOWABLE SC to control resistant fungi.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with this chemical or used containers.

STORAGE AND DISPOSAL

KEEP OUT OF REACH OF CHILDREN. Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

Unused or spent dip should be disposed of carefully to avoid contamination of water sources. Either spray the dip onto a grassed area or pour into a disposal pit specifically marked and set up for this purpose, at least 50 metres away from any streams, drains, channels, wells, bore holes or water courses.

SAFETY DIRECTIONS

Harmful if absorbed by skin contact or swallowed. Avoid contact with eyes and skin. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for STORITE FLOWABLE SC is available from FARMOZ on request. Call Customer Service on (02) 9431 7800.

CONDITIONS OF SALE: The use of FARMOZ STORITE FLOWABLE SC Fungicide being beyond the control of the manufacturer, no warranty expressed or implied is given by FARMOZ Pty. Ltd., regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and FARMOZ Pty. Ltd. accepts no responsibility for any consequence whatsoever from the use of this product.

® Registered Trademark of FARMOZ Pty. Ltd.

NOT A DANGEROUS GOOD ACCORDING TO THE AUSTRALIAN DANGEROUS GOODS (ADG) CODE.

PACK SIZES: 1 L, 5 L, 10 L, 20 L