

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE
OPENING OR USING

CHEMAG CONFRONT 800 WG INSECTICIDE

ACTIVE CONSTITUENT: 800 g/kg THIODICARB
(an anti-cholinesterase compound)

GROUP 1A INSECTICIDE

For the control of caterpillar pests in
brassicas, cotton, maize, sweet corn,
pulses, tomatoes and tobacco

THIS BOOKLET IS PART OF THE LABEL.

APVMA Approval No: 59763/0905

Imtrade Australia
ABN 13 090 151 134
Suite 22, 11 Preston Street, Como, WA 6152
Tel: 1800 171 799 Fax: 1800 171 788
Web: www.imtrade.com.au

GENERAL INSTRUCTIONS

This product is a water dispersible granule formulation that readily disperses in water for spraying by ground or air equipment. It is unstable under highly acidic or highly alkaline conditions, being hydrolytically sensitive to degradation of the active constituent by strong acids, strong bases and certain heavy metals and salts.

To improve rainfastness, add a commercially available 'sticker' to the spray-tank, at the recommended rate, before applying this product.

Insecticide Resistance Warning

For insecticide resistance management, ChemAg Confront 800 WG Insecticide is a Group 1A insecticide. Some naturally occurring insect biotypes resistant to ChemAg Confront 800 WG and other Group 1A insecticide may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if ChemAg Confront 800 WG or other Group 1A insecticides are used repeatedly. The effectiveness of ChemAg Confront 800 WG on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, ChemAg Australia Ltd accepts no liability for any losses that may result from the failure of ChemAg Confront 800 WG to control resistant insects. ChemAg Confront 800 WG may be subject to specific resistance management strategies. For further information contact your local supplier, ChemAg Australia Ltd representative or local agricultural department agronomist.

DIRECTIONS FOR USE

Restraints

DO NOT add ChemAg Confront 800 WG to water with a pH below 3.0 or above 8.5.

DO NOT store prepared spray mixture for longer than 6 hours

CROP	PEST	STATE	RATE	WHP	CRITICAL COMMENTS
Brassica crops	Cabbage Cluster Caterpillar (<i>Crociodolomia pavonana</i>), cabbage white butterfly (<i>Pieris rapae</i>)	All states	470 to 940 g/ha	7 (H) 21(G)	Spray as required depending upon pest pressure. Closely monitor pest populations. Use the lower rate for small caterpillars and the higher rate for advanced caterpillars. Add a non-ionic wetting agent at the recommended rates.

CROP	PEST	STATE	RATE	WHP	CRITICAL COMMENTS
Cotton	Heliothis (Budworms) (<i>Helicoverpa</i> <i>a spp.</i>)	Qld. NSW, WA, NT only	940 g or 1.2 kg/ha	21 (H & G)	<p>This product has ovicidal and larvicidal activity.</p> <p>Ovicidal and Larvicidal Rate: Application should be made on eggs, very small or small larvae only. The most effective timing is on egg hatch. Use the lower rate on light to moderate infestations, and the higher rate on heavier infestations. Closely monitor pest populations after application. Repeat application as needed to maintain control. Ensure thorough coverage of target areas. Larvae must feed on treated area in order to achieve effective control. Larger larvae and larvae feeding within bolls and squares may not be controlled.</p> <p>Ovicidal Rate: Ovicidal rates or ChemAg Confront 800 WG should only be used in a tank mix with the registered rate of a compatible larvicide time application on egg threshold levels prior to hatching and closely monitor the population. Use the lower rate of ChemAg Confront 800 WG where egg pressure is less than 3 eggs per terminal. Use the higher rate of ChemAg Confront 800 WG for larger larvae up to 8 mm and under higher egg pressure. Repeat application to maintain control. Ensure thorough coverage of target areas.</p>
			235 g to 470 g/ha PLUS the registered rate of a compatible larvicide		

CROP	PEST	STATE	RATE	WHP	CRITICAL COMMENTS
Maize, sweetcorn	Heliothis (Helicoverpa armigera)	Qld, NSW, Vic, Tas, WA only	700 to 940 g/ha	Maize 7 (H & G) Sweet corn 7 (H) 21 (G)	Closely monitor pest populations during tasselling-silking. If high numbers or medium to large caterpillars are in the tassels, spray at tassel emergence. Egg laying is heaviest during green silk formation. Time of application to coincide with egg hatching. Ensure thorough coverage of target areas.
Pulses e.g. Navy beans, soybeans, mungbeans, pigeon peas, chickpeas	Heliothis (Budworms) (<i>Helicoverpa spp.</i>)	All states	235 or 350 g/ha	21 (H&G)	Time application to coincide with egg hatching and closely monitor pest populations. Use the lower dose rate on eggs and very small larvae (up to 3 mm) and light to moderate infestations; and the higher rate on small larvae (up to 7 mm) and heavier infestations. Repeat application as needed to maintain control. Ensure thorough coverage of target areas.
Tomatoes	Heliothis (Budworms) (<i>Helicoverpa spp.</i>)	All states	<u>Low Volume:</u> 655 g/ha <u>High Volume:</u> 33 g/100 L water	1 (H) 21(G)	Time application to coincide with egg hatching and closely monitor pest populations. Repeat applications as needed to maintain control. Ensure thorough coverage of target areas by increasing water volume with plant growth stage. NOTE: Low volume is defined as 400 L or less total spray volume per hectare. High volume is defined as 1000 L or more total spray volume per hectare.

(continued from previous page)

CROP	PEST	STATE	RATE	WHP	CRITICAL COMMENTS
Tomatoes	Heliiothis (budworms) (<i>Helicoverpa</i> spp.) Potato moth (<i>Phthorimaea</i> <i>operculella</i>)	All states	<u>Low Volume:</u> 233 g to 655 g/ha ChemAg Confront 800WG PLUS 500 g to 1.0 kg Deifin® WG (or other compatible registered Bt product) per ha <u>High Volume:</u> 23 g to 46 g ChemAg Confront 800 WG PLUS 50 to 100 g Delfin WG (or other compatible registered Bt product) per 100 L water	1 (H) 21(G)	Time application to coincide with egg hatching and closely monitor pest populations. Repeat application as needed to maintain control. Ensure thorough coverage of target areas by increasing water volume with plant growth stage. NOTE: Low volume is defined as 400 L or less total spray volume per hectare. High volume is defined as 1000 L or more total spray volume per hectare. The lower rates or ChemAg Confront 800 WG in mixtures with Bt provide ovicidal control only. Use higher rates of ChemAg Confront 800 WG when larval control is required. Use the higher rate of Delfin WG when Potato Moth is present.
Tobacco	Heliiothis (Budworms) (<i>Helicoverpa</i> <i>a</i> spp.) Tobacco loopers (<i>Chrysodeixis</i> <i>spp.</i>)	Qld, NSW, Vic, WA only	33 g/100 L water	Nil (H) 21 (G)	Time of application to coincide with egg hatching and closely monitor pest populations. Repeat applications as needed to maintain control. Ensure thorough coverage of target areas by increasing water volume with plant growth stage.

**NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

DO NOT USE THIS PRODUCT IN THE HOME GARDEN.

WITHHOLDING PERIODS (WHP)

(H) = Harvest

Brassica, maize, sweetcorn:

DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION.

Cotton and Dried pulses:

DO NOT HARVEST FOR 21 DAYS AFTER APPLICATION.

Tomatoes:

DO NOT HARVEST FOR 1 DAY AFTER APPLICATION

Tobacco:

NOT REQUIRED WHEN USED AS DIRECTED.

(G) = Grazing

All crops except maize:

DO NOT GRAZE OR CUT FOR STOCK FEED FOR 21 DAYS AFTER APPLICATION.

Maize:

DO NOT GRAZE OR CUT FOR STOCK FEED FOR 7 DAYS AFTER APPLICATION.

Export of Treated Produce

Growers should note that suitable MRLs or import tolerances may not be established in all markets for produce treated with ChemAg Confront 800 WG Insecticide. If you are growing produce for export, please check with Imtrade Australia for the latest information on MRLs and import tolerances BEFORE using Chemag Confront 800 WG Insecticide.

Mixing

Remove oil, rust, scale, pesticide residues and other foreign matter from sprayer and strainer. Flush with clean water. Use 50 mesh or slotted strainer in spray system. Fill tank approximately half full of water, add this product and mix thoroughly by mechanical or hydraulic action. Add water to desired volume and thoroughly mix again.

Application

This product may be applied by air or ground equipment. Use a minimum prepared spray volume of 20 L/ha by air (30 L/ha for cotton, 50L/ha for maize or sweetcorn) and 50 L/ha by ground. A droplet, density of minimum 60 droplets per cm² must be attained. There needs to be some air movement, which preferably should be blowing across the flight path. After use, drain and flush sprayer with water.

Compatibility

ChemAg Confront 800 WG may be mixed with Dipel® ES, Delfin® WG, most foliar Bt formulations, Thiodan® EC, Folidol® M500, Marlin® 225, Folimat® and most EC formulations of Profenofos, Chlorothalonil and Dimethoate. ChemAg Confront 800 WG is physically compatible with Intrepid® and Reign®. However, before preparing tank mix combinations, it is advisable to mix a small amount of ChemAg

Confront 800 WG with an equal volume of water in a small container and add the other pesticide. Any mixture, which curdles, precipitates or greases should not be used. ChemAg Confront 800 WG is NOT compatible with manganese, zinc or copper based products, with Deltamethrin, with LV, ULV or other oil-based formulations, or with vegetable oil dilutant. As the formulations of other manufacturers' products are beyond the control of ChemAg Australia Ltd, all mixtures should be tested prior to mixing commercial quantities.

PRECAUTIONS

DO NOT allow children access to treated areas. When the product is being applied, human flaggers/markers must be protected by engineering controls such as vehicles with cabs.

Re-entry Period

Do not allow entry into treated areas for 1 day after treatment. When prior entry is necessary, wear cotton overalls buttoned to the neck and wrist and a washable hat, chemical resistant gloves and impervious footwear. Clothing must be laundered after each day's use.

DRIFT WARNING

DO NOT apply under weather conditions or from spraying equipment, which could be expected to cause spray to drift onto adjacent crops, crop lands pastures or livestock.

PROTECTION OF LIVESTOCK

Dangerous to bees. DO NOT spray any plants in flower while bees are foraging.

PROTECTION OF WILDLIFE, FISH CRUSTACEANS AND ENVIRONMENT AND OTHERS

Dangerous to fish and aquatic organisms. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area. Do not store for long periods in direct sunlight.

Bottle

Triple or preferably pressure rinse containers before disposal. Add rinsing to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal

pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Poisonous if absorbed by skin contact, inhaled or swallowed. May irritate eyes and skin. Avoid contact with eyes and skin. Do not inhale dust or spray mist. Obtain an emergency supply of atropine tablets 0.6 mg. When opening the container, preparing the spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist, washable hat, elbow-length PVC gloves, face shield, impervious footwear and half facepiece respirator with combined dust and gas cartridge. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shields respirator and contaminated clothing. Wash rubber respirator with detergent and warm water.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (telephone 13 11 26). If swallowed, give one atropine tablet every 5 minutes until dryness of the mouth occurs if poisoned by skin absorption or through lungs, remove any contaminated clothing, wash skin thoroughly and give tablets as above. Get a doctor or hospital quickly.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet, which can be obtained from www.imtrade.com.au

EXCLUSION OF LIABILITY

This product must be used strictly as directed, and in accordance with all instruction appearing on the label and in other reference material. So far as it is lawfully to do so, Bay Crop Science Pty Ltd accepts no liability or responsibility for loss or damage arising from failure to follow such directions and instructions.



CARBAMATE PESTICIDE, SOLID, TOXIC (contains Thiodicarb)		
UN 2757	PG II	HAZCHEM 2X
IN A TRANSPORT EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE		
FOR 24 HOUR SPECIALIST ADVICE IN EMERGENCY ONLY PHONE 13 11 26		