

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

RIZOLEX[®] 500WP

FUNGICIDE

ACTIVE CONSTITUENT: 500g/kg TOLCLOFOS-METHYL

GROUP **X** FUNGICIDE

For the control of *Rhizoctonia solani* in Cotton

GENERAL INSTRUCTIONS

MIXING:

Mix the required amount of RIZOLEX 500 WP Fungicide with a small quantity of water to form a slurry. Add slurry to the required volume and agitate thoroughly.

APPLICATION:

RIZOLEX 500 WP Fungicide does not control *Pythium spp.* and should only be applied to seed previously treated with a fungicide such as Apron* for control of *Pythium spp.* RIZOLEX 500 WP Fungicide may be applied to seed previously treated with quintozene, thiodicarb or acephate. RIZOLEX 500 WP Fungicide can be applied as an in-furrow spray or by water injection from 20 to 350 L/ha irrespective of spray band width.

COMPATIBILITY:

RIZOLEX 500 WP Fungicide is compatible with formulations of chlorpyrifos.

Fungicide Resistance Warning

GROUP **X** FUNGICIDE

For fungicide resistance management Sumitomo RIZOLEX 500 WP Fungicide is a Group X Fungicide.

Some naturally occurring individual fungi resistant to RIZOLEX 500 WP Fungicide and other Group X fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by RIZOLEX 500 WP Fungicide or any other Group X fungicides, thus resulting in a reduction of efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use, Sumitomo Chemical Australia Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant fungi.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT:

Dangerous to fish. DO NOT contaminate streams, rivers or waterways with the product or used containers.

STORAGE AND DISPOSAL:

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Shake empty bag/ keg into spray tank. DO NOT dispose of undiluted chemicals on site. Puncture or shred and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS:

Avoid contact with eyes and skin. DO NOT inhale dust or spray mist.

FIRST AID:

If poisoning occurs, contact a doctor or Poisons Information Centre (Tel. 131126).

MATERIAL SAFETY DATA SHEET:

Additional information is listed in the Material Safety Data Sheet (MSDS).

NOTICE TO BUYER

Sumitomo Chemical Australia Pty Ltd will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply or use of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatory imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such statutes the liability of Sumitomo Chemical Australia Pty Ltd is limited to the replacement of the goods or (at the option of Sumitomo Chemical Australia Pty Ltd) the refund of the price paid and is conditional upon a claim being made in writing and where possible sufficient part of the goods to enable proper examination being returned to Sumitomo Chemical Australia Pty Ltd.

[®]Registered Trademark of Sumitomo Chemical Co. Japan.

NRA Approval No: 50730/0598

THIS PRODUCT IS NOT CONSIDERED TO BE A DANGEROUS GOOD UNDER THE AUSTRALIAN CODE FOR THE TRANSPORT OF DANGEROUS GOODS BY ROAD OR RAIL	
In a Transport Emergency Dial 000 Police or Fire Brigade	SPECIALIST ADVICE IN EMERGENCY ONLY ALL HOURS - AUSTRALIA WIDE 1800 024 973

RIZOLEX[®] 500WP

DIRECTIONS FOR USE

CROP	DISEASE	STATES	RATE	CRITICAL COMMENT
Cotton	<i>Rhizoctonia solani</i>	Qld and NSW only	120g/ha or 120g/10,000m row	Apply as an in-furrow spray or by water injection at time of planting

**NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

WITHHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED.