

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

GENFARM TOLEX[®]

Liquid Fungicide

ACTIVE CONSTITUENT: 500 g/L TOLCLOFOS-METHYL

GROUP	X	HERBICIDE
--------------	----------	------------------

For the control of *Rhizoctonia solani* in
Cotton and seed borne *Rhizoctonia solani*
In Potatoes

CONTENTS: 1L, 5 L, 20L

Genfarm Crop Protection Pty Ltd
Suite 2, Level 3, 64 Talavera Road,
Macquarie Park,
NSW, 2113
Tel: (02) 9889 5400

® Registered Trademark of Genfarm Crop Protection Pty Ltd

DIRECTIONS FOR USE

CROP	DISEASE	STATES	RATE	CRITICAL COMMENTS
Cotton	<i>Rhizoctonia solani</i>	QLD and NSW only	120 mL/ha or 120 mL/10,000 m row	Apply as an in-furrow spray or by water injection at time of planting.
Potatoes	Seed borne <i>Rhizoctonia solani</i>	All States	0.4 L/tonne of seed	Do NOT apply to freshly cut seed potatoes as emergence may be affected. Allow sufficient time for cut wounds to dry and suberise before applying TOLEX Liquid or use "round" seed. DO NOT plant wet seed. Apply immediately prior to planting (within 1 day) by spray methods and allow seed to dry before planting. Seed pieces should be rotated on a roller system (e.g. grading table) to ensure complete and thorough coverage. For further information refer to the General Instructions. If conditions (e.g. high soil temperature and moisture) or seed quality favour seed piece breakdown, the addition of any moisture to the seed may enhance the problem. Under these circumstances the use of Tizolex 100D is recommended.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED.

GENERAL INSTRUCTIONS

COTTON:

TOLEX Liquid does not control *Pythium spp.* and should only be applied to seed previously treated with a fungicide such as Apron* for control of *Pythium spp.* TOLEX Liquid may be applied to seed previously treated with quintozone, thiodicarb or acephate.

Mixing

Add the required amount of TOLEX Liquid to the partly filled tank with the agitator running and complete filling the tank with water. Do not mix with alkaline water. Continue thorough agitation during spraying and after stoppage. Do not let prepared spray solution sit in spray tank overnight.

Application

TOLEX Liquid can be applied as an in furrow spray or by water injection using 20 to 350 L/ha of water irrespective of spray band width.

POTATOES:

TOLEX Liquid ensures protection of potatoes from seed borne *Rhizoctonia solani*. Best results will be obtained where other crop management practices (including crop rotation) are employed to reduce *Rhizoctonia solani* infection in the soil.

Application

Mix the required quantity of TOLEX Liquid with 5 L of water per tonne of seed. Ensure constant agitation of the spray tank during application. Apply only when seed pieces are being rotated using spraying methods designed to ensure complete coverage. Only treat enough seed to meet planting needs. Ensure the seed is dry prior to planting. DO NOT plant wet seed. Ensure that treated seed remains separated from untreated seed at all times.

FUNGICIDE RESISTANCE WARNING

For fungicide resistance management TOLEX Liquid is a Group X Fungicide. Some naturally occurring individual fungi resistant to TOLEX Liquid and other Group X fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by TOLEX Liquid or any other Group X fungicides, thus resulting in a reduction of efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use, Genfarm Crop Protection Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant fungi.

PRECAUTIONS

DO NOT use treated seed for animal or human consumption.

DO NOT allow treated seed to contaminate grain or other seed intended for animal or human consumption.

DO NOT feed treated seed or otherwise expose to wild life or domestic birds.

When treated seed potatoes are stored they should be kept apart from other potatoes and the bags or containers should be clearly marked to indicate the contents have been treated with this product. Bags and containers which have held treated seed potatoes are not to be used for any other purpose.

RE-ENTRY TO TREATED AREA

The following conditions MUST be observed when re-entry is required before the spray has dried:

Wear impervious boots. Do not re-use footwear until thoroughly aired.

RE-HANDLING TREATED SEED

The following conditions MUST be observed when handling treated seed prior to planting.

Wear cotton overalls buttoned to the neck and wrists (or equivalent clothing) and elbow-length PVC gloves. After each day's use, wash gloves and clothing.

PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT feed treated seed to animals including poultry. Dangerous to fish. DO NOT contaminate streams, rivers or waterways with the product or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. Break, crush, puncture and bury empty containers in a local authority landfill. If not available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

May irritate the eyes and skin. Avoid contact with eyes and skin. Wash hands after use.

When mixing and loading wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length PVC gloves. When using in enclosed areas wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length PVC gloves. Ensure adequate ventilation during use. After each day's use, wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia: 131126).

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet (MSDS) which is available from the supplier.

CONDITIONS OF SALE

The use of this product is beyond the control of Genfarm Crop Protection Pty Ltd. No warranty is expressed or implied regarding the suitability or efficiency for any purpose for which the buyer uses it. Genfarm Crop Protection Pty Ltd accepts no responsibility for any consequences resulting from the use of this product. Genfarm Crop Protection Pty Ltd will not be held liable for any loss, injury or damage arising from the sale, supply or use of this product, whether through negligence or otherwise. No responsibility will be accepted for any consequences whatsoever resulting from the use of this product.

*Registered trademarks

THIS PRODUCT IS NOT CONSIDERED TO BE A DANGEROUS GOOD UNDER THE AUSTRALIAN CODE FOR THE TRANSPORT OF DANGEROUS GOODS BY ROAD OR RAIL
In a Transport Emergency Dial 000 Police or Fire Brigade
SPECIALIST ADVICE IN EMERGENCY ONLY ALL HOURS – AUSTRALIA WIDE 1800 024 973

APVMA Approval Number: 62158/1L/0208, 62158/5L/0208, 62158/20L/0208

Batch No:

Date of Manufacture: