

POISON

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS**

Bayfidan[®] Turf

FUNGICIDE

Active
Constituent:
250 g/L TRIADIMENOL
Solvent: 602 g/L N-METHYL-2-PYRROLIDONE

GROUP C FUNGICIDE

For the systemic control of dollar spot
and other diseases of turf as specified
in the directions for use table

Bayer 

Chipco[®]

Bayfidan® Turf

Fungicide

Packaging: 1 L

General Instructions

Fungicide Resistance Warning

GROUP C FUNGICIDE

Bayfidan Turf is a member of the DMI group of fungicides. For fungicide resistance management the product is a Group C fungicide. Some naturally occurring individual fungi resistant to the product and other Group C fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by Bayfidan and other Group C fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Bayer Australia Limited accepts no liability for any losses that result from failure of this product to control resistant fungi.

Mixing

Add the required quantity of Bayfidan Turf to water in the spray vat while stirring or with agitators in motion.

Bayfidan Turf is a broad spectrum, systemic, foliar fungicide that is quickly taken up by plants and translocated within the plant. Bayfidan Turf provides protection on a monthly spray program and acts curatively after infection has taken place. Because of its systemic activity, Bayfidan Turf is less affected by rainfall or irrigation than contact type fungicides.

Compatibility

Bayfidan Turf is compatible with Baythroid® Turf. Do not mix concentrates together but add each to the spray tank separately. As formulations of other manufacturer's products are beyond the control of Bayer Australia Limited, all mixtures should be tested prior to mixing commercial quantities. As changes in climatic conditions can alter the sensitivity of plants to mixtures of sprays, Bayer cannot be responsible for the behaviour of such mixtures.

Protection of Crops, Native and Non-target Plants

Do not apply earlier than 5 weeks after sowing.

Protection of Wildlife, Fish, Crustaceans and Environment

Harmful to fish. DO NOT contaminate streams, river or waterways with the chemical or used containers.

Storage and Disposal

Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Rinse container before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on-site. Dispose of at a local authority landfill. If no landfill is available, bury the container below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

Safety Directions

Will irritate the eyes and skin. Avoid contact with eyes and skin. Do not inhale spray mist. When preparing spray, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow-length butyl rubber gloves and goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing.

First Aid

If poisoning occurs, contact a doctor or Poisons Information Centre (131126). If swallowed, do NOT induce vomiting. Give a glass of water. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

For further information refer to the Material Safety Data Sheet for the product.

Liability

This product must be used strictly as directed. Bayer Australia Limited accepts no responsibility for loss or damage arising from failure to follow directions for use.

®Trademark of Bayer Germany

MN 1011226 12/00

©Copyright Bayer Australia Limited

NRA Approval Number 30676/0499

Directions for use

Crop	Disease	Rate	Critical Comments
Turf	Dollar spot (Not Tasmania, NT)	Preventive 30 mL/100 m ²	Apply thoroughly on a monthly program starting immediately the disease appears.
		Curative 60 mL/100 m ²	Apply thoroughly under conditions of heavy disease pressure; then treat as necessary.
Turf	Black Helminthosporium (QLD NSW TAS ONLY)	60 mL/100 m ²	Use high volume equipment. Low volume may be used provided turf is irrigated thoroughly after spraying. Apply during period of turf growth on a monthly program.
Turf	Take-all patch (Not SA, NT)	60 mL/100 m ²	Preventive only after soil fumigation with methyl bromide. Apply thoroughly, using high volume equipment, 6-8 weeks after seeding. Repeat applications (every 8 weeks in the first two years) for continued protection.
Turf	Winter Fusarium (NSW VIC TAS ONLY)	60 mL/100 m ²	Apply thoroughly on a monthly program from end of March until danger of infection has passed.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS
AUTHORISED UNDER APPROPRIATE LEGISLATION**