


634-636 Casella Place, Kewdale WA 6239 Tel: (08) 9353 4300 Fax: (08) 9353 4144

Approved by: _____
Date: _____

Label image proudly supplied by




infopest@dpi.qld.gov.au




WESTERN AUSTRALIA
6344-128

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



TRIADIMEFON 500 FUNGICIDE WP
ACTIVE CONSTITUENT: 500g/kg TRIADIMEFON


GROUP C FUNGICIDE



259mm

7.5mm

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



TRIADIMEFON 500 FUNGICIDE WP
ACTIVE CONSTITUENT: 500g/kg TRIADIMEFON

GROUP C FUNGICIDE

WHEAT:
For control of leaf and stripe rust, septoria tritici blotch, powdery mildew and suppression of Take All.

BARLEY:
For control of powdery mildew.

PRIMARY PACK CONTAINS: 10 x 1kg MEASURE PACKS
WHICH IT IS ILLEGAL TO SELL SEPARATELY
DO NOT DESTROY OUTER CONTAINER WHILE PRODUCT STILL REMAINS.

4FARMERS PTY. LTD.
A.C.N. 067 443 485
643 MURRAY STREET,
WEST PERTH, WESTERN AUSTRALIA 6005.

10Kg NET


APVMA Approval No:50160/10/0505

Batch Number: D.O.M.

TRIADIMEFON 500 FUNGICIDE WP

307mm

4FARMERS PTY. LTD.
A.C.N. 067 443 485
643 MURRAY STREET, WEST PERTH,
WESTERN AUSTRALIA 6005.
PH: (08) 9356 3445 FAX: (08) 9356 3447
TOLL FREE: 1800 038 445



GENERAL INSTRUCTIONS

- 4Farmers Triadimefon 500 Fungicide WP may be used by Aircraft or Ground spraying equipment.
- Always follow the label directions for use.
- Determine the susceptibility of the wheat variety, eg Dept of Agriculture data, to determine spray requirements.
- Timing of application: When approx 20% of leaves show signs of infection during tillering or jointing, spray within a week.
- Application: Spray within one week if approx 10% of leaves show infection. Do not delay.
- Weather: Do not spray if rain is forecast within 24 hours.
- Temperature: Do not spray if temperature is below 10°C.
- Wind: Do not spray if wind speed is greater than 15 km/h.
- Humidity: Do not spray if relative humidity is less than 70%.
- For Stripe Rust: Re-spraying may be required as reinfection can occur. Regularly monitor crops at risk from 3 weeks after application to assess if re-application is required.

FUNGICIDE RESISTANCE WARNING:

Triadimefon is a member of the D.M.I. group of fungicides. For fungicide resistance management, 4Farmers Triadimefon 500 Fungicide WP is a Group C fungicide. Some naturally occurring individual fungi resistant to 4Farmers Triadimefon 500 Fungicide WP and other Group C fungicides may occur through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled by 4Farmers Triadimefon 500 Fungicide WP and other Group C fungicides, thus resulting in efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use, 4Farmers Pty Ltd accepts no liability for any losses that may result from the failure of 4Farmers Triadimefon 500 Fungicide WP to control resistant fungi.

MIXING:

- The required quantity of 4Farmers Triadimefon 500 Fungicide WP should be thoroughly mixed with a small quantity of clean filtered water to make a creamy mix. Add this slurry to the required quantity of water in the tank and mix thoroughly. Agitate prior to and during spraying.
- DO NOT mix more than can be used for the day's spraying.
- DO NOT tank mix with trace elements or other chemicals without first checking compatibility.

SPRAYING:

- Ground Rig: Use a convenient amount of water, eg 30 - 100 L water / Ha.
- Aircraft: Apply at not less than 10 L / Ha, preferably in a cross wind of greater than 5 knots. If this is not possible, increase the volume to 20 L / Ha and increase the droplet size. Spray in the morning and reduce the swath width. Boom, with fan nozzles set forward at 45° forward to the boom and 'Micro' nozzles with blade settings of 30 - 50° are equally satisfactory. Use proper measures for gauging wind speed, direction and variability of the wind.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT

- DO NOT feed treated leaf matter to wild or domestic birds or other animals. Observe withholding periods before collecting. Harmful to fish.
- DO NOT contaminate streams, rivers or waterways with the chemical or use containers.

STORAGE AND DISPOSAL AND PROTECTION OF OTHERS

- Store the unopened product in the closed original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Share empty bag. If using as a liquid, single rinse the bag before disposal and add rinsings to the spray tank. Do not dispose of unlabelled chemical waste. Puncture and burn the bag in a safe place. If the bag is not to be used, bury the container to a depth of 200 mm in a disposal site specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS:

- Poisonous if swallowed. May irritate the eyes and skin. Do not inhale dust or spray mist. When preparing product for use, wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves and a disposable dust mask covering mouth and nose.
- After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.
- After each day's use, wash gloves and contaminated clothing.

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 131126). MSDS: This can be obtained from your supplier or 4Farmers Pty Ltd on (08) 9356 3445 Toll Free 1800 038 445


CONDITIONS OF SALE:

The use of 4Farmers Triadimefon 500 Fungicide WP, being beyond the control of the manufacturer, no warranty expressed or implied is given by 4Farmers Pty Ltd, regarding its availability, fitness or efficacy for any purpose for which it is used by the buyer, whether in accordance with the directions or not and 4Farmers Pty Ltd accepts no responsibility for any consequences whatsoever resulting from the use of this product.

TRIADIMEFON 500 FUNGICIDE WP

262mm

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



TRIADIMEFON 500 FUNGICIDE WP
ACTIVE CONSTITUENT: 500g/kg TRIADIMEFON

GROUP C FUNGICIDE

WHEAT:
For control of leaf and stripe rust, septoria tritici blotch, powdery mildew and suppression of Take All.

BARLEY:
For control of powdery mildew.

PRIMARY PACK CONTAINS: 10 x 1kg MEASURE PACKS
WHICH IT IS ILLEGAL TO SELL SEPARATELY
DO NOT DESTROY OUTER CONTAINER WHILE PRODUCT STILL REMAINS.

4FARMERS PTY. LTD.
A.C.N. 067 443 485
643 MURRAY STREET,
WEST PERTH, WESTERN AUSTRALIA 6005.

10Kg NET

APVMA Approval No:50160/10/0505

Batch Number: D.O.M.

TRIADIMEFON 500 FUNGICIDE WP

307mm

371mm

DIRECTIONS FOR USE TABLE

Crop	Disease/Pest	Rate	WHP months	Critical Comments
Barley	Powdery mildew	All States 250g/ha	3	Apply at the first signs of mildew
Wheat	Leaf Rust	NSW, VIC, SA, WA, ONL 125 - 250g/ha		For spray timings see general instructions. Use 250g/ha where longer residual control is required particularly in periods of high risk during rust epidemics. Repeat spraying may be required with Stripe Rust. Regularly monitor crops from 3-4 weeks after spraying for signs of reinfection and treat accordingly.
	Stripe Rust	NSW, VIC, SA, WA, ONL 125 - 250g/ha		
Wheat	Septoria tritici blotch	Mid and North Western Victoria 250g/ha		Where this disease commonly affects the flag leaf, apply to susceptible varieties at the 3-4 tiller stage. Re-infestation can occur if most conditions prevail
	Stn	NSW, Vic only 125 - 250g/ha		For control of light to moderate infections. Applications should coincide with early head emergence. Tank mixes with Benzyl may improve control in severe infections

APPLIED AS A FERTILISER TREATMENT

Crop	Disease/Pest	Rate	WHP months	Critical Comments
Barley	Powdery Mildew	NSW, VIC, SA, WA, ONL 200 g/ha	3	Use the higher rate where conditions are conducive to very high rates of disease pressure. Also in years when the incidence of disease is expected to be high as forecast by Department of Agriculture or reputable disease forecasters. Application: Apply product evenly onto the fertilizer. If using DAP or similar granulated product, use as powder. If adding to a particularly dusty - dry fertilizer granule, slize with minimum water (up to a maximum of 8 Litres) ensure consistent mixing of the suspension using a 'granny pot' or similar device. Calculation of the rate required / ton of Fert is below: 1000 = (Fert rate / ha) X rate of fungicide (kg) / kg using 100 kg Fert / ha, and wetting 200 g/ha fungicide: 1000 = 100 X 0.2 = 2 kg / ton. Regularly monitor crops from about 10-12 weeks after planting assess the need for follow-up disease control (eg stripe rust) as the residual protection from the fertilizer application declines.
Wheat	Leaf Rust			
	Suppression of Take-All			

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

- DO NOT HARVEST FOR 3 MONTHS AFTER APPLICATION
- Feed treated with this product must not be used for animal consumption, poultry feed or mixed with animal feed.
- DO NOT mix leaves treated with this product with feed intended for animal consumption.

TRIADIMEFON 500 FUNGICIDE WP

259mm

(Printed on inner water soluble measure packs)
(This label is full size)

POISON

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING
OR USING**

**4Farmers
TRIADIMEFON 500
FUNGICIDE WP**

**ACTIVE CONSTITUENTS:
500 g/kg TRIADIMEFON**

1 Kg NET

**PACK CONTAINS: 1kg TRIADIMEFON 500 WP
WATER SOLUBLE PACKAGING. KEEP DRY.**

**NOT TO BE SOLD SEPARATELY
BEFORE USE READ ALL DIRECTIONS ON OUTER
PACK.**

**DO NOT DESTROY OUTER CONTAINER WHILE
PRODUCT STILL REMAINS**

APVMA Approval No: 50160/1/0505

