

CAUTION

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

Trilogy[®] 600

Herbicide



ADAMA

ESSENTIALS

**ACTIVE CONSTITUENT: 600 g/L TRIFLURALIN
SOLVENT: 208 g/L HYDROCARBON LIQUID**

GROUP	D	HERBICIDE
--------------	----------	------------------

A pre-emergence herbicide for the control of annual grasses and certain broadleaf weeds in certain horticultural and agricultural crops as specified in the Directions for Use.

adama.com

CONTENTS: 20 L, 110 L, 200 L, 1000 L

DIRECTIONS FOR USE

RESTRAINTS:

- DO NOT mix with dirty or turbid water.
- DO NOT store product mix for more than 24 hours.
- DO NOT incorporate more than four (4) hours after application unless otherwise specified.

FIELD CROPS

SITUATION	WEEDS CONTROLLED	STATE	RATE L/HA/SOIL TYPE	CRITICAL COMMENTS
Adzuki Beans, Cowpeas, Lablab, Mung Beans, Borlotti Beans, Red Kidney Beans Chickpeas	Amaranthus, Annual Ryegrass, Barnyard Grass, Caltrop, Crab Grass, Paradoxa Grass (Canary Grass), Pigweed, Wild Oats, Winter Grass, Wireweed Suppression of Fumitory	Qld, NSW, ACT only	960 mL-1.2 L/ha Light Soil 1.2 L/ha Medium Soil 1.36 L/ha Heavy Soil	Apply from 4 weeks up to just prior to sowing. For method of incorporation, refer to table: point # 3, 4 or 5.
	Annual Ryegrass, Paradoxa Grass (Canary Grass), Wireweed (Hogweed), Black Pigweed Suppression of Climbing Buckwheat (Black Bindweed), Wild Oats	Qld only	1L-1.36 L/ha All Soil Types	Use 1.0 L/ha when applying immediately prior to sowing. Use 1.36 L/ha when applying to dry soil before the planting rain.
	Red and White Fumitory, Rough Poppy, Wireweed, Annual Ryegrass, Barley Grass, Canary Grass, Sand Fescue, Suppression of Deadnettle, Speedwell, Three Cornered Jack, Yellow Burr Weed, Brome Grass, Cereal Oats and surface Wild Oats	SA only	1 L/ha	Apply to a level seed bed 0 to 4 weeks before sowing. For method of incorporation refer to table: point #6.
Cotton	Annual Ryegrass, Barnyard Grass, Black Pigweed (Qld only), Canary Grass, Caltrop, (Bullhead / Yellow Vine), Crab Grass, Fumitory, Mossman River Grass (Innocent Weed), Pigweed, <i>Phalaris</i> spp., Redroot (Amaranthus), Redshank (Prince of Wales Feather), Summer Grass, Wild Oats, Winter Grass, Wireweed (Hogweed) From seed only: Columbus Grass, Guinea Grass, Johnson Grass, Liverseed Grass (Urochloa)	Qld, NSW, ACT, WA only	960 mL Light Soils 1.36 L/ha Medium Soils 1.84 L/ha Heavy Soils	Spray between 6 weeks and just before sowing takes place. For method of incorporation, refer to table: point # 1 or 2.

FIELD CROPS – continued

SITUATION	WEEDS CONTROLLED	STATE	RATE L/HA/SOIL TYPE	CRITICAL COMMENTS
Faba Beans	Annual Ryegrass, Barley Grass, Capeweed, Corn Gromwell (Sheepweed), Fumitories, Geranium, Ivy Leaf Speedwell, Mustards, Turnips, Wireweed Suppression of Brome Grass, Soursob and Wild Oats	SA only	640 mL/ha + 1.0-1.5 L/ha Flowable Simazine (500 g/L)	Apply to bare moist soil and incorporate to a depth of 5 cm just prior to sowing. Incorporation should be made within 4 hours of application. Application should not be made to ridged or excessively cloddy soil. For fully reliable results significant rainfall (20-30 mm) is necessary within 2-3 weeks of application.
Lentils	Annual Phalaris, Annual Ryegrass, Wild Oats, Wireweed	NSW, ACT only	640 mL Light Soils 960 mL Medium and Heavy Soils	Apply 1 to 4 weeks before sowing.
	Fumitory – Red and White, Rough Poppy, Wireweed, Barley Grass, Canary Grass, Annual Ryegrass, Sand Fescue	SA only	1.0 L/ha All soil types	
Legume Seed Crop Establishment Annual Medics Clover Berseem, Red Strawberry (Sub and White)	Annual Ryegrass, Barnyard Grass, Black Pigweed(Qld only), Canary Grass, Caltrop (Bullhead Yellow Vine), Crab Grass, Fumitory, Mossman River Grass (Innocent Weed), <i>Phalaris</i> spp., Pigweed, Redroot (Amaranthus), Redshank (Prince of Wales Feather), Summer Grass, Wild Oats, Winter Grass, Wireweed (Hogweed)	NSW, ACT, Vic, Tas SA, WA only	960 mL Light and Medium Soils 1.36 L/ha Heavy Soils	Autumn Sowing – Apply from 4 weeks to 7 days before sowing takes place. For method of incorporation, refer to table: point # 6. Spring Sowing – Apply between 4 weeks and 3 days before sowing takes place. For method of incorporation, refer to table: point # 6. Warning: In both cases seedling disease, excessive moisture, cold weather and high salt may result in stand reduction. Temporary crop suppression could result
Lucerne (for hay and seed crop)	From seed only: Columbus Grass, Guinea Grass, Johnson Grass, Liverseed Grass (Urochloa)	All States		
Linseed		NSW, ACT, Vic, SA, WA only	960 mL Light Soils 1.2 L/ha Medium Soils 1.36 L/ha Heavy Soils	Spray 2-4 weeks before sowing. Sowing depth should be 1.3-2.5 cm. Deeper sowing may result in some stand reduction. For method of incorporation, refer to table: point # 6.
Lupins		All States		Spray between 4 weeks and just before sowing takes place. For method of incorporation, refer to table: point # 6. WA only: Use higher rate for heavier stubble coverage. Stubble coverage above 40-50% ground cover can reduce weed control below acceptable levels. Refer table: point 13 for method of incorporation.
	Annual Grasses and Broadleaf weeds	Qld, NSW, ACT, Vic only	640 mL/ha + 3.0 L/ha Flowable Simazine (500 g/L)	Use a low volume boom applying 50-100 Litres spray mixture per hectare. Apply to bare moist soil and incorporate to a depth of 5 cm just prior to sowing the crop. Incorporate within 4 hours of application. DO NOT apply to a ridged soil.
	Capeweed, Turnip, Radish, Doublegee, Suppression of Annual Ryegrass, Wild Oats	WA only	1 L/ha + 1.0-1.5 L/ha Flowable Simazine (500 g/L)	Rate for yellow sands. Refer to incorporation table point #11, 12 or 13
	As above plus Suppression of Brome Grass		1 L/ha + 2.0 L/ha Flowable Simazine (500 g/L)	Rate for all other soil types. Apply to bare moist soil and incorporate to a depth of 5 cm just prior to sowing. Incorporation should be made within 4 hours of application. Application should not be made to ridged or excessively cloddy soil. For Simazine to be effective, sufficient rainfall (20-30 mm) to wet the soil through the weed root zone is necessary, within 2-3 weeks of application. Results with Simazine can be variable if seasonal conditions are dry prior to sowing and Lupins are sown into dry or low moisture seed beds.

FIELD CROPS – continued

SITUATION	WEEDS CONTROLLED	STATE	RATE L/HA/SOIL TYPE	CRITICAL COMMENTS
Lupins – <i>continued</i>	Red and White Fumitory, Rough Poppy, Wireweed, Barley Grass, Canary Grass, Annual Ryegrass, Sand Fescue, suppression of Deadnettle, Speedwell, Three Cornered Jack, Yellow Burr Weed, Brome Grass, Cereal Oats, soil surface Wild Oats	SA only	1-1.36 L/ha All soil types	Spray between 4 weeks and just before sowing takes place. Refer to the incorporation table point #6 for method of incorporation.
	Above weeds plus Capeweed, Common Fumitory, Geranium, Indian Hedge Mustard, Sheepweed, Shepherd Purse, Toadrush, Turnips, suppression of Ice Plant and Soursob		1-1.36 L/ha + 1.7-2.4 L/ha of Flowable Simazine (500 g/L)	Use a low volume boom applying 50-100 Litres spray mixture per hectare. Apply to bare moist soil and incorporate to a depth of 5 cm just prior to sowing the crop. Incorporate within 4 hours of application. DO NOT apply to ridged soil.
Pigeon Peas	Amaranthus, Barnyard Grass, Canary Grass, Crowsfoot Grass, Pigweed, Spiny Burrgrass, Summer Grass, Wild Oats, Wireweed (Hogweed), Suppression of Yellow Vine (Caltrop) From seed only: Columbus Grass, Guinea Grass, Johnson Grass, Liverseed Grass (Urochloa)	NSW, ACT only	960 mL Light Soils 1.2 L/ha Medium Soils 1.36 L/ha Heavy Soils	Apply between 4 weeks and just before sowing. Refer to incorporation table points #3, 4 or 6 for suitable method of incorporation.
Tobacco	Summer Grass, Crowsfoot Grass, Red Natal Grass, Love Grass, Button Grass, Rhodes Grass, Pigweed	Qld only	640 mL for light soils 960 mL for medium soils	Apply to soil 3 to 4 weeks prior to transplanting. The longer period to be used for applications made during June and July. Incorporate to a depth of 10 cm.
	Crowsfoot Grass	NSW, ACT only	640 mL for light soils 960 mL for medium and heavy soils	Apply to light sandy soil 14 to 21 days before transplanting. DO NOT incorporate to a depth greater than 6 cm. Apply to loam (medium soil) 14 to 21 days before transplanting. DO NOT incorporate to a depth of greater than 6 cm.
Peanuts	Annual Ryegrass, Barnyard Grass, Black Pigweed (Qld only), Canary Grass, Caltrop, (Bullhead Yellow Vine), Crab Grass, Mossman River Grass (Innocent Weed), Pigweed, Redroot (Amaranthus), Redshank (Prince of Wales Feather), Summer Grass, Wild Oats, Winter Grass, Wireweed (Hogweed)	Qld, WA only	960 mL Light Soils 1.2 L/ha Medium Soils 1.36 L/ha Heavy Soils	Spray between 4 weeks and just before sowing takes place. For method of incorporation, refer to table: point # 3, 4 or 5 for method of incorporation.
Peas Canola (Rapeseed) Safflower	From seed only: Columbus Grass, Guinea Grass, Johnson Grass Liverseed Grass (Urochloa)	All States		Spray between 4 weeks and just before sowing takes place. For method of incorporation, refer to table: point # 6 or 11 for method of incorporation.
Sugarcane – Early Season Late Season		Qld, NSW, ACT only	2.4 L/ha All soil types	Apply to plant cane after emergence to “out of hand” stage. Apply to Ratoon cane immediately after harvest.
			1.84 L/ha All soil types	For method of incorporation, refer to table: point # 3, 7 or 10.
Sunflowers		All States	960 mL Light Soils 1.2 L/ha Medium Soils 1.36 L/ha Heavy Soils	Spray between 4 weeks and just before sowing takes place. For method of incorporation, refer to table: point # 3, 4 or 5.
Wheat, Barley, Triticale Pre-Sowing ONLY Pre-Sowing & Post- Sowing (Self- Mulching Soils)	Annual Ryegrass, <i>Phalaris</i> spp., Wireweed (Hogweed)	NSW, ACT, Vic, WA only	640 mL/ha All soil types	Apply 1-4 weeks before sowing. Sowing depth should be at least 5 cm. Use cover harrows behind combine. Ground should be left flat. DO NOT use pre- sowing on self-mulching soils as damage may occur from wheel tracking and poor control of wild oats.
	Fumitory	WA only		For method of incorporation, refer to table: point # 6 for method of incorporation.
	Canary Grass	Vic only		
	As above except Fumitory			Pre-Sowing – Apply more than 4 weeks before sowing to prevent crop damage. Post-Sowing – Apply within 2 days after sowing to well prepared seedbed. For method of incorporation, refer to table: point # 6.

FIELD CROPS – continued

SITUATION	WEEDS CONTROLLED	STATE	RATE L/HA/SOIL TYPE	CRITICAL COMMENTS
Wheat	Annual Ryegrass, Canary Grass (Paradoxa Grass), soil surface Wild Oats, Wireweed (Hogweed)	Qld only	640 mL/ha All soil types	On non-self mulching soils apply 1-4 weeks before sowing. Sowing depth should be at least 5 cm. Use cover harrows behind combine. Ground should be left flat. On self-mulching soils, as above except apply more than 4 weeks before sowing to prevent crop damage. For method of incorporation, refer to table: point # 6.
Barley				Apply to self-mulching and non-self-mulching from 1-4 weeks before sowing. Sowing depth should be at least 5 cm. Use cover harrows behind combine. Ground should be left flat. For method of incorporation, refer to table: point # 6.
Wheat, Triticale, Rye	Annual Ryegrass, Red and White Fumitory, <i>Phalaris</i> spp., Wireweed, suppression of Deadnettlles, Rough Poppy, Yellow Burr Weed	SA only	1 L/ha for all soil types	Apply 1-4 weeks before sowing. Sowing depth should be at least 5 cm. Use cover harrows behind combine. Ground should be left flat. DO NOT use pre-sowing on self mulching soils as damage may occur from wheel tracking and poor control of wild oats. Refer incorporation table: point #6 for method of incorporation.
Barley	As above for SA plus Sand Fescue and suppression of Brome Grass			
Wheat and Triticale Only	Annual Phalaris	NSW, ACT only	80 mL/ha plus + 20 g/ha Chlorsulfuron (750 g/kg)	If possible spray and incorporate into the soil in one operation. If this is not possible incorporation should take place within 4 hours of spraying. Delay may cause inferior weed control.

VEGETABLES, ORCHARDS AND VINEYARDS

SITUATION	WEEDS CONTROLLED	STATE	RATE L/HA/SOIL TYPE	CRITICAL COMMENTS
Transplants only Broccoli, Cauliflowers, Cabbage, Tomatoes	Annual Ryegrass, Barnyard Grass, Black Pigweed, Canary Grass, Caltrop (Bullhead Yellow Vine), Crab Grass, Mossman River Grass (Innocent Weed), Pigweed, Redroot (Amaranthus), Redshank (Prince of Wales Feather), Summer Grass, Wild Oats, Winter Grass, Wireweed (Hogweed) From seed only: Columbus Grass, Guinea Grass, Johnson Grass, Liverseed Grass (Urochloa)	All States	960 mL Light Soils 1.36 L/ha Medium Soils 1.84 L/ha Heavy Soils	Spray between 4 weeks and just before sowing takes place. For method of incorporation, refer to table: point # 3, 4 or 5.
Direct seeded Broccoli, Brussels Sprouts, Cabbage				
Direct Seeded Cauliflower				
Carrots				
Chicory				
Green Beans, Navy Beans				
Soy Beans			960 mL Light Soils 1.2 L/ha Medium Soils 1.36 L/ha Heavy Soils	
			960 mL/ha Light Soils 1.36 L/ha Medium Soils 1.84 L/ha Heavy Soils	

VEGETABLES, ORCHARDS AND VINEYARDS – continued

SITUATION	WEEDS CONTROLLED	STATE	RATE L/HA/SOIL TYPE	CRITICAL COMMENTS
Orchards and Vineyards	Annual Ryegrass, Barnyard Grass, Black Pigweed, Canary Grass, Caltrop (Bullhead Yellow Vine), Crab Grass, Mossman River Grass (Innocent Weed), Pigweed, Redroot (Amaranthus), Redshank (Prince of Wales Feather), Summer Grass, Wild Oats, Winter Grass, Wireweed (Hogweed) From seed only: Columbus Grass, Guinea Grass, Johnson Grass, Liverseed Grass (Urochloa)	Qld, SA, WA, Vic, Tas only	960 mL Light Soils 1.36 L/ha Medium Soils 1.84 L/ha Heavy Soils	Apply to new planting during pre-plant cultivation. Apply to established crops in spring after weeds and green manure crop has been ploughed into ground. For method of incorporation, refer to table: point # 8 or 9.
Vetch	Annual Ryegrass, Deadnettle, Wireweed, soil surface Wild Oats, suppression of Brome Grass, Rough Poppy, Speedwell, Three Cornered Jack, Burr Weed, Sheepweed	SA, WA only	1.36 L/ha	Apply to level seedbed 0 to 4 weeks before sowing. Refer to incorporation table: point # 6 for method of incorporation.

FIELD CROPS: FOR USE IN NO-TILL; MIN-TILL CROPPING SYSTEMS OR INCORPORATED BY SOWING (IBS) (SEE TABLE 13)

SITUATION	WEEDS CONTROLLED	STATE	RATE L/HA/SOIL TYPE	CRITICAL COMMENTS
Wheat, Barley, Triticale	Annual Ryegrass, <i>Phalaris</i> spp., Wireweed, Fumitory	NSW, ACT, Vic, SA, WA, only	1.2-1.6 L/ha	Use the higher rate on lighter sandy loam soils. DO NOT use on heavy soils. Use with Knife/Blade point sowing equipment. Use the higher rate for heavier stubble coverage. Stubble coverage above 40-50% ground cover can reduce weed control below acceptable levels. Refer to Point 13 in the Incorporation Table. WA only: Application can occur 0-24 hours prior to incorporation by sowing. For best results apply as close as possible to sowing (within 12 hours). Application 12-24 hours before sowing may be more adversely affected by above average soil moisture, warm temperatures and high weed seed densities. These factors individually or combined may reduce final weed control levels.
Chickpeas	Annual Ryegrass, Wireweed, <i>Phalaris</i> spp., Fumitory	WA only	1 to 1.36 L/ha + 2 L/ha Flowable Simazine (500 g/L)	Incorporate as per point #13 in the incorporation table.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED

GENERAL INSTRUCTIONS

This product must be incorporated into the soil within 4 hours of application except in Western Australia where the crop is sown with minimum tillage sowing equipment (fitted with knife points or blades less than 12 mm wide, usually with press wheels) where application may occur up to 24 hours before incorporation by the sowing process. Emerging roots of susceptible species take up Trilogy 600 Herbicide as seedlings germinate. Selectivity is improved by sowing below the herbicide band, and impeded by poor growing conditions and compacted soil.

RESISTANT WEEDS WARNING:

Trilogy® 600 Herbicide is a member of the Dinitroanilines group of herbicides.

GROUP D	HERBICIDE
----------------	------------------

Trilogy 600 Herbicide has the Inhibitors of Tubulin Formation mode of action. For weed resistance management Trilogy 600 Herbicide is a Group D herbicide. Some naturally occurring weed biotypes resistant to Trilogy 600 Herbicide and other Group D herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Trilogy 600 Herbicide or other Group D herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Adama Australia Pty. Ltd. accepts no liability for any losses that may result from the failure of Trilogy 600 Herbicide to control resistant weeds.

METHOD OF INCORPORATION TABLE

- Prior to furrowing out: 2 workings at an angle required using Offset OR Tandem disc harrows.
- After furrowing out: 2 workings required using Go-Devil discs or Lilliston cultivators set at 10 cm depth.
- Rotary Hoe: 1 working required at 5-7.5 cm depth.
Sugarcane: 7.5-13 cm depth.

- Offset or Tandem Disc Harrows: (Preferably with spiked harrows in tandem) 2 workings at an angle required at 7.5-15 cm depth at 6.5-10 km per hour.
- Heavy Diamond or Stump Jump Harrows: (Weighted 20-30 kg per section) at 10-13 km/hr speed. Then cross work with offset or tandem disc harrows set to 7.3-15 cm depth at speed 6.5-10 km per hour.
- Weighted Heavy Diamond OR Stump Jump Harrows: (Weight with 20-30 kg per section) at 10-13 km per hour. Cross work with combine at 5-7.5 cm depth at speed of 10-13 km per hour.
- Disc Ratoon Cultivator: 2 workings needed with discs and cultivator set at 7.5-13 cm depth.
- Offset or Tandem Disc Harrows: Set at 7.5-15 cm depth. A second discing is required working in opposite direction with discs set to throw treated soil into tree or vine row.
- Rotary Hoe: 1 working needed at 5-10 cm depth.
- Offset Discs (Bumpers): 2 workings needed at depth 7.5-13 cm.
- Incorporation by Sowing (IBS) on suitably prepared seedbed with heavy diamond harrows trailing or as a separate operation.
- Post-sowing pre-emergence: Use heavy diamond harrows cross working at right angles to the direction of sowing. Do not attempt this method of incorporation on poorly prepared, clumpy or cloddy soils.
- Incorporated by sowing (IBS) with knife or blade points. Use press wheels to avoid dragging treated soil back into the furrow. Maintain a slow to moderate speed, so that soil is not thrown into adjacent furrows. **NOTE:** Knife or blade point systems can result in poor weed control in the seed furrow, as chemical can be displaced from this zone. Stubble coverage above 40-50% ground cover can reduce weed control below acceptable levels. A knife or blade point is 12 mm or less, no wings, inverted T or blade and generally at a min of 8 inch spacing. Heavy rain between sowing and crop emergence that washes soil into the furrow will reduce emergence.

COMPATIBILITY

Trilogy 600 Herbicide is compatible with a range of herbicides including Farnoz Tackle® WG Herbicide, Simazine and Diuron 500 Flowable.

MIXING AND APPLICATION

This product is an emulsifiable concentrate, which mixes readily with water. Add the recommended amount to the spray tank during filling operation and apply 70-450 L of water per hectare (Broadacre basis) dependent on soil type. Under hot conditions or if possible spray and incorporate into the soil in one operation. Delay may cause inferior weed control. Use properly calibrated standard low pressure 170-340 kilopascal boom type sprayer with flat fan tips.

CONDITIONS FOR BEST RESULTS

This product must be thoroughly incorporated as recommended. Soil should be well-worked and free of weeds at time of application. Product effectiveness may be reduced by inadequate incorporation, high organic matter, excess clods, crop or trash residues, stones or other foreign matter and in areas of unnaturally high weed seed population such as header tracks or livestock rest areas. Trifluralin is volatile and disappears from exposed surfaces. High temperatures, winds or warm moist soil hastens loss.

WILD OATS

Germinating Wild Oats lying on the soil surface will not be controlled. Hence, specific Wild Oat control is only possible with shallow cultivation. Poor control will occur on self-mulching soils and all soil types where deep cultivation is practiced.

PROTECTION OF CROPS, LIVESTOCK, WILDLIFE, FISH, ENVIRONMENT AND OTHERS

DO NOT use in high winds.

DO NOT exceed rates specified, to avoid crop damage.

DO NOT plant sensitive grasses such as oats, sorghum, millets, *Phalaris* spp., ryegrass, or wheat for 12 months following the use of this product except where wheat follows wheat or other winter crops.

DO NOT plant oil seed poppies for two years following use.

DO NOT apply to orchards and vineyards after first flush of growth or when residues can lodge on or in fruit.

DO NOT allow spray drift onto susceptible crops or ornamentals.

DO NOT contaminate streams, rivers, or waterways with chemical or used containers.

Reduced germination of wheat and barley may occur due to the following reasons with use of this product:

- Short coleoptile cultivars

- Use of seed dressings

- Shallow or uneven seeding depth

Drift Warning: DO NOT apply under meteorological conditions or from spray equipment which allow spray drift onto susceptible crops or pastures.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, well-ventilated area, as cool as possible. DO NOT store for prolonged periods in direct sunlight. DO NOT store below 5°C. Extended storage below 5°C can result in the formation of crystals on the bottom of the container. If crystallisation does occur, store the container on its side at room temperature and rock occasionally until the crystals redissolve. Ensure crystals are dissolved before adding to spray tank.

Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilizers.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulation. DO NOT burn empty containers or product.

Envirodrum Micro Matic Valve (110 L)

Store the original sealed Envirodrum in a cool well-ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the Micro Matic valve or the security seal. DO NOT contaminate the Envirodrum with water or any foreign matter. After each use of the product, please ensure that the Micro Matic coupler delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the Envirodrum to the point of purchase. The Envirodrum remains the property of Adama Australia Pty. Ltd.

1000 L

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Storage must be secure so that contents cannot be tampered with. All locks and/or seals must be in order. If locks or seals are broken prior to initial use then the integrity of this product cannot be assured. If this occurs Adama Australia Pty. Ltd. should be advised immediately. This minibulk container is reusable and remains the property of Adama Australia Pty. Ltd. DO NOT rinse empty container. Empty contents fully into application equipment. Close all valves and return to the point of supply for refill or storage. No other liquid, solid or pesticide product should be put into it. When empty return to Adama Australia Pty. Ltd. for cleaning, relabelling and refilling.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Repeated exposure may cause allergic disorders. Sensitive workers should use protective clothing. Avoid contact with eyes and skin. DO NOT inhale spray mist. When opening the container and using the prepared spray wear cotton overalls, buttoned to the neck and wrist and a washable hat and elbow length PVC gloves and face shield or goggles. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If swallowed, DO NOT induce vomiting. Give a glass of water.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for Trilogy 600 Herbicide is available from Adama on request. Call Customer Service on (02) 9431 7800.

CONDITIONS OF SALE: The use of Trilogy® 600 Herbicide being beyond the control of the manufacturer, no warranty expressed or implied is given by Adama Australia Pty. Ltd., regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Adama Australia Pty. Ltd. accepts no responsibility for any consequence whatsoever resulting from the use of this product.

® Registered trademark of an ADAMA Agricultural Solutions Ltd Company

NOT A DANGEROUS GOOD ACCORDING TO THE AUSTRALIAN DANGEROUS GOODS (ADG) CODE.

Adama Australia Pty. Ltd. ABN 55 050 328 973

Suite 1, Level 4, Building B,

207 Pacific Highway

St Leonards NSW 2065 Australia

Tel: (02) 9431 7800 Fax: (02) 9431 7700

APVMA Number: 68897/59332

Batch No.

Date of Manufacture