

## STORAGE AND DISPOSAL

### Keep out of reach of children

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to a recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots in compliance with relevant Local, State or Territory government regulations. Do not burn empty containers or product.

### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Telephone Australia 13 11 26.

### MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet (MSDS) that can be obtained from the supplier.

## MANUFACTURERS WARRANTY AND EXCLUSION OF LIABILITY

Chemtura Australia Pty Ltd has no control over storage, handling and manner of use of this product. Where this material is not stored, handled or used correctly and in accordance with directions, no express or implied representations or warranties concerning this product (other than non-excludable statutory warranties) will apply. Chemtura Australia Pty Ltd accepts no responsibility for any loss arising from incorrect storage, handling or use.

In a Transport Emergency  
Dial **000**  
Police or Fire Brigade



Batch No.  
DOM

APVMA Approval No.: 66746/54314

File name	xxxxx xxxxx 1kg B MPL V4.pdf
Dimensions	Size 75mm H x 86mm W
Scale	Prints at 100% A4
Date	08/11/2011

**CAUTION**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

*Chemtura*  
**Triasul**  
750 WG HERBICIDE

**ACTIVE CONSTITUENT: 750 g/kg TRIASULFURON**

**GROUP B HERBICIDE**

For pre-plant control of annual ryegrass, Paradoxa grass and certain broadleaf weeds in wheat and for post-emergent control of wild radish in wheat, oats and barley as per Directions for Use

**IMPORTANT: READ THIS BOOKLET BEFORE USING THIS PRODUCT**

**Contents: 1 kg**

Chemtura Australia Pty Ltd  
Level 7, 435 King William St, Adelaide, SA, 5000  
Tel: (08) 8112 0900 Fax: (08) 8112 0999



## TABLE OF CONTENTS

Directions for Use – Western Australia .....	4
Directions for Use – South Australia .....	8
Directions for Use – New South Wales .....	12
Directions for Use – Queensland .....	16
Directions for Use – Victoria and Tasmania .....	20
General Instructions .....	24
Resistant Weeds Warning .....	24
Mixing .....	24
Application .....	25
Sprayer Cleanup .....	25
Compatibility .....	25
Crop Rotation Guidelines .....	26
Precaution .....	29
Re-entry Period .....	29
Protection of Crops .....	29
Protection of Wildlife .....	30
Storage and Disposal .....	30
First Aid .....	30
Material Safety Data Sheet .....	30
Manufacturer's Warranty .....	30

**DIRECTIONS FOR USE****WESTERN AUSTRALIA ONLY****PRE-EMERGENT APPLICATION RESTRAINTS:**

DO NOT apply to crops undersown with legumes.

DO NOT use if another Group B herbicide (ALS Inhibitor) has been used during the current season.

DO NOT apply to paddocks where there is a high risk of weeds resistant to Group B Herbicides.

Apply no more than two Group B Herbicides in any four-year period on the same paddock.

<b>CROP</b>	<b>WEEDS CONTROLLED</b>	<b>RATE/HA</b>
<b>Wheat</b>	Burr Medic, Common Cotula, Corn Gromwell, (White Ironweed, Sheepweed), Deadnettle, Denseflower Fumitory, Hedge Mustard, Indian Hedge Mustard, Matricaria, Paterson's Curse, Rough Poppy, Smallflower Fumitory, Wards Weed, wild Turnip, Yellow Burrweed (Amsinckia), and suppression of Crassula.	30 g
	Annual Ryegrass, Capeweed, Doublegee or Three Cornered Jack (Spiny Emex), wireweed and suppression of Wild Radish and also of 60–80% suppression of Soursob.	35 g
	Annual Ryegrass, Corn Gromwell, (White Ironweed, Sheepweed), Deadnettle, Hedge Mustard, Indian Hedge Mustard, Prickly Lettuce (Whipthistle), Smallflower Fumitory, Wild Turnip, Wireweed (Hogweed), Ward's Weed	10 g to 15 g plus 830 mL of trifluralin (480 g/L)

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL  
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**CRITICAL COMMENTS**

Apply to bare moist soil prior to sowing or at sowing and incorporate by the sowing operation using low profile 10 cm combine points. Application should not be made to ridged or excessively cloddy soil. For best results apply to moist soil, when follow up rain is likely to occur within 7 – 10 days. In conservation tillage situations where weeds and grasses have emerged apply as a tank mixture with the recommended rate of knockdown herbicide prior to sowing.

When used on Kulin wheat on very acid soils (pH 5.5 in water) or under poor fertility conditions increases stem breakage may occur. Early season crop retardation may occur where the product is used on soils with a pH greater than 8, and which are prone to zinc deficiency.

This mixture to be only used on alkaline soils where the pH is greater than 8 (1:5 soil: water suspension method). For best results apply mix to bare moist soil that has a minimum of trash and incorporate to a depth of 5 cm just prior to sowing. Incorporation should be made within 4 hours of application. Heavy rainfall (greater than 50 mm) within 7 days of application may affect efficacy especially at the 10 g rate. Late germination of some weeds e.g. Annual Ryegrass will not be controlled in seasons of above average rainfall. Use the higher rate where high density of weeds is expected.

**WITHHOLDING PERIODS**

**PRE-EMERGENT: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 WEEKS AFTER APPLICATION**

**POST-EMERGENT: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 14 DAYS AFTER APPLICATION**

**WESTERN AUSTRALIA ONLY POST-EMERGENT APPLICATION RESTRAINTS:**

DO NOT apply to crops undersown with legumes. DO NOT spray when very dry conditions prevail. DO NOT spray under dry frosty conditions. DO NOT use if another Group B herbicide (ALS Inhibitor) has been used during the current season. DO NOT apply to paddocks where there is a high risk of weeds resistant to Group B Herbicides. Apply no more than two Group B Herbicides in any four-year period on the same paddock. If RAIN FALLS within 6 hours of application, the effect could be diminished.

<b>CROP</b>	<b>WEEDS CONTROLLED</b>	<b>RATE/HA</b>
<b>Wheat, Oats, Barley</b>	Wild Radish	10–15 g plus crop oil at 1 L per 100 L of spray mixture or surfactant at recommended label rates.
<b>Barley, Oats, Triticale, wheat</b> From 3 leaf to early tillering stage	Turnip Weed	6.5 g + 300 mL terbutryn (500 g/L)
	Denseflower Fumitory, Hedge Mustard, Smallflower Fumitory	6.5 g + 300 mL terbutryn (500 g/L) to 10 g + 440 mL terbutryn (500 g/L)
	Ball Mustard, Field peas (volunteer)	10 g + 440 mL terbutryn (500 g/L)
	Deadnettle	10 g + 440 mL terbutryn (500 g/L) to 13 g + 600 mL terbutryn (500 g/L)
	Australian Crassula, Doublegee or Three cornered Jack (Spiny Emex), Volunteer Lupins.	13 g + 600 mL terbutryn (500 g/L)

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**CRITICAL COMMENTS**

**EARLY POST EMERGENT APPLICATION:** Spray prior to the crop reaching mid-tillering (Zadoks 23) and when Wild Radish is in the 2–6 leaf stage. Use rates towards the lower end of the range, when weeds are small and growing conditions ideal. Spray only when weeds are actively growing.

**LATE POST EMERGENT APPLICATION:** Spray during early flowering of the wild radish. DO NOT apply to the crop during crop anthesis or flowering (Zadoks 60–69). DO NOT spray on weeds under stress.

**WARNING:** Application to oats may cause some yellowing. DO NOT use on weeds resistant, or thought to be resistant, to Group B herbicides. Petroleum based crop oils are recommended as well as non-ionic surfactants (1000 g/L).

Spray when weeds are in 2 to 6 leaf stage (up to 6 node/leaflet for field peas, 10 leaflet for vetch), except 2 to 4 leaf for Doublegee. Use the rates towards the lower end of the range (where applicable) when weeds are small and soil conditions are very moist.

Spray only when weeds are actively growing.

Spray only after good rain and when topsoil is moist. Best results are obtained when good soil moisture has been present since planting.

**WITHHOLDING PERIODS**

**PRE-EMERGENT: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 WEEKS AFTER APPLICATION**

**POST-EMERGENT: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 14 DAYS AFTER APPLICATION**

**SOUTH AUSTRALIA ONLY****PRE-EMERGENT APPLICATION RESTRAINTS:**

DO NOT apply to crops undersown with legumes.

DO NOT use if another Group B herbicide (ALS Inhibitor) has been used during the current season.

DO NOT apply to paddocks where there is a high risk of weeds resistant to Group B Herbicides.

Apply no more than two Group B Herbicides in any four-year period on the same paddock.

<b>CROP</b>	<b>WEEDS CONTROLLED</b>	<b>RATE/HA</b>
<b>Wheat</b>	Ball Mustard, Burr Medic, Corn Gromwell (White Ironweed, Sheepweed), Deadnettle, Denseflower Furnitory, Hedge Mustard, Indian Hedge Mustard, Prickly Lettuce, Smallflower Furnitory, Stemless Thistle, Vetch, Wild Turnip, Yellow Burrweed (Amsinckia), Rough Poppy, Wards Weed.	30 g
	Annual Ryegrass, Capeweed, Three cornered Jack or Doublegee (Spiny Emex), Volunteer Chickpeas, Faba Beans and Field Peas, Wireweed, and suppression of Wild Radish, also 60–80% control of Soursob and suppression of remaining plants. Suppression of Skeleton Weed.	35 g
	Annual Ryegrass, Corn Gromwell, (White Ironweed, Sheepweed), Deadnettle, Hedge Mustard, Indian Hedge Mustard, Prickly Lettuce (Whipthistle), Smallflower Furnitory, Wild Turnip, Wireweed (Hogweed), Ward's Weed	10 g to 15 g plus 830 mL of trifluralin (480 g/L)

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL  
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**



**CRITICAL COMMENTS**

Apply to bare moist soil prior to sowing or at sowing and incorporate by the sowing operation using low profile 10 cm combine points. Application should not be made to ridged or excessively cloddy soil. For best results apply to moist soil, when follow up rain is likely to occur within 7–10 days. In conservation tillage situations where weeds and grasses have emerged apply as a tank mixture with the recommended rate of knockdown herbicide prior to sowing.

Chemtura Triasul will provide good control of volunteer grain legumes, however a small proportion of plants may survive and require an overspray to eliminate the potential for grain contamination.

For Skeleton Weed a significant degree of control will be achieved on soil types of a predominantly sandy clay loam mixture with a pH greater than 8. Best control is observed where Skeleton Weed germinates in the very early stages of the crop. Surviving plants will be stunted.

**DO NOT** use on weeds resistant, or thought to be resistant, to Group B Herbicides.

This mixture to be only used on alkaline soils where the pH is greater than 8 (1:5 soil: water suspension method). For best results apply mix to bare moist soil that has a minimum of trash and incorporate to a depth of 5 cm just prior to sowing. Incorporation should be made within 4 hours of application. Heavy rainfall (greater than 50 mm) within 7 days of application may affect efficacy especially at the 10 g rate. Late germination of some weeds e.g. Annual Ryegrass will not be controlled in seasons of above average rainfall. Use the higher rate where high density of weeds is expected.

**WITHHOLDING PERIODS**

**PRE-EMERGENT: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 WEEKS AFTER APPLICATION**

**POST-EMERGENT: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 14 DAYS AFTER APPLICATION**

**SOUTH AUSTRALIA ONLY****POST-EMERGENT APPLICATION RESTRAINTS:**

DO NOT apply to crops undersown with legumes. DO NOT spray when very dry conditions prevail. DO NOT spray under dry frosty conditions. DO NOT use if another Group B herbicide (ALS Inhibitor) has been used during the current season. DO NOT apply to paddocks where there is a high risk of weeds resistant to Group B Herbicides. Apply no more than two Group B Herbicides in any four-year period on the same paddock. If RAIN FALLS within 6 hours of application, the effect could be diminished.

<b>CROP</b>	<b>WEEDS CONTROLLED</b>	<b>RATE/HA</b>
<b>Wheat, Oats, Barley</b> Late Post emergent application	Wild Radish	15 g plus crop oil at 1 L per 100 L of spray mixture or surfactant at recommended label rates.
<b>Barley, Oats, Triticale, Wheat</b> From 3 leaf to early tillering stage	Faba Beans (volunteer), Long Fruited Turnip, Long Headed Poppy, Paterson's Curse, Turnip Weed	6.5 g + 300 mL terbutryn (500 g/L)
	Hedge Mustard, Indian Hedge Mustard, Wild Turnip	6.5 g + 300 mL terbutryn (500 g/L) to 10 g + 440 mL terbutryn (500 g/L)
	Ball Mustard, Crassula, Medic, Prickly Lettuce (Whipthistle)	10 g + 440 mL terbutryn (500 g/L)
	Field peas (volunteer), Wild Radish	10 g + 440 mL terbutryn (500 g/L) to 13 g + 600 mL terbutryn (500 g/L)
	Volunteer Lupins, Doublegee or Three Cornered Jack (Spiny Emex), Vetch, Wireweed (less than 3 leaves) suppression.	13 g + 600 mL terbutryn (500 g/L)

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

### **CRITICAL COMMENTS**

**LATE POST EMERGENT APPLICATION:** Spray during early flowering of the wild radish. DO NOT apply to the crop during or after anthesis and flowering (Zadoks 60). DO NOT spray on weeds under stress.

**WARNING:** Application to oats may cause some yellowing.

DO NOT use on weeds resistant, or thought to be resistant, to Group B herbicides.

Petroleum based crop oils are recommended as well as non-ionic surfactants (1000 g/L).

Spray when weeds are in 2 to 6 leaf stage (up to 6 node/leaflet for field peas, 10 leaflet for vetch), except 2 to 4 leaf for Doublegee.

Use the rates towards the lower end of the range (where applicable) when weeds are small and soil conditions are very moist.

Spray only when weeds are actively growing.

Spray only after good rain and when topsoil is moist. Best results are obtained when good soil moisture has been present since planting.

### **WITHHOLDING PERIODS**

**PRE-EMERGENT: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 WEEKS AFTER APPLICATION**

**POST-EMERGENT: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 14 DAYS AFTER APPLICATION**

**NEW SOUTH WALES ONLY****PRE-EMERGENT APPLICATION RESTRAINTS:**

DO NOT apply to crops undersown with legumes.

DO NOT use if another Group B herbicide (ALS Inhibitor) has been used during the current season.

DO NOT apply to paddocks where there is a high risk of weeds resistant to Group B Herbicides.

Apply no more than two Group B Herbicides in any four-year period on the same paddock.

<b>CROP</b>	<b>WEEDS CONTROLLED</b>	<b>RATE/HA</b>
<b>Wheat</b>	Black Bindweed (Climbing Buckwheat), Burr Medic, Common Cotula, Corn Gromwell, (White Ironweed, Sheepweed), Deadnettle, Denseflower Fumitory, Hedge Mustard, Indian Hedge Mustard, Lesser Swinecress, Paterson's Curse, Prickly Lettuce, Rough Poppy, Smallflower Fumitory, Sowthistle, Stagger Weed, Turnip Weed, Wild Turnip, Yellow Burrweed (Amsinckia), and suppression of Variegated Thistle.	30 g
	Annual Ryegrass, Capeweed, Doublegee or Three Cornered Jack (Spiny Emex), Paradoxa Grass (Annual Phalaris), Wireweed and suppression of Wild Radish and Mexican Poppy	35 g
	Annual Ryegrass, Corn Gromwell, (White Ironweed, Sheepweed), Deadnettle, Hedge Mustard, Indian Hedge Mustard, Prickly Lettuce (Whipthistle), Smallflower Fumitory, Wild Turnip, Wireweed (Hogweed), Ward's Weed	10 g to 15 g plus 830 mL of trifluralin (480 g/L)

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL  
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**CRITICAL COMMENTS**

Apply to bare moist soil prior to sowing or at sowing and incorporate by the sowing operation using low profile 10 cm combine points.

Application should not be made to ridged or excessively cloddy soil.

For best results apply to moist soil, when follow up rain is likely to occur within 7–10 days.

In conservation tillage situations where weeds and grasses have emerged apply as a tank mixture with the recommended rate of knockdown herbicide prior to sowing.

For best results, for Paradoxa Grass control, apply to dry soil before the sowing rain.

DO NOT use on weeds resistant, or thought to be resistant, to Group B Herbicides.

This mixture to be only used on alkaline soils where the pH is greater than 8 (1:5 soil: water suspension method). For best results apply mix to bare moist soil that has a minimum of trash and incorporate to a depth of 5 cm just prior to sowing. Incorporation should be made within 4 hours of application. Heavy rainfall (greater than 50 mm) within 7 days of application may affect efficacy especially at the 10 g rate. Late germination of some weeds e.g. Annual Ryegrass will not be controlled in seasons of above average rainfall. Use the higher rate where high density of weeds is expected.

**WITHHOLDING PERIODS**

**PRE-EMERGENT: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 WEEKS AFTER APPLICATION**

**POST-EMERGENT: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 14 DAYS AFTER APPLICATION**

## NEW SOUTH WALES ONLY

### POST-EMERGENT APPLICATION RESTRAINTS:

DO NOT apply to crops undersown with legumes.

DO NOT spray when very dry conditions prevail.

DO NOT spray under dry frosty conditions.

DO NOT use if another Group B herbicide (ALS Inhibitor) has been used during the current season.

If RAIN FALLS within 6 hours of application, the effect could be diminished.

DO NOT apply to paddocks where there is a high risk of weeds resistant to Group B Herbicides.

Apply no more than two Group B Herbicides in any four-year period on the same paddock.

CROP	WEEDS CONTROLLED	RATE/HA
<b>Wheat, Oats, Barley</b> Late Post emergent application	Wild Radish	15 g plus crop oil at 1 L per 100 L of spray mixture or surfactant at recommended label rates.
<b>Barley, Oats, Triticale, wheat</b> From 3 leaf to early tillering stage	Turnip Weed	6.5 g + 300 mL terbutryn (500 g/L)
	Hedge Mustard, Indian Hedge Mustard, London Rocket, Wild Turnip	6.5 g + 300 mL terbutryn (500 g/L) to 10 g + 440 mL terbutryn (500 g/L)
	Black Bindweed (Climbing Buckwheat), Faba Beans (volunteer), Field peas (volunteer), Medic, Shepherds Purse	10 g + 440 mL terbutryn (500 g/L)
	Deadnettle, Mexican Poppy, Wild Radish	10 g + 440 mL terbutryn (500 g/L) to 13 g + 600 mL terbutryn (500 g/L)
	Coreopsis, Sunflower (volunteer), Vetch	13 g + 600 mL terbutryn (500 g/L)

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**CRITICAL COMMENTS**

**LATE POST EMERGENT APPLICATION:** Spray during early flowering of the wild radish. DO NOT apply to the crop during crop anthesis or flowering (Zadoks 60–69). DO NOT spray on weeds under stress.

**WARNING:** Application to oats may cause some yellowing.

DO NOT use on weeds resistant, or thought to be resistant, to Group B Herbicides.

Petroleum based crop oils are recommended as well as non-ionic surfactants (1000 g/L).

Spray when weeds are in 2 to 6 leaf stage (up to 6 node/leaflet for field peas, 10 leaflet for vetch), except 2 to 4 leaf for Doublegee.

Use the rates towards the lower end of the range (where applicable) when weeds are small and soil conditions are very moist.

Spray only when weeds are actively growing.

Spray only after good rain and when top soil is moist.

Best results are obtained when good soil moisture has been present since planting.

**WITHHOLDING PERIODS**

**PRE-EMERGENT: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 WEEKS AFTER APPLICATION**

**POST-EMERGENT: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 14 DAYS AFTER APPLICATION**

**QUEENSLAND ONLY****PRE-EMERGENT APPLICATION RESTRAINTS**

DO NOT apply to crops undersown with legumes.

DO NOT use if another Group B herbicide (ALS Inhibitor) has been used during the current season.

DO NOT apply to paddocks where there is a high risk of weeds resistant to Group B Herbicides.

Apply no more than two Group B Herbicides in any four-year period on the same paddock.

<b>CROP</b>	<b>WEEDS CONTROLLED</b>	<b>RATE/HA</b>
<b>Wheat</b>	African Turnip Weed, Black Bindweed (Climbing Buckwheat), Corn Gromwell, (White Ironweed, Sheepweed), Deadnettle, Hedge Mustard, Indian Hedge Mustard, New Zealand Spinach, Prickly Lettuce, Slender Celery, Smallflower Fumitory, Turnip Weed, Yellow Burrweed (Amsinckia)	30 g
	Annual Ryegrass, Common Peppergrass, Doublegee or Three Cornered Jack (Spiny Emex), London Rocket, Paradoxa Grass, Stagger Weed and suppression of Wild Radish and Wireweed	35 g

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL  
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**



**CRITICAL COMMENTS**

Apply to weed free soil prior to sowing or at sowing and incorporate by the sowing operation using low profile 10cm combine points.

Application should not be made to ridged or excessively cloddy soil.

For best results apply when follow up rain is likely to occur within 7 – 10 days. In conservation tillage situations where weeds and grasses have emerged apply as a tank mixture with the recommended rate of knockdown herbicide prior to sowing.

For Paradoxa Grass control, apply to dry soil before the sowing rain.

DO NOT use on weeds resistant, or thought to be resistant, to Group B Herbicides.

**TO AVOID UNACCEPTABLE RESIDUES**

Do not graze or cut for stock food for 7 weeks after application.

**WITHHOLDING PERIODS**

**PRE-EMERGENT: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 WEEKS AFTER APPLICATION**

**POST-EMERGENT: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 14 DAYS AFTER APPLICATION**

## QUEENSLAND ONLY

### POST-EMERGENT APPLICATION RESTRAINTS

DO NOT apply to crops undersown with legumes.

DO NOT spray when very dry conditions prevail.

DO NOT spray under dry frosty conditions.

DO NOT use if another Group B herbicide (ALS Inhibitor) has been used during the current season.

DO NOT apply to paddocks where there is a high risk of weeds resistant to Group B Herbicides.

Apply no more than two Group B Herbicides in any four-year period on the same paddock.

If RAIN FALLS within 6 hours of application, the effect could be diminished.

CROP	WEEDS CONTROLLED	RATE/HA
<b>Barley, Wheat</b> From 3 leaf to early tillering stage	Turnip Weed	6.5 g + 300 mL terbutryn (500 g/L)
	Denseflower Fumitory, Indian Hedge Mustard, Smallflower Fumitory, Wild Turnip	6.5 g + 300 mL terbutryn (500 g/L) to 10 g + 440 mL terbutryn (500 g/L)
	Black Bindweed (Climbing Buckwheat), London Rocket	10 g + 440 mL terbutryn (500 g/L)
	Wild Radish	10 g + 440 mL terbutryn (500 g/L) to 13 g + 600 mL terbutryn (500 g/L)
	Coreopsis, Corn Gromwell (Sheepweed or White Ironweed)	13 g + 600 mL terbutryn (500 g/L)

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL  
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**CRITICAL COMMENTS**

Spray when weeds are in 2 to 6 leaf stage, except 2 to 4 leaf for Doublegee.

Use the rates towards the lower end of the range (where applicable) when weeds are small and soil conditions are very moist.

Spray only when weeds are actively growing.

Best results are obtained when good soil moisture has been present since planting.

**TO AVOID UNACCEPTABLE RESIDUES**

Do not graze or cut for stock food for 14 days after application.

**WITHHOLDING PERIODS**

**PRE-EMERGENT: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 WEEKS AFTER APPLICATION**

**POST-EMERGENT: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 14 DAYS AFTER APPLICATION**

**VICTORIA ONLY PRE-EMERGENT APPLICATION RESTRAINTS**

DO NOT apply to crops undersown with legumes.

DO NOT use if another Group B herbicide (ALS Inhibitor) has been used during the current season.

DO NOT apply to paddocks where there is a high risk of weeds resistant to Group B Herbicides.

Apply no more than two Group B Herbicides in any four-year period on the same paddock.

<b>CROP</b>	<b>WEEDS CONTROLLED</b>	<b>RATE/HA</b>
<b>Wheat</b> Pre-emergent application	Annual Ryegrass, Burr Medic, Capeweed, Corn Gromwell, (White Ironweed, Sheepweed), Deadnettle, Hedge Mustard, Indian Hedge Mustard, Prickly Lettuce (Whipthistle), Smallflower Fumitory, Volunteer Chickpeas, Faba Beans and Field Peas, Wild Turnip, Wireweed (Hogweed), Yellow Burrweed (Amsinckia) and suppression of Wild Radish and Skeleton Weed	30 to 35 g
	Annual Ryegrass, Corn Gromwell (White Ironweed, Sheepweed), Deadnettle, Hedge Mustard, Indian Hedge, Mustard, Prickly Lettuce, (Whipthistle), Smallflower, Fumitory, Wild Turnip, Wireweed (Hogweed), Ward's Weed.	10 g – 15 g plus 830 mL/ha of trifluralin (480 g/L)

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL  
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

### **CRITICAL COMMENTS**

Apply to bare moist soil prior to sowing or at sowing and incorporate by the sowing operation using low profile 10cm combine points.

Applications should not be made to ridged or excessively cloddy soil. Use rates towards the lower end of the range where broadleaf weeds are the major problem.

Use the higher rate where Capeweed, Volunteer Chickpeas, Faba Beans and Field Peas, Wild Radish and Wireweed are the problem. Chemtura Triasul will provide good control of volunteer grain legumes, however a small proportion of plants may survive and require an overspray to eliminate the potential for grain contamination.

For best results apply to moist soil when follow up rain is likely to occur within 7 – 10 days. Use the lower rate (30 g/ha) on sandy clay loams with a pH greater than 8.5.

For Skeleton Weed a significant degree of control will be achieved on soil types of a predominantly sandy clay loam mixture with a pH greater than 8. Best control is observed where Skeleton Weed germinates in the very early stages of the crop. Surviving plants will be stunted.

**DO NOT** use on weeds resistant, or thought to be resistant, to Group B Herbicides.

This mixture to be only used on alkaline soils where pH is greater than 8 (1:5 Soil: Water suspension method). For best results apply mix to bare moist soil that has a minimum of trash and incorporate to a depth of 5cm just prior to sowing. Incorporation should be made within 4 hours of application. Heavy rainfall (greater than 50 mm) within 7 days of application may affect efficacy especially at the 10 g/ha rate. Late germination of some weeds, e.g. Annual Ryegrass will not be controlled in seasons of above average rainfall.

Use the higher rate where heavy density of weeds is expected.

### **WITHHOLDING PERIODS**

**PRE-EMERGENT: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 WEEKS AFTER APPLICATION**

**POST-EMERGENT: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 14 DAYS AFTER APPLICATION**

**VICTORIA ONLY****POST-EMERGENT APPLICATION RESTRAINTS:**

DO NOT apply to crops undersown with legumes. DO NOT spray when very dry conditions prevail. DO NOT spray under dry frosty conditions. DO NOT use if another Group B herbicide (ALS Inhibitor) has been used during the current season. DO NOT apply to paddocks where there is a high risk of weeds resistant to Group B Herbicides.

Apply no more than two Group B Herbicides in any four-year period on the same paddock.

If RAIN FALLS within 6 hours of application, the effect could be diminished.

<b>CROP</b>	<b>WEEDS CONTROLLED</b>	<b>RATE/HA</b>
<b>Wheat, Oats, Barley</b> Late Post emergent application	Wild Radish	15 g plus crop oil at 1 L per 100 L of spray mixture or surfactant at recommended label rates.
<b>Barley, Oats, Triticale, Wheat</b> From 3 leaf to early tillering stage	Paterson's Curse, Turnip Weed	6.5 g + 300 mL terbutryn (500 g/L)
	Hedge Mustard, Indian Hedge Mustard, Wild Turnip	6.5 g + 300 mL terbutryn (500 g/L) to 10 g + 440 mL terbutryn (500 g/L)
	Crassula, Faba Beans (volunteer), Hyssop	10 g + 440 mL terbutryn (500 g/L)
	Loosestrife, Medic, Prickly Lettuce (Whipthistle)	
	Deadnettle, Field peas (volunteer), Wild Radish	10 g + 440 mL terbutryn (500 g/L) to 13 g + 600 mL terbutryn (500 g/L)
	Volunteer Lupins, Vetch, and Wireweed (less than 3 leaves) suppression.	13 g + 600 mL terbutryn (500 g/L)

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL  
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**CRITICAL COMMENTS**

**LATE POST EMERGENT APPLICATION:** Spray during early flowering of the wild radish. DO NOT apply to the crop during or after anthesis and flowering (Zadoks 60). DO NOT spray on weeds under stress.

**WARNING:** Application to oats may cause some yellowing.

DO NOT use on weeds resistant, or thought to be resistant, to Group B herbicides.

Petroleum based crop oils are recommended as well as non-ionic surfactants (1000 g/L).

Spray when weeds are in 2 to 6 leaf stage (up to 6 node/leaflet for field peas, 10 leaflet for vetch), except 2 to 4 leaf for Doublegee.

Use the rates towards the lower end of the range (where applicable) when weeds are small and soil conditions are very moist.

Spray only when weeds are actively growing.

Spray only after good rain and when top soil is moist.

Best results are obtained when good soil moisture has been present since planting.

**WITHHOLDING PERIODS**

**PRE-EMERGENT: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 WEEKS AFTER APPLICATION**

**POST-EMERGENT: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 14 DAYS AFTER APPLICATION**

## **GENERAL INSTRUCTIONS**

Chemtura Triasul 750 WG Herbicide is a water dispersible granular, herbicide for the pre-plant, incorporated by sowing, control of annual ryegrass, Paradoxa grass and certain broadleaf weeds in wheat, and for post-emergent control of wild radish in wheat, oats and barley.

Crops other than wheat, barley, oats, triticale and cereal rye can be very sensitive to low soil concentrations of Chemtura Triasul 750 WG Herbicide, thus prior to using the product, careful consideration should be given to crop rotation plans.

## **RESISTANT WEEDS WARNING**

### **GROUP B HERBICIDE**

Chemtura Triasul 750 WG Herbicide is a member of the Sulfonylurea group of herbicides. The product has the inhibitors of acetolactate synthase mode of action. For weed resistance management Chemtura Triasul 750 WG Herbicide is a Group B Herbicide.

Some naturally occurring weed biotypes resistant to Chemtura Triasul 750 WG Herbicide and other Group B herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Chemtura Triasul 750 WG Herbicide or any other Group B herbicides.

**DO NOT** rely exclusively on Chemtura Triasul 750 WG Herbicide for weed control. Use as part of an integrated weed management program involving herbicides with other modes of action and non-chemical means of control.

Croplife resistance management strategies are available from your local agricultural chemical supplier. Since the occurrence of resistant weeds is difficult to detect prior to use, Chemtura Australia Pty Ltd accepts no liability for any losses that may result from the failure of Chemtura Triasul 750 WG Herbicide to control resistant weeds.

Advice as to strategies and alternative treatments that can be used should be obtained from your local supplier, consultant, local Department of Agriculture, Primary Industries Department.



## **MIXING**

Chemtura Triasul 750 WG Herbicide is a water dispersible granular herbicide, which mixes readily with water and is applied as a spray.

- Partly fill the spray tank with water
- Start the agitation
- Add the correct amount of product to the spray tank with the agitation system running
- Continue agitation while topping up the tank with water and while spraying.
- Use the spray mix within 24 hours of preparation.

## **APPLICATION**

### **Ground application**

Apply by boom spray, applying 30 to 100 L of water per hectare. Avoid overlapping of boom runs.

### **Aerial application**

Always spray in a crosswind of less than 5 knots. Ensure good spray coverage is obtained. Apply 20 to 40 litres per hectare.

## **SPRAYER CLEANUP**

When the sprayer is being used to spray cereal crops, rinse the sprayer thoroughly with water.

When the sprayer is being used to spray crops other than cereals:

1. Drain tank then rinse tank and spray boom with clean water for at least 10 minutes.
2. Fill the tank with clean water and add to it 300 mL of household chlorine bleach (containing 4% chlorine) per 100 L of water. Rinse hoses and boom and leave in tank for 15 minutes whilst agitating. Drain through nozzle.
3. Repeat step 2 and then rinse thoroughly with clean water to remove all traces of chlorine bleach.
4. Nozzles and filters should be cleaned separately.
5. Dispose of all water used for cleaning.

## COMPATIBILITY

When using a tank mix of Chemtura Triasul 750 WG Herbicide and paraquat/diquat, add Chemtura Triasul, with constant agitation, to approximately half the total volume of water to be used.

- Ensure that the Chemtura Triasul is fully dispersed.
- Add the paraquat/diquat, fill the spray tank to full volume with water and mix thoroughly.
- Apply tank mix immediately, under constant agitation.

Chemtura Triasul 750 WG Herbicide is compatible with S-metolachlor, tri-allate, paraquat/diquat, glyphosate, 2,4-D amine, metsulfuron-methyl, bromoxynil, trifluralin, sodium molybdate, zinc sulphate, manganese sulphate, copper sulphate, chlorpyrifos, permethrin.

## CROP ROTATION GUIDELINES PRE-EMERGENT APPLICATION

**Where the product is applied at the rate of 30–35 g/ha:**

Unless otherwise specified (see table below), wheat, barley, oats, triticale and cereal rye can be planted the following season without restrictions.

For other specified crops Chemtura Triasul 750 WG Herbicide treated area may be replanted after the interval indicated in the table below. These recommendations are made on the assumption that Chemtura Triasul 750 WG Herbicide is applied to a wheat crop that reaches maturity in the season of application.

## CROP ROTATION GUIDELINES (PRE-EMERGENT APPLICATION)

Soil pH (1:5 Soil: Water Suspension Method)	State	Replanting Interval	Minimum Rainfall Requirements between application and sowing the following crop	Crop
6.5 or less	WA, SA, NSW, Vic, Qld only	12 months	300 mm	Field Peas, Linseed, Lucerne, Lupins, Medics**, Subterranean Clover**, Faba Beans, Chickpeas and Canola.
	NSW, Qld only	15 months	700 mm	Sorghum, Maize, Soybean, Cotton, Cowpea and Mung Bean.
		18 months	900 mm	Sunflowers.

<b>Soil pH (1:5 Soil: Water Suspension Method)</b>	<b>State</b>	<b>Replanting Interval</b>	<b>Minimum Rainfall Requirements between application and sowing the following crop</b>	<b>Crop</b>
<b>6.6 to 7.5</b>	NSW, Qld only	12 months	500 mm	Chickpeas and Canola
		15 months	700 mm	Sorghum, Maize, Soybean, Cotton, Cowpea and Mung Bean.
		18 months	900 mm	Sunflowers.
	WA, SA, Vic only	22 months	500 mm	Field Peas, Linseed, Lucerne, Lupins, Medics**, Subterranean Clover**, Faba Beans, Chickpeas, Canola, Sorghum, Maize, Soybean and Cotton
<b>7.6 and 8.5</b>	Vic, SA only	12 Months	250 mm	Barley, Oats, Cereal Rye for grain crops.
			300 mm	Barley, Oats, Cereal Rye for hay crops.
	NSW, Qld only	12 months	500 mm	Chickpeas and Canola.
			18 months	700 mm
	WA, SA, Vic only	24 months	700 mm	Field Peas, Linseed, Lucerne, Lupins, Medics**, Subterranean Clover**, Faba Beans, Chickpeas, Canola, Sorghum, Maize, Soybean and Cotton
<b>8.6 and above</b>	Vic, SA only	12 months	250 mm	Barley, Oats, Cereal Rye for grain crops.
			300 mm	Barley, Oats, Cereal Rye for hay crops.
	WA, SA, NSW, Vic, Qld only	24 months	700 mm	Field Peas, Linseed, Lucerne, Lupins, Medics**, Subterranean Clover**, Faba Beans, Chickpeas, Canola, Sorghum, Maize, Soybean and Cotton

\*\* Includes natural regeneration of subterranean clover and medics.

For all other crops a replanting interval of 24 months has to be observed.

**Where the product is applied at 10–15 g/ha plus 830 mL of trifluralin (480 g/L):**

Where the pH is less than 7.5 the following crops can be replanted from 9 months after application providing 300 mm of rainfall has been recorded; field peas, canola, chickpeas, medics, clover, lucerne, safflower, lupins, cereal rye, barley, oats, wheat, Triticale. Where rainfall is less than 300 mm for this period further advice should be sought from Chemtura.

Where the pH is above 7.5 the following crops can be replanted 9 months after application of Chemtura Triasul–cereal rye, wheat, oats, barley, triticale.

Where the pH is above 7.5 the following crops can be replanted 12 months after application of Chemtura Triasul providing 350 mm of rainfall has been recorded; field peas, canola, chickpeas, medics, clover, lucerne, safflower, lupins. Where less than 350 mm of rain has fallen between application and sowing the crop, further advice should be sought from a Chemtura representative.

**CROP ROTATION GUIDELINES (POST-EMERGENT APPLICATION)**

<b>Soil pH (1:5 Soil: Water Suspension Method)</b>	<b>State</b>	<b>Replanting Interval</b>	<b>Crop</b>
<b>6.5 or less</b>	WA, SA, NSW, Vic only	7 months	Field Peas, Linseed, Lucerne, Lupins, Medics**, Subterranean Clover**, Faba Beans, Chickpeas and Canola.
		14 months	Sorghum, Maize, Soybean and Cotton,
<b>6.6 to 8.0</b>	WA, SA, NSW, Vic only	20 months	Chickpeas, canola, sorghum, maize, soybean, cotton, field peas, linseed, lucerne, medics**, subterranean clover**, faba beans.

\*\* Includes natural regeneration of Subterranean Clover and Medics.

For all other crops a replanting interval of 24 months has to be observed.

These recommendations are made on the assumption that Chemtura Triasul 750 WG Herbicide is applied to a wheat crop that reaches maturity in the season of application. Where Chemtura Triasul 750 WG Herbicide is used on soil types with pH greater than 8 (1:5 soil: water suspension method), further advice should be sought from Chemtura regarding crop rotation guidelines, except for barley, cereal rye, oats, triticale and wheat.

**CROP ROTATION GUIDELINES (tank mixture of Chemtura Triasul 750 WG Herbicide and terbutryn)**

The following crops can be planted after an application of Chemtura Triasul and terbutryn.

In areas where pH is less than 7.5–

**9 months** after an application: barley, cereal rye, chickpeas, faba beans, field peas, lupins, medics, oats, canola, safflower, sub clover, Triticale, wheat.

In areas where pH is greater than 7.5–

**9 months** after an application: barley, cereal rye, oats, Triticale, wheat.

**14 months** after an application: cotton, maize, sorghum, soybeans, and sunflowers. Where residual herbicides are applied following an application, the crop rotational guidelines for these products must be followed.

**22 months** after an application: chickpeas, faba beans, field peas, lupins, medics, canola, safflower, and sub clover.

For all other crops, advice should be sought from Chemtura Australia Pty Ltd.

## **PRECAUTIONS**

Some crop yellowing or crop retardation may occur where a stress factor such as water logging, drought, excessive soil acidity or alkalinity or trace element deficiency, disease—Rhizoctonia, Take All, cereal cyst nematodes or soil insects are present or occur following application.

Special care should be taken with regard to the application of Chemtura Triasul to durum wheats as these may be more sensitive where the above stresses are present. In these situations crop recovery will be rapid provided the stress factors DO NOT continue exerting a negative effect on the crops growth.

Crop retardation may also occur in some instances where considerable late summer/early autumn weed growth occurs. Weeds such as goosefoot *Chenopodium spp* can release herbicidally active compounds into the soil.

## **RE-ENTRY PERIOD**

DO NOT enter treated area without protective clothing until spray has dried.

## **PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS**

DO NOT apply on or near shrubs, trees, lawns or crops other than wheat, oats and barley.

DO NOT drain or flush equipment on or near desirable trees or other plants, where their roots may extend or in situations where by movements of soil, or seepage, absorption of the herbicide may occur.

DO NOT apply under weather conditions or from spraying equipment that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

DO NOT allow spray to drift onto adjacent crops and non-target desirable plants.

DO NOT allow spray to drift onto adjacent fallow land.

## **STORAGE AND DISPOSAL**

### **Keep out of reach of children**

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight.

Triple rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to a recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots in compliance with relevant Local, State or Territory government regulations. Do not burn empty containers or product.

### **FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre. Telephone Australia 13 11 26.

## **MATERIAL SAFETY DATA SHEET**

Additional information is listed in the Material Safety Data Sheet (MSDS) that can be obtained from the supplier.

## **MANUFACTURERS WARRANTY AND EXCLUSION OF LIABILITY**

Chemtura Australia Pty Ltd has no control over storage, handling and manner of use of this product. Where this material is not stored, handled or used correctly and in accordance with directions, no express or implied representations or warranties concerning this product (other than non-excludable statutory warranties) will apply. Chemtura Australia Pty Ltd accepts no responsibility for any loss arising from incorrect storage, handling or use.

In a Transport Emergency  
Dial **000**  
Police or Fire Brigade

APVMA Approval No.: 66746/54314

