

POISON

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

ZIRAGRANZ
FUNGICIDE

ACTIVE CONSTITUENT : 760 g/kg ZIRAM

GROUP	Y	FUNGICIDE
--------------	----------	------------------

**Controls black spot & other fungal diseases of
Celery, Grapes, Pome & Stone Fruits & Roses**

NET CONTENTS: 20kg

Crop Care Australasia Pty Ltd, Unit 15/16 Metroplex Avenue, Murarrie Qld 4172

POISON

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

ZIRAGRANZ
FUNGICIDE

ACTIVE CONSTITUENT : 760 g/kg ZIRAM

GROUP Y FUNGICIDE

**Controls black spot & other fungal diseases of
Celery, Grapes, Pome & Stone Fruits & Roses**

**CONTAINS 15 x 700g MEASURE PACKS
WHICH IT IS ILLEGAL TO SELL SEPARATELY**

NET CONTENTS: 10.5kg

Crop Care Australasia Pty Ltd, Unit 15/16 Metroplex Avenue, Murarrie Qld 4172

DIRECTIONS FOR USE

All States

CROP	DISEASE	RATE	CRITICAL COMMENTS
Grapes	Black spot (<i>Elsinoe ampelina</i>)	150g per 100L	Apply at bud swell, bud burst and 2 to 3 weeks later as a foliar cover spray. If bad weather persists apply at 10 to 14 day intervals.
POME FRUIT Apples	Black spot (<i>Venturia inaequalis</i>)		Apply after early copper sprays. Apply at spur burst pink bud stage and then every 14 days or as required in cover sprays.
Pears	Black spot (<i>Venturia pirina</i>)		Apply after early copper sprays. Apply at spur burst at white bud stage and calyx stage, and then every 14 days or as required in cover sprays.
STONE FRUIT Cherries	Brown rot: Blossom (<i>Monilinia laxa</i>)		Do not apply this product to apricots. Apply at mid full bloom, early petal fall and at shuck fall. Apply cover sprays at 14 day intervals after fruit commences to ripen. Also apply at 21 and 7 days before harvest.
Nectarines Peaches	Brown rot: Fruit (<i>Monilinia fructicola</i>)		
Celery	Septoria Leaf Spot (<i>Septoria apiicola</i>)		Apply as a protective spray every 7 to 10 days from seedling to harvest.
Roses	Black spot (<i>Diplocarpon rosae</i>)		Spray early spring and every 10 to 14 days until end of flowering.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIOD

DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION

GENERAL INSTRUCTIONS

Apply before disease symptoms appear and spray to obtain thorough coverage of stems, leaves, buds and flowers.

MIXING : This product mixes readily with water, no pre-mixing is required. During mixing the spray vat should be no more than 25% full of water. Vigorous agitation of vat contents is required during the addition of the product. Pour product steadily when adding the required amount to the spray vat. DO NOT DUMP all the required product into the vat at once. When the product has been thoroughly dispersed and spray vat is nearly full add 'Agral' Spray Activator and Spreading Agent at a rate of 16mL/100L of spray solution.

COMPATIBILITY : Do not mix with lime sulphur. Do not tank mix this product with fungicides containing copper.

Fungicide Resistance Warning

GROUP	Y	FUNGICIDE
-------	---	-----------

For fungicide resistance management Ziragranz Fungicide is a Group Y fungicide.

Some naturally occurring individual fungi resistant to Ziragranz Fungicide and other Group Y fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by Ziragranz Fungicide and other Group Y fungicides, thus resulting in a reduction in efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use, Crop Care Australasia Pty Ltd accepts no liability for any losses that may result from the failure of Ziragranz Fungicide to control resistant fungi.

PROTECTION OF LIVESTOCK

Low hazard to bees. May be applied on any plants at any time.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to fish. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in dry, cool, well-ventilated area out of direct sunlight. Do not allow product to become wet or overheated in storage.

Shake box empty into spray tank. Single rinse bag before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on-site. Puncture or shred and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Poisonous if inhaled. Harmful if swallowed. Product is corrosive. Attacks eyes and will irritate the skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. Do not inhale dust or spray mist. When preparing spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves and goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 131126. Avoid giving alcohol. If skin contact occurs, remove contaminated clothing and wash skin thoroughly. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

Additional information is listed in the Material Safety Data Sheet.

Conditions of sale

Crop Care Australasia Pty Ltd will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply or use of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Crop Care Australasia Pty Ltd is limited to the replacement of the goods or (at the option of Crop Care Australasia Pty Ltd) the refund of the price paid and is conditional upon a claim being made in writing and where possible sufficient part of the goods to enable proper examination being returned to Crop Care Australasia Pty Ltd within thirty days of delivery.

In a Transport Emergency Dial 000 Police or Fire Brigade	SPECIALIST ADVICE IN EMERGENCY ONLY 1800 033 498 ALL HOURS - AUSTRALIA WIDE
--	---

Agral is a Registered Trade Mark.



Crop Care Australasia Pty Ltd
ACN 061 362 347
Unit 15/16 Metroplex Avenue
Murarrie Qld 4172

Bar codes
to be
inserted
here

APVMA Approval No. : 48755/1102

POISON

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

ZIRAGRANZ

FUNGICIDE

ACTIVE CONSTITUENT : 760 g/kg ZIRAM

GROUP	Y	FUNGICIDE
--------------	----------	------------------

**NOT TO BE SOLD SEPARATELY
BEFORE USE READ ALL DIRECTIONS ON THE OUTER PACK**

NET CONTENTS: 700g

Crop Care Australasia Pty Ltd, Unit 15/16 Metroplex Avenue, Murarrie Qld 4172

APVMA Approval No. 48755/1102